

Box Dimensions of Two Dimensional Cantor like Sets and Special Property of Cantor like Sets Similar to Sets of Positive Measure

Dr. S. M. Padhye¹, Satish B. Khobragade²

Associate Professor, HOD, Department of Mathematics, Shri R.L.T. College of Science, Akola, Maharashtra, India¹

Research Student, Department of Mathematics, Shri R.L.T. College of Science, Akola, Maharashtra, India²

ABSTRACT: In this paper we determine the estimates for upper / lower Box dimension of two dimensional generalized Cantor like sets S . The two dimensional Cantor like sets S have been constructed in [3], for any sequence $\{\epsilon_n\}$ with $0 < \epsilon_n < 1$ with the help of sequence of sets $\{E_n\}$ of subsets of $[0, 1] \times [0, 1]$ such that $E_n \supset E_{n+1}$, $S = \bigcap E_n$ and $m(S) = [\prod_{n=1}^{\infty} (1 - \epsilon_n)]^2$. Further if $\sum \epsilon_n < \infty$, $m(S) \neq 0$ and if $\sum \epsilon_n = \infty$, $m(S) = 0$.

In particular it is proved that the upper and lower Box dimension of the set S satisfy $\underline{dim}_B(S) \geq \lim_{k \rightarrow \infty} \frac{\log 4}{-\log\{[\prod_{n=1}^k \frac{(1-\epsilon_n)}{2}]^{1/k} [\epsilon_{k+1}]^{1/k}\}}$ and $\overline{dim}_B(S) \leq \overline{lim}_{k \rightarrow \infty} \frac{\log 4}{-\log\{[\prod_{n=1}^k \frac{(1-\epsilon_n)}{2}]^{1/k}\}}$. Further we generalize the property of sets of positive measure to Cantor like sets namely "If T is a measurable set of positive measure then $T^* = \{x - y : x \in T, y \in T\} = T - T$ contains an interval $(-\alpha, \alpha)$ for some $\alpha > 0$ ". We in fact prove that if S is Cantor like subset of $[0, 1]$ then $S^* = S - S = [-1, 1]$.

KEYWORDS: Cantor set, Cantor like set, upper Box dimension and lower Box dimension.

I. INTRODUCTION

Definition 1.1 Box Counting Dimension

Let F be any nonempty bounded subset of \mathbb{R}^n and let $N_\delta(F)$ be the smallest number of sets of diameter at most δ which can cover F . The lower and upper box dimensions of F are defined respectively as $\underline{dim}_B(F) = \lim_{\delta \rightarrow 0} \frac{\log N_\delta(F)}{-\log \delta}$ and $\overline{dim}_B(F) = \overline{lim}_{\delta \rightarrow 0} \frac{\log N_\delta(F)}{-\log \delta}$.

If the lower and upper box-counting dimensions of F are equal then the common value is called box-counting dimension of F or box- dimension of F and $dim_B(F) = \lim_{\delta \rightarrow 0} \frac{\log N_\delta(F)}{-\log \delta}$.

Above definition is taken from Kenneth Falconer [2].

II. MAIN RESULTS

A. CONSTRUCTION OF TWO DIMENSIONAL GENERALIZED CANTOR LIKE SET

At first stage we divide unit square E_0 into three columns and three rows consisting of nine rectangles out of which 4 are corner squares each of side $\frac{1-\epsilon_1}{2}$, 4 are rectangles each of size $\epsilon_1 \times (\frac{1-\epsilon_1}{2})$ and 1 centre square is of side ϵ_1 .

Let E_1 be a set obtained by removing middle rectangle from each column and each row. Thus E_1 is the union of 4 corner squares each of side $\frac{1-\epsilon_1}{2}$.

International Journal of Innovative Research in Science, Engineering and Technology

(An ISO 3297: 2007 Certified Organization)

Vol. 5, Issue 12, December 2016

At second stage we divide each remaining square again into three columns and three rows consisting of nine rectangles out of which 4 are corner squares each of side $\frac{(1-\epsilon_1)(1-\epsilon_2)}{4}$, 4 are rectangles each of length $\frac{(1-\epsilon_1)\epsilon_2}{2}$ and breadth $\frac{(1-\epsilon_1)(1-\epsilon_2)}{4}$ and 1 centre square is of side $\frac{(1-\epsilon_1)\epsilon_2}{2}$.

Let E_2 be a set obtained by removing middle rectangle from each column and each row from each remaining square. Thus E_2 is union of 4^2 squares each of side $\frac{(1-\epsilon_1)(1-\epsilon_2)}{4}$.

At third stage we divide each remaining squares again into three columns and three rows consisting of nine rectangles out of which 4 are corner squares each of side $\frac{(1-\epsilon_1)(1-\epsilon_2)(1-\epsilon_3)}{8}$, 4 are rectangles each of length $\frac{(1-\epsilon_1)(1-\epsilon_2)\epsilon_3}{8}$ and breadth $\frac{(1-\epsilon_1)(1-\epsilon_2)(1-\epsilon_3)}{8}$ and 1 centre square is of side $\frac{(1-\epsilon_1)(1-\epsilon_2)\epsilon_3}{4}$.

Let E_3 be a set obtained by removing middle rectangle from each column and each row from each remaining square. Thus E_3 is union of 4^3 squares each of side $\frac{(1-\epsilon_1)(1-\epsilon_2)(1-\epsilon_3)}{8}$.

Continuing in this way at k^{th} stage E_k consists of union of 4^k squares each of side $\prod_{i=1}^k \frac{(1-\epsilon_i)}{2}$.

We define

$$S = \bigcap_{k=0}^{\infty} E_k$$

Where $E_k = \bigcup J_{j,k}$, $j = 1, 2, \dots, 4^k$, each $J_{j,k}$ being a square of side $\prod_{i=1}^k \frac{(1-\epsilon_i)}{2}$.

Above construction is from [3].

We determine of Box dimension of two dimensional generalized Cantor like set constructed in [3].

Theorem 2.2

Lower and upper box-dimensions of two dimensional generalized Cantor like set S satisfy $\underline{dim}_B(S) \geq \lim_{k \rightarrow \infty} \frac{\log 4}{-\log\{\left[\prod_{n=1}^k \frac{(1-\epsilon_n)}{2}\right]^{1/k} [\epsilon_{k+1}]^{1/k}\}}$ and $\overline{dim}_B(S) \leq \lim_{k \rightarrow \infty} \frac{\log 4}{-\log\left[\prod_{n=1}^k \frac{(1-\epsilon_n)}{2}\right]^{1/k}}$.

Proof

We know that in the construction of two dimensional generalized Cantor like set S , at k^{th} stage the set E_k consists of union of 4^k disjoint squares each of diameter

$$\leq \sqrt{2} \left[\prod_{n=1}^k \frac{(1-\epsilon_n)}{2}\right]. \text{ Take } \delta_k = \sqrt{2} \left[\prod_{n=1}^k \frac{(1-\epsilon_n)}{2}\right]. \text{ Then } \delta_k \rightarrow 0 \text{ as } k \rightarrow \infty.$$

\therefore At most 4^k squares of diameter $\leq \delta_k$ are required to cover S .

$$\therefore N_{\delta_k}(S) \leq 4^k$$

By definition 1.1 we get,

$$\begin{aligned} \overline{dim}_B(S) &= \lim_{\delta_k \rightarrow 0} \frac{\log N_{\delta_k}(S)}{-\log \delta_k}. \text{ Since } \delta_k \rightarrow 0 \text{ as } k \rightarrow \infty, \text{ we have} \\ \overline{dim}_B(S) &\leq \lim_{k \rightarrow \infty} \frac{\log 4^k}{-\log\{\sqrt{2} \left[\prod_{n=1}^k \frac{(1-\epsilon_n)}{2}\right]\}} \\ &\leq \lim_{k \rightarrow \infty} \frac{\log 4^k}{-\log\{\sqrt{2} \left[\prod_{n=1}^k \frac{(1-\epsilon_n)}{2}\right]\}^{k/k}} \\ &= \lim_{k \rightarrow \infty} \frac{k \log 4}{-k \log\{\sqrt{2} \left[\prod_{n=1}^k \frac{(1-\epsilon_n)}{2}\right]\}^{1/k}} \\ &= \lim_{k \rightarrow \infty} \frac{\log 4}{-\log\{\sqrt{2} \left[\prod_{n=1}^k \frac{(1-\epsilon_n)}{2}\right]\}^{1/k}} = \lim_{k \rightarrow \infty} \frac{\log 4}{-\log[\sqrt{2}]^{1/k} - \log\left[\prod_{n=1}^k \frac{(1-\epsilon_n)}{2}\right]^{1/k}} \\ \therefore \overline{dim}_B(S) &\leq \lim_{k \rightarrow \infty} \frac{\log 4}{-\log\left[\prod_{n=1}^k \frac{(1-\epsilon_n)}{2}\right]^{1/k}} \end{aligned}$$

Now we show that $\underline{dim}_B(S) \geq \lim_{k \rightarrow \infty} \frac{\log 4}{-\log\{\left[\prod_{n=1}^k \frac{(1-\epsilon_n)}{2}\right]^{1/k} [\epsilon_{k+1}]^{1/k}\}}$, consider a rectangle U of diameter $\leq \delta$, $\delta > 0$

and choose k such that

$$\sqrt{2} \left\{ \left[\prod_{n=1}^k \frac{(1-\epsilon_n)}{2}\right] \epsilon_{k+1} \right\} \leq \delta < \sqrt{2} \left\{ \left[\prod_{n=1}^{k-1} \frac{(1-\epsilon_n)}{2}\right] \epsilon_k \right\}, \text{ then } U \text{ intersects at most one of the basic square of diameter } \leq \sqrt{2} \left[\prod_{n=1}^k \frac{(1-\epsilon_n)}{2}\right] \text{ used in the construction of } S.$$

International Journal of Innovative Research in Science, Engineering and Technology

(An ISO 3297: 2007 Certified Organization)

Vol. 5, Issue 12, December 2016

For if U intersects more than one square of E_k then $|U| \geq$ distance between two disjoint squares of E_k
i.e. $|U| \geq \sqrt{2} \{ [\prod_{n=1}^{k-1} \frac{(1-\epsilon_n)}{2}] \epsilon_k \}$,

i.e. $\delta > \sqrt{2} \{ [\prod_{n=1}^{k-1} \frac{(1-\epsilon_n)}{2}] \epsilon_k \}$,

which is false because $\delta < \sqrt{2} \{ [\prod_{n=1}^{k-1} \frac{(1-\epsilon_n)}{2}] \epsilon_k \}$.

These are 4^k such squares.

\therefore Atleast 4^k squares of diameter $\leq \delta$ are required to cover $E_k \supset S$.
 $\therefore N_\delta(S) \geq 4^k$

$$\text{Now } \delta \geq \sqrt{2} \{ [\prod_{n=1}^k \frac{(1-\epsilon_n)}{2}] \epsilon_{k+1} \}$$

$$\therefore \frac{1}{-\log \delta} = \frac{1}{\log(\frac{1}{\delta})} \geq \frac{1}{-\log(\sqrt{2} \{ [\prod_{n=1}^k \frac{(1-\epsilon_n)}{2}] \epsilon_{k+1} \})}$$

By definition 1.1 we get,

$$\begin{aligned} \underline{\dim}_B(S) &= \lim_{\delta \rightarrow 0} \frac{\log N_\delta(S)}{-\log \delta} \\ &\geq \lim_{k \rightarrow \infty} \frac{\log 4^k}{-\log(\sqrt{2} \{ [\prod_{n=1}^k \frac{(1-\epsilon_n)}{2}] \epsilon_{k+1} \})} \\ &= \lim_{k \rightarrow \infty} \frac{\log 4^k}{-\log \{ \sqrt{2} \{ [\prod_{n=1}^k \frac{(1-\epsilon_n)}{2}] \epsilon_{k+1} \} \}^{k/k}} \\ &= \lim_{k \rightarrow \infty} \frac{k \log 4}{-k \log \{ \sqrt{2} \{ [\prod_{n=1}^k \frac{(1-\epsilon_n)}{2}] \epsilon_{k+1} \} \}^{1/k}} \\ &= \lim_{k \rightarrow \infty} \frac{\log 4}{-\log[\sqrt{2}]^{1/k} - \log \{ [\prod_{n=1}^k \frac{(1-\epsilon_n)}{2}] \epsilon_{k+1} \}^{1/k}} \\ &\therefore \underline{\dim}_B(S) \geq \lim_{k \rightarrow \infty} \frac{\log 4}{-\log \{ [\prod_{n=1}^k \frac{(1-\epsilon_n)}{2}]^{1/k} [\epsilon_{k+1}]^{1/k} \}} \end{aligned}$$

Now we take particular cases corresponding to different values of $\epsilon_n = \epsilon$, a constant, $0 < \epsilon < 1$, in Theorem 2.2 which gives Box dimension of corresponding two dimensional Cantor like set.

Particular Case 2.3

When $\epsilon_n = \frac{1}{5}$, in Theorem 2.2 box dimension of two dimensional Cantor - $\frac{2}{5}$ set is $\frac{\log 4}{\log(\frac{5}{2})}$.

Let $\epsilon_n = \frac{1}{5}$

$$\begin{aligned} \underline{\dim}_B(S) &\geq \lim_{k \rightarrow \infty} \frac{\log 4}{-\log \{ [\prod_{n=1}^k \frac{(1-\epsilon_n)}{2}]^{1/k} [\epsilon_{k+1}]^{1/k} \}} \\ &= \lim_{k \rightarrow \infty} \frac{\log 4}{-\log \{ [\prod_{n=1}^k \frac{(1-\frac{1}{5})}{2}]^{1/k} [\frac{1}{5}]^{1/k} \}} = \lim_{k \rightarrow \infty} \frac{\log 4}{-\log \{ (\frac{2}{5}) [\frac{1}{5}]^{1/k} \}} = \frac{\log 4}{\log(\frac{5}{2})} \end{aligned}$$

$$\therefore \underline{\dim}_B(S) \geq \frac{\log 4}{\log(\frac{5}{2})} \dots\dots\dots(1)$$

$$\begin{aligned} \text{Now, } \overline{\dim}_B(S) &\leq \lim_{k \rightarrow \infty} \frac{\log 4}{-\log \{ [\prod_{n=1}^k \frac{(1-\epsilon_n)}{2}]^{1/k} \}} \\ &= \lim_{k \rightarrow \infty} \frac{\log 4}{-\log \{ [\prod_{n=1}^k \frac{(1-\frac{1}{5})}{2}]^{1/k} \}} = \lim_{k \rightarrow \infty} \frac{\log 4}{-\log \{ [\frac{2}{5}]^k \}^{1/k}} = \frac{\log 4}{\log(\frac{5}{2})} \\ &\therefore \overline{\dim}_B(S) \leq \frac{\log 4}{\log(\frac{5}{2})} \dots\dots\dots(2) \end{aligned}$$

From equations (1) and (2) we get,

$$\underline{\dim}_B(S) = \frac{\log 4}{\log(\frac{5}{2})}$$

which is box dimension of two dimensional Cantor - $\frac{2}{5}$ set.

International Journal of Innovative Research in Science, Engineering and Technology

(An ISO 3297: 2007 Certified Organization)

Vol. 5, Issue 12, December 2016

Particular Case 2.4

When $\epsilon_n = \frac{1}{7}$, in Theorem 2.2 box dimension of two dimensional Cantor - $\frac{3}{7}$ set is $\frac{\log 4}{\log(\frac{7}{3})}$.

Let $\epsilon_n = \frac{1}{7}$,

$$\begin{aligned} \underline{dim}_B(S) &\geq \lim_{k \rightarrow \infty} \frac{\log 4}{-\log\{ [\prod_{n=1}^k \frac{(1-\epsilon_n)}{2}]^{1/k} [\epsilon_{k+1}]^{1/k} \}} \\ &= \lim_{k \rightarrow \infty} \frac{\log 4}{-\log\{ [\prod_{n=1}^k \frac{(1-\frac{1}{7})}{2}]^{1/k} [\frac{1}{7}]^{1/k} \}} = \lim_{k \rightarrow \infty} \frac{\log 4}{-\log\{ (\frac{3}{7}) [\frac{1}{7}]^{1/k} \}} = \frac{\log 4}{\log(\frac{7}{3})}. \\ \therefore \underline{dim}_B(S) &\geq \frac{\log 4}{\log(\frac{7}{3})} \dots\dots\dots(3) \end{aligned}$$

Now

$$\begin{aligned} \overline{dim}_B(S) &\leq \overline{\lim}_{k \rightarrow \infty} \frac{\log 4}{-\log [\prod_{n=1}^k \frac{(1-\epsilon_n)}{2}]^{1/k}} = \overline{\lim}_{k \rightarrow \infty} \frac{\log 4}{-\log [\prod_{n=1}^k \frac{(1-\frac{1}{7})}{2}]^{1/k}} \\ &= \overline{\lim}_{k \rightarrow \infty} \frac{\log 4}{-\log \{ [2]^{1/k} [(\frac{3}{7})^k]^{1/k} \}} = \frac{\log 4}{\log(\frac{7}{3})}. \\ \therefore \overline{dim}_B(S) &\leq \frac{\log 4}{\log(\frac{7}{3})} \dots\dots\dots(4) \end{aligned}$$

From equations (3) and (4) we get,

$$dim_B(S) = \frac{\log 4}{\log(\frac{7}{3})}.$$

which is box dimension of two dimensional Cantor - $\frac{3}{7}$ set.

Particular Case 2.5

When $\epsilon_n = \frac{1}{2}$, in Theorem 2.2 box dimension of two dimensional Cantor - $\frac{1}{4}$ set is 1.

Let $\epsilon_n = \frac{1}{2}$,

$$\begin{aligned} \underline{dim}_B(S) &\geq \lim_{k \rightarrow \infty} \frac{\log 4}{-\log\{ [\prod_{n=1}^k \frac{(1-\epsilon_n)}{2}]^{1/k} [\epsilon_{k+1}]^{1/k} \}} \\ &= \lim_{k \rightarrow \infty} \frac{\log 4}{-\log\{ [\prod_{n=1}^k \frac{(1-\frac{1}{2})}{2}]^{1/k} [\frac{1}{2}]^{1/k} \}} = \lim_{k \rightarrow \infty} \frac{\log 4}{-\log\{ (\frac{1}{4}) [\frac{1}{2}]^{1/k} \}} = \frac{\log 4}{\log 4} = 1 \\ \therefore \underline{dim}_B(S) &\geq 1 \dots\dots\dots(5) \end{aligned}$$

Now

$$\begin{aligned} \overline{dim}_B(S) &\leq \overline{\lim}_{k \rightarrow \infty} \frac{\log 4}{-\log [2 \prod_{n=1}^k \frac{(1-\epsilon_n)}{2}]^{1/k}} \\ &= \overline{\lim}_{k \rightarrow \infty} \frac{\log 4}{-\log [2 \prod_{n=1}^k \frac{(1-\frac{1}{2})}{2}]^{1/k}} = \overline{\lim}_{k \rightarrow \infty} \frac{\log 4}{-\log \{ [2]^{1/k} [(\frac{1}{4})^k]^{1/k} \}} = 1. \\ \therefore \overline{dim}_B(S) &\leq 1 \dots\dots\dots(6) \end{aligned}$$

From equations (5) and (6) we get,

$$dim_B(S) = 1$$

which is box dimension of two dimensional Cantor - $\frac{1}{4}$ set.

Remark 2.6

If $\epsilon_n = \epsilon$, a constant, $0 < \epsilon < 1$, then the estimates for lower and upper box-dimensions of two dimensional generalized Cantor like set S determined in Theorem 2.2 are equal and the common value is $\frac{\log 4}{-\log(\frac{1-\epsilon}{2})}$.

Proof

Let $\epsilon_n = \epsilon$,

$$\underline{dim}_B(S) \geq \lim_{k \rightarrow \infty} \frac{\log 4}{-\log\{ [\prod_{n=1}^k \frac{(1-\epsilon_n)}{2}]^{1/k} [\epsilon_{k+1}]^{1/k} \}}$$

International Journal of Innovative Research in Science, Engineering and Technology

(An ISO 3297: 2007 Certified Organization)

Vol. 5, Issue 12, December 2016

$$= \lim_{k \rightarrow \infty} \frac{\log 4}{-\log\{[\prod_{n=1}^k \frac{(1-\epsilon)}{2}]^{1/k} [\epsilon]^{1/k}\}} = \lim_{k \rightarrow \infty} \frac{\log 4}{-\log\{(\frac{1-\epsilon}{2})^{1/k} [\epsilon]^{1/k}\}} = \frac{\log 4}{-\log(\frac{1-\epsilon}{2})}$$

$$\therefore \underline{\dim}_B(S) \geq \frac{\log 4}{-\log(\frac{1-\epsilon}{2})} \dots\dots\dots(1)$$

Now

$$\overline{\dim}_B(S) \leq \overline{\lim}_{k \rightarrow \infty} \frac{\log 4}{-\log [2 \prod_{n=1}^k \frac{(1-\epsilon n)}{2}]^{1/k}}$$

$$= \overline{\lim}_{k \rightarrow \infty} \frac{\log 4}{-\log [2 \prod_{n=1}^k \frac{(1-\epsilon)}{2}]^{1/k}} = \overline{\lim}_{k \rightarrow \infty} \frac{\log 4}{-\log \{ [2]^{1/k} [(\frac{1-\epsilon}{2})^k]^{1/k} \}} = \frac{\log 4}{-\log(\frac{1-\epsilon}{2})}$$

$$\therefore \overline{\dim}_B(S) \leq \frac{\log 4}{-\log(\frac{1-\epsilon}{2})} \dots\dots\dots(2)$$

From equations (1) and (2) we get,

$$\dim_B(S) = \frac{\log 4}{-\log(\frac{1-\epsilon}{2})}$$

Now we generalize the following property of sets of positive measure, to Cantor like sets. “ If T is a measurable set of positive measure then $T^* = \{ x - y : x \in T, y \in T \} = T - T$ contains an interval $(-\alpha, \alpha)$ for some $\alpha > 0$ ”.

This result is from G. de. Barra [1], p – 43, Example 20.

In fact we prove that if S is a Cantor like set even if their measures are zero they satisfy the property that $S^* = S - S$ contains an interval. The precise statement of the theorem is as follows.

Theorem

If S is one of the Cantor like sets then $S^* = S - S = [-1, 1]$.

This is proved in the form of Lemma and Theorem as follows.

Lemma 2.7

If $x \in [0, 1]$ and x has finite expansion in base 5 of length n then there exists b, c of Cantor - $\frac{1}{5}$ set such that $x = b - c$ and expansion of b in base 5 is not longer than that of a and expansion of c in base 5 is either no longer than that of a or all the digits from n + 1 onwards are 4.

Proof

We prove this result by induction

Let P(n) : If $x \in [0, 1]$ and x has finite expansion in base 5 of length n then there b, c of Cantor - $\frac{1}{5}$ set such that $x = b - c$ and expansion of b in base 5 is not longer than that of a and expansion of c in base 5 is either no longer than that of a or all the digits from n + 1 onwards are 4.

For n = 1, there are just five reals of length 1.

$\therefore x = 0 \cdot 0 = 0 \cdot 0 - 0 \cdot 0$

$$x = 0 \cdot 1 = \frac{1}{5} = \frac{2}{5} - \frac{1}{5} = \frac{2}{5} - \frac{\frac{4}{3^2}}{1 - \frac{1}{5}} = \frac{2}{5} - (\frac{0}{5} + \frac{4}{5^2} + \frac{4}{5^3} + \dots) = 0 \cdot 2 - 0 \cdot 0444\dots$$

$x = 0 \cdot 2 = 0 \cdot 2 - 0 \cdot 0$

$$x = 0 \cdot 3 = \frac{3}{5} = \frac{4}{5} - \frac{1}{5} = \frac{4}{5} - \frac{\frac{4}{3^2}}{1 - \frac{1}{5}} = \frac{4}{5} - (\frac{0}{5} + \frac{4}{5^2} + \frac{4}{5^3} + \dots) = 0 \cdot 4 - 0 \cdot 0444\dots$$

$x = 0 \cdot 4 = 0 \cdot 4 - 0 \cdot 0$

\therefore The result is true for n = 1.

Let the result be true for n-1.

We prove that it is true for n

Let $x = 0 \cdot a_1 a_2 \dots a_n = 0 \cdot a_1 a_2 \dots a_{n-1} \cdot a_n$, where $a_i \in \{0, 1, 2, 3, 4\}$, $i = 1, 2, 3, \dots$

Since the result is true for n-1

\therefore We can write

- (i) $0 \cdot a_1 a_2 \dots a_{n-1} = 0 \cdot b_1 b_2 \dots b_{n-1} - 0 \cdot c_1 c_2 \dots c_{n-1}$ or
- (ii) $0 \cdot a_1 a_2 \dots a_{n-1} = 0 \cdot b'_1 b'_2 \dots b'_{n-1} - 0 \cdot c'_1 c'_2 \dots c'_{n-1} 444\dots$

International Journal of Innovative Research in Science, Engineering and Technology

(An ISO 3297: 2007 Certified Organization)

Vol. 5, Issue 12, December 2016

There are five cases where $a_n = 0, 1, 2, 3, 4$.

Case I $a_n = 0$

$$\begin{aligned} \text{For (i) } x &= 0 \cdot a_1 a_2 \dots a_{n-1} 0 \\ &= 0 \cdot b_1 b_2 \dots b_{n-1} 0 - 0 \cdot c_1 c_2 \dots c_{n-1} 0 \\ &= b - c \end{aligned}$$

Where b, c satisfy required conditions.

$$\begin{aligned} \text{For (ii) } x &= 0 \cdot a_1 a_2 \dots a_{n-1} 0 \\ &= 0 \cdot b'_1 b'_2 \dots b'_{n-1} - 0 \cdot c'_1 c'_2 \dots c'_{n-1} 444\dots \\ &= b' - c' \end{aligned}$$

Where b', c' satisfy required conditions.

Case II For $a_n = 1$

$$\begin{aligned} \text{For (i) } x &= 0 \cdot a_1 a_2 \dots a_{n-1} 1 \\ &= 0 \cdot a_1 a_2 \dots a_{n-1} + 0 \cdot 0 \dots 1^{n^{\text{th}} \text{ place}} \\ &= 0 \cdot b_1 b_2 \dots b_{n-1} - 0 \cdot c_1 c_2 \dots c_{n-1} + 0 \cdot 0 \dots 1^{n^{\text{th}} \text{ place}} \\ &= 0 \cdot b_1 b_2 \dots b_{n-1} - 0 \cdot c_1 c_2 \dots c_{n-1} + 0 \cdot 0 \dots 2^{n^{\text{th}} \text{ place}} - 0 \cdot 0 \dots 4^{(n+1)^{\text{th}} \text{ place}} 444\dots \\ (\because 0 \cdot 0 \dots 1^{n^{\text{th}} \text{ place}} &= \frac{1}{5^n} = \frac{2}{5^n} - \frac{1}{5^n} = \frac{2}{5^n} - \frac{4/5^{n+1}}{1-1/5} = \frac{2}{5^n} - \left(\frac{0}{5^n} + \frac{4}{5^{n+1}} + \frac{4}{5^{n+2}} + \dots \right) \\ &= 0 \cdot 0 \dots 2^{n^{\text{th}} \text{ place}} - 0 \cdot 0 \dots 4^{(n+1)^{\text{th}} \text{ place}} 444\dots) \\ x &= (0 \cdot b_1 b_2 \dots b_{n-1} + 0 \cdot 0 \dots 2^{n^{\text{th}} \text{ place}}) - (0 \cdot c_1 c_2 \dots c_{n-1} + 0 \cdot 0 \dots 4^{(n+1)^{\text{th}} \text{ place}} 444\dots) \\ &= 0 \cdot b_1 b_2 \dots b_{n-1} 2 - 0 \cdot c_1 c_2 \dots c_{n-1} 0444\dots \\ &= b - c \end{aligned}$$

Where b, c satisfy required conditions.

$$\begin{aligned} \text{For (ii) } x &= 0 \cdot a_1 a_2 \dots a_{n-1} 1 \\ &= 0 \cdot a_1 a_2 \dots a_{n-1} + 0 \cdot 0 \dots 1^{n^{\text{th}} \text{ place}} \\ &= 0 \cdot b'_1 b'_2 \dots b'_{n-1} - 0 \cdot c'_1 c'_2 \dots c'_{n-1} 444\dots + 0 \cdot 0 \dots 00^{n^{\text{th}} \text{ place}} 444\dots \\ (\because 0 \cdot 0 \dots 1^{n^{\text{th}} \text{ place}} &= \frac{1}{5^n} = \frac{1}{5^n} = \frac{4/5^{n+1}}{1-1/5} = \frac{0}{5^n} + \frac{4}{5^{n+1}} + \frac{4}{5^{n+2}} + \dots = 0 \cdot 0 \dots 4^{(n+1)^{\text{th}} \text{ place}} 444\dots) \\ x &= 0 \cdot b'_1 b'_2 \dots b'_{n-1} - (0 \cdot c'_1 c'_2 \dots c'_{n-1} 444\dots - 0 \cdot 0 \dots 00^{n^{\text{th}} \text{ place}} 444\dots) \\ &= 0 \cdot b'_1 b'_2 \dots b'_{n-1} - 0 \cdot c'_1 c'_2 \dots c'_{n-1} 400\dots \\ &= 0 \cdot b'_1 b'_2 \dots b'_{n-1} - 0 \cdot c'_1 c'_2 \dots c'_{n-1} 4 \\ x &= b' - c' \end{aligned}$$

Where b', c' satisfy required conditions.

Case III For $a_n = 2$

$$\begin{aligned} \text{For (i) } x &= 0 \cdot a_1 a_2 \dots a_{n-1} 2 \\ &= 0 \cdot a_1 a_2 \dots a_{n-1} + 0 \cdot 0 \dots 2^{n^{\text{th}} \text{ place}} \\ &= 0 \cdot b_1 b_2 \dots b_{n-1} - 0 \cdot c_1 c_2 \dots c_{n-1} + 0 \cdot 0 \dots 02^{n^{\text{th}} \text{ place}} \\ &= 0 \cdot b_1 b_2 \dots b_{n-1} 2 - 0 \cdot c_1 c_2 \dots c_{n-1} \\ &= b - c \end{aligned}$$

Where b, c satisfy required conditions.

$$\begin{aligned} \text{For (ii) } x &= 0 \cdot a_1 a_2 \dots a_{n-1} 2 \\ &= 0 \cdot a_1 a_2 \dots a_{n-1} + 0 \cdot 0 \dots 2^{n^{\text{th}} \text{ place}} \\ &= 0 \cdot b'_1 b'_2 \dots b'_{n-1} - 0 \cdot c'_1 c'_2 \dots c'_{n-1} 444\dots + 0 \cdot 0 \dots 2^{n^{\text{th}} \text{ place}} \\ &= 0 \cdot b'_1 b'_2 \dots b'_{n-1} 2 - 0 \cdot c'_1 c'_2 \dots c'_{n-1} 444\dots \\ x &= b' - c' \end{aligned}$$

Where b', c' satisfy required conditions.

Case IV For $a_n = 3$

$$\text{For (i) } x = 0 \cdot a_1 a_2 \dots a_{n-1} 3$$

International Journal of Innovative Research in Science, Engineering and Technology

(An ISO 3297: 2007 Certified Organization)

Vol. 5, Issue 12, December 2016

$$\begin{aligned}
 &= 0 \cdot a_1 a_2 \dots a_{n-1} + 0 \cdot 0 \dots 3^{nth\ place} \\
 &= 0 \cdot b_1 b_2 \dots b_{n-1} - 0 \cdot c_1 c_2 \dots c_{n-1} + 0 \cdot 0 \dots 3^{nth\ place} \\
 &= 0 \cdot b_1 b_2 \dots b_{n-1} - 0 \cdot c_1 c_2 \dots c_{n-1} + 0 \cdot 0 \dots 4^{nth\ place} - 0 \cdot 0 \dots 4^{(n+1)th\ place} 444 \dots \\
 (\because 0 \cdot 0 \dots 3^{nth\ place} &= \frac{3}{5^n} = \frac{4}{5^n} - \frac{1}{5^n} = \frac{4}{5^n} - \frac{4/5^{n+1}}{1-1/5} = \frac{4}{5^n} - \left(\frac{0}{5^n} + \frac{4}{5^{n+1}} + \frac{4}{5^{n+2}} + \dots \right) \\
 &= 0 \cdot 0 \dots 4^{nth\ place} - 0 \cdot 0 \dots 4^{(n+1)th\ place} 444 \dots)
 \end{aligned}$$

$$\begin{aligned}
 x &= (0 \cdot b_1 b_2 \dots b_{n-1} + 0 \cdot 0 \dots 4^{nth\ place}) - (0 \cdot c_1 c_2 \dots c_{n-1} + 0 \cdot 0 \dots 4^{(n+1)th\ place} 444 \dots) \\
 &= 0 \cdot b_1 b_2 \dots b_{n-1} 4 - 0 \cdot c_1 c_2 \dots c_{n-1} 0444 \dots \\
 &= b - c
 \end{aligned}$$

Where b, c satisfy required conditions.

For (ii) $x = 0 \cdot a_1 a_2 \dots a_{n-1} 3$

$$\begin{aligned}
 &= 0 \cdot a_1 a_2 \dots a_{n-1} + 0 \cdot 0 \dots 3^{nth\ place} \\
 &= 0 \cdot b'_1 b'_2 \dots b'_{n-1} - 0 \cdot c'_1 c'_2 \dots c'_{n-1} 444 \dots + 0 \cdot 0 \dots 2^{nth\ place} + 0 \cdot 0 \dots 00^{nth\ place} 444 \dots \\
 (\because 0 \cdot 0 \dots 3^{nth\ place} &= \frac{3}{5^n} = \frac{2}{5^n} + \frac{1}{5^n} = \frac{2}{5^n} + \frac{4/5^{n+1}}{1-1/5} = \frac{2}{5^n} + \left(\frac{0}{5^n} + \frac{4}{5^{n+1}} + \frac{4}{5^{n+2}} + \dots \right) \\
 &= 0 \cdot 0 \dots 2^{nth\ place} + 0 \cdot 0 \dots 4^{(n+1)th\ place} 444 \dots)
 \end{aligned}$$

$$\begin{aligned}
 x &= 0 \cdot b'_1 b'_2 \dots b'_{n-1} + 0 \cdot 0 \dots 2^{nth\ place} - (0 \cdot c'_1 c'_2 \dots c'_{n-1} 444 \dots - 0 \cdot 0 \dots 00^{nth\ place} 444 \dots) \\
 &= 0 \cdot b'_1 b'_2 \dots b'_{n-1} 2 - 0 \cdot c'_1 c'_2 \dots c'_{n-1} 400 \dots \\
 &= 0 \cdot b'_1 b'_2 \dots b'_{n-1} 2 - 0 \cdot c'_1 c'_2 \dots c'_{n-1} 4 \\
 x &= b' - c'
 \end{aligned}$$

Where b' , c' satisfy required conditions.

Case V For $a_n = 4$

For (i) $x = 0 \cdot a_1 a_2 \dots a_{n-1} 4$

$$\begin{aligned}
 &= 0 \cdot a_1 a_2 \dots a_{n-1} + 0 \cdot 0 \dots 4^{nth\ place} \\
 &= 0 \cdot b_1 b_2 \dots b_{n-1} - 0 \cdot c_1 c_2 \dots c_{n-1} + 0 \cdot 0 \dots 04^{nth\ place} \\
 &= 0 \cdot b_1 b_2 \dots b_{n-1} 4 - 0 \cdot c_1 c_2 \dots c_{n-1} \\
 &= b - c
 \end{aligned}$$

Where b, c satisfy required conditions.

For (ii) $x = 0 \cdot a_1 a_2 \dots a_{n-1} 4$

$$\begin{aligned}
 &= 0 \cdot a_1 a_2 \dots a_{n-1} + 0 \cdot 0 \dots 4^{nth\ place} \\
 &= 0 \cdot b'_1 b'_2 \dots b'_{n-1} - 0 \cdot c'_1 c'_2 \dots c'_{n-1} 444 \dots + 0 \cdot 0 \dots 4^{nth\ place} \\
 &= 0 \cdot b'_1 b'_2 \dots b'_{n-1} 4 - 0 \cdot c'_1 c'_2 \dots c'_{n-1} 444 \dots
 \end{aligned}$$

$$x = b' - c'$$

Where b' , c' satisfy required conditions.

\therefore Result is true for $P(1)$ and $P(n-1) \implies P(n)$

\therefore Result is true for all n.

\therefore For any finite fraction x having expansion in base 5, we can find two elements b and c from Cantor - $\frac{1}{5}$ set such that $x = b - c$.

Theorem 2.8

i) $[0, 1] \subset S - S$

ii) $[-1, 1] = S - S$, where S is Cantor - $\frac{1}{5}$ set i.e. $S = \{ x \in [0, 1] : x \text{ has base 5 expansion containing } 0, 2, 4 \text{ only} \}$

Proof i) We show that $[0, 1] \subset S - S$, where S is Cantor - $\frac{1}{5}$ set. i.e. $S = \{ x \in [0, 1] : x \text{ has base 5 expansion containing } 0, 2, 4 \text{ only} \}$

Let $a \in [0, 1]$ and $a = 0 \cdot a_1 a_2 \dots$ be a base 5 expansion of a.

If $x_k = 0 \cdot a_1 a_2 \dots a_k$, $k \geq 1$. Then $a = \lim_{i \rightarrow \infty} x_i$.

International Journal of Innovative Research in Science, Engineering and Technology

(An ISO 3297: 2007 Certified Organization)

Vol. 5, Issue 12, December 2016

From Lemma 2.7 for $k \geq 1$ there exist members of the Cantor- $\frac{1}{5}$ set b_k and c_k such that $x_k = b_k - c_k$.

Since $\{b_i\} \subset S$, where S is compact there is a convergent subsequence $\{b_{ij}\}$ which converges. Similarly corresponding to $\{b_{ij}\}$ there is a sequence $\{c_{ij}\}$ which has a convergent subsequence $\{c_{ijr}\}$.

Since each subsequence must converge to the same limit as that of its parent sequence, subsequence $\{b_{ijr}\}$ and $\{c_{ijr}\}$ are also convergent. Let their limits be b and c respectively, where $b, c \in S$.

$$\begin{aligned} \text{Also } a &= \lim_{i \rightarrow \infty} x_i \\ &= \lim_{r \rightarrow \infty} x_{ijr} \\ &= \lim_{r \rightarrow \infty} (b_{ijr} - c_{ijr}) \\ &= b - c \\ \therefore a &= b - c, \text{ where } b, c \in S \\ \therefore a &\in S - S \\ \therefore a \in [0, 1] &\implies a \in S - S \\ [0, 1] &\subset S - S \end{aligned}$$

ii) We show that $[-1, 1] = S - S$, where S is Cantor $-\frac{1}{5}$ set.

By part (i) we know that $[0, 1] \subset S - S$ (1)

We show that $[-1, 0] \subset S - S$

$$\begin{aligned} \text{Let } x &\in [-1, 0] \\ \therefore -x &\in [0, 1] \\ \therefore -x &= b - c, \text{ where } b, c \in S && \text{(By part (i))} \\ \therefore x &= c - b, \text{ where } b, c \in S \\ \therefore x &\in S - S \\ \therefore x \in [-1, 0] &\implies x \in S - S \end{aligned}$$

$$[-1, 0] \subset S - S \quad \dots \quad (2)$$

From equations (1) and (2), we get

$$[-1, 1] \subset S - S \quad \dots \quad (3)$$

Now we show that $S - S \subset [-1, 1]$

$$\begin{aligned} \text{Let } x, y &\in S \\ \therefore 0 &\leq x \leq 1 \text{ and } 0 \leq y \leq 1 \\ \therefore 0 &\leq x \leq 1 \text{ and } -1 \leq -y \leq 0 \\ \therefore -1 &\leq x - y \leq 1 \\ \therefore x - y &\in [-1, 1] \\ \therefore S - S &\subset [-1, 1] \quad \dots \quad (4) \end{aligned}$$

From equations (3) and (4), we get
 $[-1, 1] = S - S$, where S is Cantor $-\frac{1}{5}$ set

Similar proofs can be given for Cantor ternary set, Cantor $-\frac{1}{7}$ set, Cantor $-\frac{1}{2n+1}$ set, $n \geq 1$.

Lemma 2.9

If $x \in [0, 1]$ and x has finite ternary expansion of length n then there exists b, c of Cantor ternary set such that $x = b - c$ and expansion of b is not longer than that of a , expansion of c is either no longer than that of a or all the digits from $n + 1$ onwards are 2.

Theorem 2.10

- i) $[0, 1] \subset S - S$
- ii) $[-1, 1] = S - S$, where S is Cantor ternary set.

Lemma 2.11

If $x \in [0, 1]$ and x has finite expansion in base 7 of length n then there exists b, c of Cantor $-\frac{1}{7}$ set such that $x = b - c$ and expansion of b in base 7 is not longer than that of a and expansion of c in base 7 is either no longer than that of a or all the digits from

International Journal of Innovative Research in Science, Engineering and Technology

(An ISO 3297: 2007 Certified Organization)

Vol. 5, Issue 12, December 2016

n + 1 onwards are 6.

Theorem 2.12

- i) $[0, 1] \subset S - S$
- ii) $[-1, 1] = S - S$, where S is Cantor $-\frac{1}{7}$ set.

III. CONCLUSION

We have obtained estimates for upper / lower Box dimension of such two dimensional generalized Cantor like sets S . In particular it is proved that the upper and lower Box dimension of the set S satisfy $\underline{dim}_B(S) \geq \lim_{k \rightarrow \infty} \frac{\log 4}{-\log\{[\prod_{n=1}^k \frac{(1-\epsilon_n)}{2}]^{1/k} [\epsilon_{k+1}]^{1/k}\}}$ and $\overline{dim}_B(S) \leq \overline{\lim}_{k \rightarrow \infty} \frac{\log 4}{-\log\{[\prod_{n=1}^k \frac{(1-\epsilon_n)}{2}]^{1/k}\}}$ and for the particular case $\epsilon_n = \epsilon$, a constant, $0 < \epsilon < 1$ the equality holds namely $\underline{dim}_B(S) = \frac{-\log 4}{\log(\frac{1-\epsilon}{2})}$. Also we observe that the property $S^* = S - S$ contains an interval is also satisfied by Cantor $-\frac{1}{5}$ set, Cantor $-\frac{1}{7}$ set even though they have measure zero.

REFERENCES

- [1] G.de. Barra, "Measure Theory And Integration", New Age International (p) Limited, 2003.
- [2] Kenneth Falconer, "Fractal Geometry : Mathematical Foundations and Applications", John Willey and Sons.1990
- [3] Dr. S.M. Padhye, Satish B. Khobragade " On The Construction Of Two Dimensional Cantor Like Sets And Determination Of Hausdorff Dimension", IJIRSET, Volume 5, Issue 11, pages 19448 - 19456, November 2016.
- [4] J. Shah. Hausdorff dimension and its application. Manuscript. <http://math.uchicago.edu/~may/VIGRE/VIGRE2009/REUPapers/Shah.pdf>