

(A High Impact Factor, Monthly, Peer Reviewed Journal) Visit: <u>www.ijirset.com</u> Vol. 7, Issue 5, May 2018

Clinical Evaluation of *sivappukukilthaila paste* with *panchathuvarpikasayam* in the Management of Wound

Dr. S. R. Pholtan Rajeev¹, Dr. P. Rajavathana²

PG Scholar, Govt. Siddha Medical College, Palayamkottai, Tirunelveli, India¹

Lecturer(Visiting), Trincomalee Campus, EUSL, Sri Lanka²

ABSTRACT :Ayurvedic hospital, vaharai managed the unhealed wounds with wound dressing with herbal medicine successfully. Therefore this research was done to evaluate efficacy of research dressing package as *sivappukukilthila paste* with*panchathuvarpikasayam*. These were prepared by hospital labourers under my guidance. The*panchathuvarpikasayam*used as cleansing the wound area and *sivappukukilthila paste* apply on wounds as healing paste. Research method was Clinical Trial, Research period was 04 months (September to December, 2013), Research samples was Random sample from OPD patients, total Samples was 42 and Research assessment was measurement and visual record of wound healing. According to results and observation; Sample Age defined as; 12 –below and 29 years, 28 -30 years to59years and only 02 –above and 60 years. Sex of the samples were got result was 31 samples were male and 11 samples were Females. Associate symptoms of samples were; no sample complaints only fever, 3% of only lymphnode enlargement, 5% of only Oozing, only pain and lymphnode enlargement and Oozing. 12% of fever with Oozing.19% of fever with lymphnode enlargement and fever with pain.24% of lymphnode enlargement and pain. Character of wound of samples were; Clean-17, Pus-25, granulation tissue formation -13, regular edge -21, irregular edge -21, bad smell -18 and Slough -20. These all characteristic of unhealed wounds were healed within 04 months of routine wound dressing by using of research drug *panchathuvarpikasayam*with*sivappukukilthila paste* significantly.

KEYWORDS: wounds, wound dressing, sivappukukilthila paste andpanchathuvarpikasayam.

I. INTRODUCTION

Wound healing is a complex and dynamic process with the wound environment changing with the changing health status of the individual. The knowledge of the physiology of the normal wound healing trajectory through the phases of hemostasis, inflammation, granulation and maturation provides a framework for an understanding of the basic principles of wound healing. Through this understanding the health care professional can develop the skills required to care for a wound and the body can be assisted in the complex task of tissue repair.

Wounds are occurs various way however healing of wounds are same physiology. In Vaharai area public doing their work in fishing and farming which are create wounds seldom in their life. Some People who has wounds they are treating themselves with surrounding natural herbs. Another people were gone to hospital to treat the wounds. Vaharai has only one Government allopathic hospital and one Ayurveda hospital. These both hospitals were 10km distance gap. I worked as Medical Officer In-charge at Ayurvedichospital, Vaharai. I faced so many patients were came their wound complaints several time Therefore I started wound dressing with herbal medicine successfully.

This research was done in there to evaluate efficacy of research dressing package as *sivappukukilthila paste* with *panchathuvarpikasayam*. These were prepared by hospital labourers under my guidance. The *panchathuvarpikasayam* used as cleansing the wound area and *sivappukukilthila paste* apply on wounds as healing paste.



(A High Impact Factor, Monthly, Peer Reviewed Journal)

Visit: www.ijirset.com

Vol. 7, Issue 5, May 2018

II. MATERIALS AND METHODS

Research method: Clinical Trial, Research area: Ayurvedic Central Dispensary, Vaharai, Batticaloa, Research period: 04 months (September to December, 2013), Research samples: Random sample from OPD patients.Samples: 42, Research assessment: measurement and visual record

Inclusion criteria:

Both sex with any age patient with fresh wound (onset by within 01 month). Incision wound, laceration, injured stab wounds, bed sores, scabies, severe acne, eczema, burn and abscess wound.

Exclusion criteria:

Too old wound with internal pain but external wound was healed. Deep wound with uncontrolled blood sugar Patient with *Diabetes mellitus*. Amputated condition, necrotic condition, too deep erode to bony level, bone fractured wound, deep head injuries, eye involvement, haemo-thorax stab wounds, abdominal stab injuries and severe internal organ involvement wounds. Etc.

Methods: Literature review, Drug preparation

sivappukukilthila paste withpanchathuvarpikasayam

Sample selection, Sample Enrollment to research and initiate data collection, Follow-up with data analyzing, Study with result and observation

- Research Drug preparation: *sivappukukilthila paste* with *panchathuvarpikasayam* method of preparation from siddha pharmacopoeia.
- Prepared drugs (photography attached in appendix)
- Medical consumable items such as cotton wool, gauze, plaster and sterile gloves took from Regional Medical Supply Division (RMSD), Batticaloa.

Symptoms	Assessment variable	Score / Tool	Final assessment	
Fever	Degree Celsius	Thermometer	Normal (36.5°C)	
Lymph node enlargement	Size in mm / Cm	Ruler	Fully disappear	
Oozing	Area mark	Paper cutting area	No oozing	
Pain	Qualitative	Visual Analogue Score (0- 10)	No pain	

Samples were follow-up with routine wound dressing on every 48hrs period.

Table 1: symptoms with wounds

III. LITERATURE REVIEW

 Modern medical view:

 Phase of Healing

 HemostasisImmediate Platelets Capping off conduits

 Inflammation Day 1 - 4 Neutrophils Unskilled laborers to clean up the site

 Proliferation, Granulation

 Contracture Day 4 - 21 Macrophages

 RemodelingDay 21 - 2 years FibrocytesRemodelers

 Basic Principles of Wound Care

 There are three basic principles which underlie wound healing.

 1. Identify and control as best as possible the underlying causes.2. Support patient centered concerns

 3. Optimize local wound care.

 Optimize Local Wound Care

wound healing have become more widely recognized and applied in clinical practice. Some of the advantages include the following:



(A High Impact Factor, Monthly, Peer Reviewed Journal)

Visit: <u>www.ijirset.com</u>

Vol. 7, Issue 5, May 2018

Decreased dehydration and cell death. Increased angiogenesis., Enhanced autolytic debridement, Increased reepithelialization. Bacterial barrier and decreased infection rates. Decreased pain.Decreased costs.

wound healing is a complex physiological process that is dependent on a number of inter-related factors. Wound assessment and treatment should be based on an understanding of normal tissue repair and factors affecting the process. **The process of wound healing**

All tissues in the body are capable of healing by one of two mechanisms: regeneration or repair. Regeneration is the replacement of damaged tissues by identical cells and is more limited than repair.

Local conditions for good wound healing

The provision of a supportive microenvironment at the wound surface is of the utmost importance when trying to maximize a wound's healing potential. Maintaining a controlled set of local conditions that is able to sustain the complex cellular activity occurring in wound healing should be the primary aim of wound management. 1. The vascular response, 2. The inflammatory response, 3. The proliferative stage, 4. The formation of granulation tissue, 5. Wound contraction, 6. Re-epithelialization, 7. The maturation stage.

Ayurvedic view:Vrana (Wound)

Diagnosis: Exogenous Vs. Innate types V-, P-, and K-type wounds

Examination, Locations, Wound discharges, Prognosis, Treatment

Two types of wound have been listed earlier (see the section on enumeration of disorders).

They are:1)InnateThis arises from the doshas within the body.

2)ExogenousThis is due to from external causes. -Ci25#5-6

Etiology of Exogenous Wounds:Blowing, binding, falling down, injury with fangs, teeth or nails, poisonous contact, fire, weapons. Uniqueness of Treatment of Exogenous Wounds:

'These are different from the innate ones in respect of treatment with incantations, anti-poison pastes, etc. etiology and partially symptoms.' -Ci25#8

Etiology of Innate-type Wounds: The dosas get vitiated in the external passages and thus cause wound.

Etiology:	V-type	P-type	K-type	
Symptoms:	it is stiff, hard, it has slow	Thirst, confusion	it has much sliminess, it is heavy	
	exudation, it has excruciating	Fever, sweating,	it is unctuous, wet, it has only mild	
	pain, piercing pain, it has	burning sensation	pain, it is pale of color, it has little	
	throbbing, it has blackish-ness	impurity, tearing	fluid[it is slow to develop and slow	
	_	foul smell, discharge	to recover]	
Treatment:	Saturation	anointing and	anointing and sprinkling with	
	Intake of unction substance	sprinkling with	astringent, pungent, rough and hot	
	Unctuous fomentation	cold, sweet and	drugs	
	and poultices,	bitter drugs intake of	lightening digestive measures	
	ointments and sprinklings	ghee purgation		

Table 2: About the Wound Types

Treatment of [Inflammation/ When Inflammation is a Prodromal Sign]:

Where inflammation is a prodromal sign:

The physician should apply in the beginning of treatment:blood-letting (This will prevent manifestation of the wound). -Ci25#44

Treatment of Patients with Plenty Vs. Little Impurity:

Treatment of Patients with Plenty of Impurity:do evacuation - Ci25#45

Treatment of Patients with Little Impurity:do lightening measures -Ci25#45 (Gabriel Van Loon, 2003)



(A High Impact Factor, Monthly, Peer Reviewed Journal)

Visit: <u>www.ijirset.com</u>

Vol. 7, Issue 5, May 2018

Table 3: Drug ingredients

Drug	Ingredients	Tamil name	Sinhala Name	Part used	
SivappuKukilTh	SivappuKukilTh Sesamumindicum Linn.		Thalathel	Seed oil	
ailam	ailam Rubiacordifolia Linn		Welmathatta	Root	
(mezhukuthailam	Hemidesmusindicus Linn.	Nannari	Iramusu	Root	
)	Bee's wax	Then mezhuku	Mee-iti	Wax	
	ShorearobustaRoxb.	Kungiliyam	Dummala	Resin	
	Water	neer	Wathura	-	
Panchathuvarpi	Ficusrecemosa Linn	Atthi	Attikka	Bark	
(5Astringents)	FicusglobuousWilld	Iththi	Walnuga	Bark	
	Ficusbengalensislinn	Aal	Nuga	Bark	
	Ficusreligiosa Linn	Arasu	Bho	Bark	
	SyzygiumcuminiRoxb.	Naval	Dhang	Bark	

(Ramanathan, 2000)

Drug preparation:

1. Water – 3.2 l, 2. Gingili oil – 0.8kg, 3. Manchisti – 50g, 4. Nannari – 50g, 5. Bee's wax – 50g, 6.Kungiliyam – 50 g

Method:Grind Manchisti and Nannari. Grind powder mixed with water. Heat gingili oil and water mixed powder put into oil when come suitable oil stage, powdered kungiliyam and scraped bee's wax put into oil and filter the oil.Paste took from filtered remains of oil. (Ramanathan, 2000)

Panchathuvarpi:

Atthi – *Ficusrecemosa* Linn, Iththi – *Ficusglobuous*Willd, Aal – *Ficusbengalensis*linn, Arasu – *Ficusreligiosa* Linn, Naval - *Syzygiumcumini*Roxb.(Ramanathan, 2000)

Kasayam (Decoction):

Kudineersaraku – 01 palam-40g, Water – 16 palam -625ml, boiled to 1/6 part. (Ramanathan, 2000)

IV. RESULTS AND OBSERVATIONS



Table 4: Sex of samples



Chart 1: Age of samples



(A High Impact Factor, Monthly, Peer Reviewed Journal)

Visit: <u>www.ijirset.com</u>

Vol. 7, Issue 5, May 2018



Chart 2: Characteristic of sample wounds ; Character of wound of samples were; Clean-17, Pus -25, granulation tissue formation -13, regular edge -21, irregular edge -21, bad smell -18 and Slough-20



Chart 3: Associate symptoms of samples

Associate symptoms of samples were; no sample complaints only fever, 3% of only lymphnode enlargement, 5% of only Oozing, only pain and lymphnode enlargement and Oozing. 12% of fever with Oozing.19% of fever with lymphnode enlargement and fever with pain.24% of lymphnode enlargement and pain.



(A High Impact Factor, Monthly, Peer Reviewed Journal)

Visit: www.ijirset.com

Vol. 7, Issue 5, May 2018

No.	Characters of Wound	Initial assessment	Final assessment
01	Clean	positive	Clean
02	Pus	Negative	No pus
03	Granulation tissue formation	Less	More
04	Regular edge	Positive	Edge less
05	Irregular edge	Negative	Edge less
06	Bad smell	Negative	No odour
07	Slough	Negative	Removed

Table 5: comparison of assessment of wound's characters

Clean and granulation tissue formation were good sign of wound healing these were improve every visit and finally result was healed wound. Pus, regular edge, irregular edge, bad smell, Slough were removed and repair for healing of wounds by each visits with research drug package.

Table 6:Assessment scores: (0-5)

No.	Characters of Wound	0	1	2	3	4	5
01	Clean	fully	marked	moderate	mild	Very mild	unclean
02	Pus	No pus	Very mild	mild	moderate	marked	severe
03	Granulation tissue formation	fully	marked	moderate	mild	Very mild	Not present
04	Regular edge	No edge	Very mild	mild	moderate	marked	defined edge
05	Irregular edge	No edge	Very mild	mild	moderate	marked	undefined edge
06	Bad smell	No	Very mild	mild	moderate	marked	Unpleasant smell
07	Slough	No	Very mild	mild	moderate	marked	severe

Table 7: Score Analysis

No.		Initial	2 nd month	3 rd month	4 th month	Final
01	Clean	5	4	2	1	0
02	Pus	5	3	2	1	0
03	Granulation tissue formation	5	4	2	1	0
04	Regular edge	5	4	2	1	0
05	Irregular edge	5	4	2	1	0
06	Bad smell	5	3	2	1	0
07	Slough	5	3	2	1	0



(A High Impact Factor, Monthly, Peer Reviewed Journal)

Visit: <u>www.ijirset.com</u>

Vol. 7, Issue 5, May 2018

Chart 4: progress of wound characters



V. CONCLUSION

Ayurvedic hospital, vaharai management the wounds with wound dressing with herbal medicine successfully.therefore this research was done in there to evaluate efficacy of research dressing package as *sivappukukilthila paste* with *panchathuvarpikasayam*. These were prepared by hospital labourers under my guidance. The *panchathuvarpikasayam* as cleansing the wound area and *sivappukukilthila paste* apply on wounds as healing paste. Research method was Clinical Trial, Research period was 04 months (September to December, 2013), Research samples was Random sample from OPD patients, total Samples was 42 and Research assessment was measurement and visual record of wound healing.

According to results and observation; Sample Age defined as; 12 –below and 29 years, 28 -30 years to59 years and only 02 –above and 60 years. Sex of the samples were got result was 31 samples were male and 11 samples were Females. Associate symptoms of samples were; no sample complaints only fever, 3% of only lymphnode enlargement, 5% of only Oozing, only pain and lymphnode enlargement and Oozing. 12% of fever with Oozing.19% of fever with lymphnode enlargement and fever with pain.24% of lymphnode enlargement and pain.Character of wound of samples were; Clean-17, Pus -25, granulation tissue formation -13, regular edge -21, irregular edge -21, bad smell -18 and Slough -20. These all characteristic of unhealed wounds were healed within 04 months of routine wound dressing by using of research drug *panchathuvarpikasayam* with *sivappukukilthila paste* significantly.

Because of antimicrobial activity of *panchathuvarpikasayam* was act as good cleanser of wound debris. *sivappukukilthila paste* acted good wound healer by promote regeneration or repair of wound tissues by reduced Inflammation, reorganized Proliferation, Granulation, Contracture and Remodelling naturally.

REFERENCES

- 1. Baxter CR. Immunologic reactions in chronic wounds. American Journal of Surgery 1994;167(1A Suppl):12S-14S
- 2. Baxter, C.R. Immunologic reactions in chronic wounds. Am J Surgery 1994; 167: S: 12S-14S.
- 3. Bennett, G., Moody, M. Wound Care for Health Professionals. London: Chapman and Hall, 1995.
- 4. Brown, G.L. Acceleration of tensile strength of incisions treated with EGF and TGF. Annals of Surgery 1988; 208: 788-794.
- 5. Clark, R.A.F. Overview and general considerations of wound repair. In Clarke, R.A.F., Henson, P.M. (eds). *The Molecular and Cellular Biology of Wound Repair*. New York:
- 6. David Keast MSc, MD, CCFP, Heather Orsted RN, BN, ET, THE BASIC PRINCIPLES OF WOUND HEALING, 2010
- 7. Diegelmann, R. et al. The role of macrophages in wound repair: a review. *Plastic Reconstructive Surgery* 1991; 68: 107-113.
- 8. Eisenbeiss, W., Peter, P.W., Bakhtiari, C. et al. Hypertrophic scars and keloids. J Wound Care 1998; 7: 5, 255-257.
- 9. Fincham-Gee, C. Nutrition and wound healing. *Nursing* 1990; 4: 18, 26-28.
- 10. Flanagan, M. Wound Management. Edinburgh: Churchill Livingstone, 1997.



(A High Impact Factor, Monthly, Peer Reviewed Journal)

Visit: www.ijirset.com

Vol. 7, Issue 5, May 2018

- 11. Flanagan. M, MA, BSc, DipN, Cert Ed, ONC, RGN, Physiology; Wound healing-REVIEW, JOURNAL OF WOUND CARE JUNE, VOL 9, NO 6, 2000, 299
- 12. Gabriel Van Loon, CharakaSamhita, Handbook on Ayurveda, Volume II, 2003
- 13. Garrett, B. Re-epithelialisation. J Wound Care 1998; 7: 7, 358-359.
- 14. GT, Sibbald RG. (eds) Wayne, PA: Chronic Wound Care: A Clinical Source Book for Healthcare Professionals, Third Edition, Health Management Publications, 2001,pp 7-17.
- 15. Haimowitz JE, Margolis DJ, Wayne, PA: Moist wound healing, in Krasner D, Kane D (eds): Chronic Wound Care: A Clinical Source Book for Healthcare Professionals, Health Management Publications, 1997, 49-55
- 16. Harding, K., Cutting, K. Criteria for identifying wound infection. J Wound Care 1994; 3: 4, 198-201.
- 17. Hutchinson JJ, McGuckin M. Occlusive dressings: A microbiologic and clinical review. Am J Infect Control 1990;18:257-268
- 18. Hutchinson, J.J. Prevalence of wound infection under occlusive dressings: a collective survey of reported research. Wounds 1989; 1: 123-133.
- 19. Kane D, Krasner D, Rodeheaver: Chronic wound healing and chronic wound management, 2000.
- 20. Katz, M.H., Alvarez, A.F., Kirsner, R.S. et al. Human wound fluid from acute wounds stimulates fibroblasts and endothelial cell growth. J Am Academic Derma 1991; 25: 1054-1058.
- 21. Kerstein MD: Introduction: moist wound healing. American Journal of Surgery 1994; 167(1A Suppl): 1S-6S
- 22. Kerstein MD: The scientific basis of healing. Adv Wound Care 1997; 10(3):30-36
- 23. Kiecolt-Glaser, J.K., Marucha, P.T., Malarkey, W.B. et al. Slowing of wound healing by psychological stress. Lancet 1995; 346: 1194-1196.
- 24. Knighton DR, Silver JA, Hunt TK. Regulation of wound-healing angiogenesis: effect of oxygen gradients and inspired oxygen concentration. Surgery 1981;90:262-270
- 25. Knighton, D. et al. Regulation of wound healing angiogenesis: effect of oxygen gradients and inspired oxygen concentration. *Surgery* 1981; 90: 262-270.
- 26. Krasnor, D. Chronic Wound Care: A clinical sourcebook for healthcare professionals (2nd edn). Wayne, Pa: Health Management Publications, 1996.
- 27. Lazarus G, Cooper D, Knighton D, Margolis D, Pecoraro R, Rodeheaver G, Robson. Definitions and guidelines for assessment of wounds and evaluation of healing. *Archives of Dermatology* 1994;130:489-493
- 28. Lock, P. The Effects of Temperature on Mitotic Activity at the Edge of Experimental Wounds. Lock Research Laboratories Paper. Kent: Lock Laboratories, 1979.
- 29. MacLeod J (ed): Davidson's Principles and Practice of Medicine, Thirteenth Edition. Edinburgh UK, 1981, pp 590-592