



e-ISSN: 2319-8753 | p-ISSN: 2347-6710

# IJIRSET

International Journal of Innovative Research in  
**SCIENCE | ENGINEERING | TECHNOLOGY**

# INTERNATIONAL JOURNAL OF INNOVATIVE RESEARCH

IN SCIENCE | ENGINEERING | TECHNOLOGY

Volume 10, Issue 12, December 2021

**ISSN** INTERNATIONAL  
STANDARD  
SERIAL  
NUMBER  
INDIA

Impact Factor: 7.569

 9940 572 462

 6381 907 438

 [ijirset@gmail.com](mailto:ijirset@gmail.com)

 [www.ijirset.com](http://www.ijirset.com)



# Extraction of Oleic Acid from Neem Leaves

K. B. Zare, R.V. Thakur, P. N. Vardhe, V. S. Yadav, S.S.Datir

P.G. Student, Department of Chemical Engineering, DYPIEMR, Akurdi, Pune, India

Assistant Professor, Department of Chemical Engineering, DYPIEMR, Akurdi, Pune, India

**ABSTRACT:** Herbs play a vital function in human's everyday life. They are utilized in especially 3 sectors food, pharmaceutical and nutraceutical in industries. Today we worry to have a fruit or vegetable sold from marketplace because of the involvement of the chemical insecticides with inside the farming. Thus there arises a want of natural insecticides with the intention to assist each the farmer in addition to patron at the identical hand. There numerous herbs with lively additives which might also additionally save you the pest effect at the plants which must be extracted from respective plant life and needs to be jumbled in a set proportion. The specific uncooked substances are kokum, cardamom, neem (*Azadirachta indica*), fennel, etc. Thus our predominant goal is to extract out a natural pesticide from natural extracts. Raw substances are decided on primarily based totally at the anti-microbial, anti-inflammatory, anti-tumour properties. Oleic acid turned into discovered to be 25-54% neem oils fatty acid content material and has belongings to provide soap. By the use of UV spectrometer, we discovered the attention of Oleic acid with inside the Neem leaves. From the evaluations and experimentation, the attention of oleic acid will increase with boom in temperature in addition to RPM.

**KEYWORDS:** Extraction, Oleic Acid, Neem Leaves, Soxhlet.

## I. INTRODUCTION

Herbs: are any plants used for flavouring, food, medicine, or fragrance from past days. cookery use generally distinguishes herbs as relating the foliate inexperienced components of a spice, a product from another a part of the plant, as well as seeds, berries, bark, roots and fruits. piece of writing or Indian system of medicines is one such conceive to enhance quality of life. Ayurveda herbs or Indian herbs are used since 1000' of years to supply seasoned remedies. these days it's a well-established incontrovertible fact that herbal remedies are additional appropriate to anatomy than isolated chemical medicines [1]. Aspirin, atropine, artemisinin, colchicine, digoxin, ephedrine, morphine, physostigmine, pilocarpine, quinine, quinidine, reserpine, taxol, neuromuscular blocking agent and Oncovin are many vital samples of what meditative plants have given America within the past. nowadays in several countries modern medication has displaced plants with many manmade merchandise however virtually 30% of pharmaceutical preparations are still obtained directly or indirectly from plants. the trendy era has seen some decline in use of medicinal plants and their extracts as therapeutic agent. The strategy of analytic the active principles from the medicinal plants and producing a pharmaceutical preparation then became common [2]. Plants have the power to synthesize a wide variety of chemical compounds that are wont to perform vital biological functions, and to defend against attack from predators such as insects, fungi and phytophagous mammals. several of those phytochemicals have useful effects on long-run health once consumed by humans, and may be used to effectively treat human diseases. At least 12,000 such compounds are isolated thus far; a number calculable to be under 10% of the total. Chemical compounds in plants mediate their effects on the anatomy through processes a twin of those already well understood for the chemical compounds in standard medication; therefore seasoned medicines don't take issue greatly from conventional drugs in terms of however they work. This allows herbal medicines to be as effective as conventional medicines, however conjointly offers them an equivalent potential to cause harmful facet effects [11]. the employment of plants as medicines predates written human history. Ethno-botany (the study of ancient human uses of plants) is recognized as a good way to discover future medications. In 2001, researchers known 122 compounds utilized in trendy medicine that were derived from "ethno-medical" plant sources; 80% of those have had associate ethno-medical use identical or involving the present use of the active components of the plant. Several of the prescribed drugs presently obtainable to physicians have a protracted history of use as seasoned remedies, as well as aspirin, digitalis, quinine, and-opium [12]. Herbs have a spread of uses including cookery, meditative, and in some cases religious usage. General usage of the term "herb" differs between culinary herbs and medicinal herbs. In medicinal or religious use any of the components of the plant could be thought of "herbs", as well as leaves, roots, flowers, seeds, resin, root bark, inner bark (and cambium), berries and typically the natural covering or alternative parts of the plant [13]. Culinary herbs: Culinary herbs are prominent from veggies in that, like spices, they may be utilized in small quantities and offer taste instead of substance to food. Herbs may be perennials which includes thyme or lavender, biennials which includes parsley, or


annuals like basil. Perennial herbs can be shrubs such as rosemary, *Rosmarinus officinalis*, or timber such as bay laurel, *Laurus nobilis* – this contrasts with botanical herbs, which via way of means of definition can't be woody plant life. Some plant life are used as each herbs and spices, which includes dill weed and dill seed or coriander leaves and seeds. Also, there are a few herbs which includes the ones with inside the mint own circle of relatives which are used for each culinary and medicinal purposes [13]. Medicinal herbs: Some flowers include phytochemicals which have consequences at the body. There can be a few consequences whilst ate up with inside the small tiers that typify culinary "spicing", and a few herbs are poisonous in large quantities. For instance, a few forms of natural extract, inclusive of the extract of St. John's-wort (*Hypericum perforatum*) or of kava (*Piper methysticum*) may be used for clinical functions to alleviate despair and stress. However, massive quantities of those herbs can also additionally cause poisonous overload that could contain complications, a number of a extreme nature, and need to be used with caution.

## II. METHODOLOGY

Extraction is one widely used separation technique. During this the component is separated employing a suitable solvent. Extraction are often broadly classified into two categories:

1. Liquid- Liquid Extraction
2. Solid- Liquid Extraction

Liquid- Liquid Extractions: When a liquid mixture isn't easily separable by distillation, liquid extraction is used. This system is usually employed when:



Liquid- Liquid Extraction

- i. The components are having very close boiling points.
- ii. They can't withstand the temperature of distillation.
- iii. Materials are heat sensitive.

### 3.2 Solid- Liquid Extraction:

Leaching may be a separation technique during which the solid mixture is contacted with a liquid solvent for removal of 1 or more soluble constituents of the solid mixture.

The process of leaching consists of two steps:

- i. Contacting the solid with the selective solvent so as to dissolve the soluble solute within the solvent.
- ii. Separation of insoluble phases i.e. separation of the liquid phase physically from the solid phase physically settling, filtration etc.

The various solvents utilized in the method of leaching are- hexane, iso-octane, toluene, chloroform, dichloromethane, tetrahydrofuran, ether, ester, acetone, acetonitrile, isopropanol, methanol, water, etc.

Factors affecting rate of leaching are:

1. Particle size: - Smaller the particle size, greater the interfacial area between the solid and liquid and greater the speed of leaching.

1. Solvent: - The solvent used should be selective and will have low viscosity for free of charge circulation.

2. Temperature: - The solubility of soluble material increases with increase in temperature, so heat favors the leaching operation.


3. Agitation: - Agitation increases the eddy diffusion and thus increases the speed of leaching. Agitation also

prevents the settling of solids, keeps them in suspension. The algorithm automatically generates mask image without user interaction that contains only text regions to be in painted.

### III. EXPERIMENTAL RESULTS

Conventional Method:

1. Soxhlet Extraction: In a conventional Soxhlet system, material is placed during a thimble-holder, and crammed with condensed fresh solvent from a distillation flask. When the liquid reaches the overflow level, a siphon aspirates the answer of the thimble-holder and unloads it back to the distillation flask, carrying extracted solutes into the majority liquid. Within the solvent flask, solute is separated from the solvent using distillation. Solute islet within the flask and fresh solvent passes back to the plant solid bed. The operation is repeated until complete extraction is achieved.



Schematic of Soxhlet extraction


Material used: Then they had been washed properly with distilled water and dried below solar rays (at 28°C for five-6 days)/ tray drier and grinded it then the grinded powder became taken for similarly experimentation. b) Apparatus: Glass reactor of capability 500 ml, clear out out paper, beaker, conical flask, stirring rod, UV- spectrophotometer unit, micropipette, volumetric flasks, funnel, weighing pan, pattern bottles, water tubtub, batch setup, soxhlet setup. c) Batch extraction: 20 gm of shed dry and overwhelmed leaves (particle length 300 - seventy five  $\mu\text{m}$ ) became subjected to completely baffled a hundred and fifty ml stirred borosilicate glass vessel at the side of ethanol (2 hundred ml) as solvent. A 4 blade turbine kind agitator became used for stirring at special rpm to make sure all debris continue to be in suspended condition. Extraction of oleic acids became achieved at special temperature for 2.five hrs. The schematic experimental setup is as proven in Figure. The samples had been amassed in an c program languageperiod of 15 minutes. The amassed samples had been filtered thru a everyday clear out out paper in most cases after which filtered thru what man clear out paper and analysed in UV spectrophotometer for willpower of Oleic acid contents in mixtures. d) Solet extraction: It became run on the way to discover max awareness that may be extracted out with inside the special time intervals. For this purpose, the we had positioned 200ml solvent in conical flask and 20gm neem leaves powder with inside the thimble and run for 3hrs and 5hrs. Conc. of oleic acid became checked in UV spectrometer. Four.1.1 Procedure: a. Pre-Processing: • Fresh Neem leaves in which amassed from the university premises. From this the small twigs, huge particle, horrific leaves and small braches of leaves in which removed. • They had been washed properly in order to do away with the dirt debris. • Then stored in tray drier for 10 min in order to do away with moisture content material after which stored below solar for five-6 days. • Then the leaves are grinded/ powdered in mixer properly. b. Process: • Water tub tub is switched ON. • 20gms of powdered neem leaves (uncooked material) in which measured the use of weighing balance. • Weighed powder at the side of 200ml of ethanol (Solvent) became poured into the extractor. • Then the baffle became inserted in to the extractor and it became nicely tightened. • The agitator became switched ON and the RPM became set (as 2 hundred, 400 & 600). • Samples are amassed from extractor at an c program language period of 15 minutes up-until 2:30 hrs. • Then the setup became switched OFF completely. • Each of the pattern bottles are classified and amassed in a beaker containing water is both moved for refrigeration (on the way to save you infection or save you microbial activity) and without delay taken for UV



evaluation in analytical lab. • Before evaluation the samples are nicely filtered with the aid of using in most cases everyday clear out paper after which with what man clear out paper. • The last pattern in extractor is ship to Solet for solvent healing and to get neem oil separately.

#### IV. RESULT AND DISCUSSION

In the experimental paintings we studied exceptional temperature (300C, 400C, 500C & 600C) & exceptional rpm's (200rpm, 400rpm & 600rpm). Sample became pipette out after each 15min for UV evaluation while the batch extraction started. Results are acquired through UV that indicates the attention of Oleic acid with inside the extract segment at that specific time. By converting parameters pace of agitator and temperature with inside the reactor we've acquired exceptional values at exceptional time.



Concentration of oleic acid obtained from Soxhlet Extraction

#### V. CONCLUSION

The new integrated approach consisting of a screening of extraction conditions and an in vitro measuring of functional activities alongside an exhaustive chemical characterization will provide us with a replacement tool to get new bioactive compounds and to assist the further design of processes to get such products within the greenest, sustainable, and efficient way, complying with the principles of green chemistry and green engineering. During this chapter we discuss and present some selected applications of the extraction of target compounds with, among others, antioxidant, antimicrobial, and anti-proliferative activities from different sustainable marine sources. We have shown many possibilities at laboratory scale, which are the idea of the knowledge, which are often used because the initiative for a much bigger production scale. Undoubtedly, the replacement of environmentally burdensome solvents like acetonitrile, methanol, dichloromethane, and toluene, long utilized in such extraction processes, for more green solvents like CO<sub>2</sub>, ethanol, and water is one among the goals and one among the proposals. Some technologies are mature enough to be used at large scale; others require more study and development but, in any case, it's advisable that new steps be taken to assist build a more rational use of our natural resources. The likelihood, mentioned during this chapter as a future trend, to create new platforms able, during a sustainable way, to run integrated processes including pre-treatments, extractions, reactions, and transformations during a more integrated way is one among our main goals and might help all folks to create a far better future. We may conclude that extraction of chemicals from neem is important for reducing numerous diseases like cancer, skin, digestive, etc. It's therefore expected that neem oil thus extracted could also be useful for the prevention and reduction of diseases and can provide an efficient pesticide for eco-friendly environment.

#### REFERENCES

- [1] Usman J.G., Okonkwo P.C., Shehu M.S. "Investigation into The Usage of Solvent for Extracting Neem Oil from Neem Seed for Industrial Application Academic", "Journal of Interdisciplinary Studies Mcser Publishing, Rome-Italy", Vol 3 No 5, Issn 2281-3993, July 2014
- [2] R. Carl Millican and J. B. Brown, "The Isolation and Properties of Some Naturally Occurring Octadecenoic (Oleic) Acids", "J. Biol. Chem.", 1944, 154:437-450.
- [3] Bushra Sultana 1, Farooq Anwar 1,\* and Muhammad Ashraf, "Effect of Extraction Solvent/ Technique on the Antioxidant Activity of Selected Medicinal Plant Extracts", "Molecules" 2009, 14, 2167-2180.



- [4] Narsing Rao, G., \*Prabhakara Rao, P. G. and Satyanarayana, AShahidi, "Chemical, fatty acid, volatile oil composition and antioxidant activity of shade dried neem (AzadirachtaIndicalL.) flower powder", "International Food Research Journal", 21(2): 807-813(2014).
- [5] Dr. V. V. L. N. Prasad1, Dr. Prakash V Diwan, "Studies on extraction and HPLC Analysis of Azadirachtin from Kernels of Neem Seeds", Journal of Advanced Pharmacy Education & Research Jan-Mar 2013 Vol 3 Issue 1, 2
- [6] Sutar.R.L.,Mane.S.P,Ghosh.J.S,'International FoodResearchJournal',19(3),1207-1210,2012.
- [7]S.N.Pravinkumar,D.G.BhadreGowda,K.Mantelingu,K.S.Rangappa,'Journalofpharmacy Research', 7, 103-106, 2013.
- [8]Chetan A. Nayak, Navin K. Rastogi, 'Separation and purification technology', 71, 144-151, 2010.
- [9] VipulUpadhyay, NeeruShrama, Himanshu Mohan Joshi, Ajay Singh, Manoj Mishra, Sanjeev Tripathi, Brijpal Singh, 'International Journal of Pharmacy and Integrated Life Sciences', Vol: 1(6) May2013.
- [10] G. K.Jayaprakasha, K.K., 'Sakariah Journal of Pharmaceutical and Biomedical Analysis', Vol.28, 379-384, 2002.
- [11] Y.C. Tripathi, "Research towards Productivity Enhancement and Value Addition", "Multifaceted Applications of Neem", Vol. 1, issue 9, October 2000, 220-237.
- [12] Skoog D. A., West D. M., Holler F. J., 'Fundamentals of Analytical Chemistry', 7th Edition, Thomson Learning, Inc, USA, 1996.
- [13] Sheen R.T. and Kahler H. L., 'Effects of Ions on Mohr Method for Chloride Determination', Ind.Eng. Chem. Anal. Ed.,10(11), 628-629,1938.
- [14]Kraemer E. O. and Stamm A. J., 'Mohr's Method for the Determination of Silver and Halogens in other than Neutral Solutions', J. Am. Chem. Soc.; 46(12); 2707 - 2709, 1924.
- [15] T. Farhat, R. J. Iannotti, and B. G. Simons-Morton, 'AmericanJournal of Preventive Medicine', vol. 38, no. 3, pp. 258-267, 2010.
- [16] C. Roongpisuthipong, R. Kantawan, and W. Roongpisuthipong, 'AsiaPacific Journal of Clinical Nutrition, vol. 16, no. 1, pp. 25-29, 2007.
- [17] B. W. Downs, M. Bagchi, G. V. Subbaraju, M. A. Shara, H. G. Preuss, and D. Bagchi, 'Bioefficacy of a novel calciumpotassium salt of (-)-hydroxycitric acid', Mutation Research,vol. 579, no. 1-2, pp. 149-162, 2005.
- [18] H. G. Preuss, D. Bagchi, M. Bagchi, C. V. S. Rao, S. Satyanarayana, and D. K. Dey, 'Efficacy of a novel, natural extract of (-)-hydroxycitric acid (HCA-SX) and a combination of HCA-SX, niacin-bound chromium and Gymnemasylvestre extract in weight management in human volunteers: a pilot study', Nutrition Research, vol. 24, no. 1, pp. 45-58, 2004.
- [19] E. Toromanyan, G. Aslanyan, E. Amroyan, E. Gabrielyan, and A. Panossian, 'Efficacy of Slim339\_ in reducing body weight of overweight and obese human subjects', PhytotherapyResearch, vol. 21, no. 12, pp. 1177-1181, 2007.
- [20] T. Yamada, H. Hida, and Y. Yamada, 'Chemistry, physiological properties, and microbial production of hydroxycitric acid', AppliedMicrobiology and Biotechnology, vol. 75, no. 5, pp. 977- 982, 2007.



**Impact Factor:  
7.569**



# INTERNATIONAL JOURNAL OF INNOVATIVE RESEARCH

IN SCIENCE | ENGINEERING | TECHNOLOGY

 9940 572 462  6381 907 438  [ijirset@gmail.com](mailto:ijirset@gmail.com)



[www.ijirset.com](http://www.ijirset.com)

Scan to save the contact details