



e-ISSN: 2319-8753 | p-ISSN: 2320-6710

IJIRSET

International Journal of Innovative Research in
SCIENCE | ENGINEERING | TECHNOLOGY

INTERNATIONAL JOURNAL OF INNOVATIVE RESEARCH

IN SCIENCE | ENGINEERING | TECHNOLOGY

Volume 10, Issue 2, February 2021

ISSN INTERNATIONAL
STANDARD
SERIAL
NUMBER
INDIA

Impact Factor: 7.512

9940 572 462

6381 907 438

ijirset@gmail.com

www.ijirset.com

Pesticides and Their Residual Effects in Soil and Water of District Ganganagar, Rajasthan

Iqbal Singh

Associate Professor in Zoology, Dr. Bhim Rao Ambedkar Govt. College, Sri Ganganagar, Rajasthan, India

ABSTRACT: Pesticides are substances that are meant to control pests. [1] This includes herbicide, insecticide, nematicide, molluscicide, piscicide, avicide, rodenticide, bactericide, insect repellent, animal repellent, microbicide, fungicide, and lampricide. [2][3] The most common of these are herbicides, which account for approximately 50% of all pesticide use globally. [4] Most pesticides are intended to serve as plant protection products (also known as crop protection products), which in general, protect plants from weeds, fungi, or insects. As an example, the fungus *Alternaria solani* is used to combat the aquatic weed *Salvinia*.

In general, a pesticide is a chemical (such as carbamate) or biological agent (such as a virus, bacterium, or fungus) that deters, incapacitates, kills, or otherwise discourages pests. Target pests can include insects, plant pathogens, weeds, molluscs, birds, mammals, fish, nematodes (roundworms), and microbes that destroy property, cause nuisance, or spread disease, or are disease vectors. Along with these benefits, pesticides also have drawbacks, such as potential toxicity to humans and other species.

KEYWORDS- Pesticides, Effects, Soil, Water, Ganganagar, Rajasthan

I. INTRODUCTION

The Food and Agriculture Organization (FAO) has defined pesticide as:

any substance or mixture of substances intended for preventing, destroying, or controlling any pest, including vectors of human or animal disease, unwanted species of plants or animals, causing harm during or otherwise interfering with the production, processing, storage, transport, or marketing of food, agricultural commodities, wood and wood products or animal feedstuffs, or substances that may be administered to animals for the control of insects, arachnids, or other pests in or on their bodies. The term includes substances intended for use as a plant growth regulator, defoliant, desiccant, or agent for thinning fruit or preventing the premature fall of fruit. Also used as substances applied to crops either before or after harvest to protect the commodity from deterioration during storage and transport in soil and water of district Ganganagar, Rajasthan. [1,2,3]

Pesticides can be classified by target organism (e.g., herbicides, insecticides, fungicides, rodenticides, and pediculicides^[6] – see table), chemical structure (e.g., organic, inorganic, synthetic, or biological (biopesticide)^[7] although the distinction can sometimes blur), and physical state (e.g. gaseous (fumigant)).^[7] Biopesticides include microbial pesticides and biochemical pesticides.^[8] Plant-derived pesticides, or "botanicals", have been developing quickly. These include the pyrethroids, rotenoids, nicotinoids, and a fourth group that includes strychnine and scilliroside.^{[9]:15}

Many pesticides can be grouped into chemical families. Prominent insecticide families include organochlorines, organophosphates, and carbamates. Organochlorine hydrocarbons (e.g., DDT) could be separated into dichlorodiphenyl ethanes, cyclodiene compounds, and other related compounds. They operate by disrupting the sodium/potassium balance of the nerve fiber, forcing the nerve to transmit continuously. Their toxicities vary greatly, but they have been phased out because of their persistence and potential to bioaccumulate.^{[9]:239-240} Organophosphate and carbamates largely replaced organochlorines. Both operate through inhibiting the enzyme acetylcholinesterase, allowing acetylcholine to transfer nerve impulses indefinitely and causing a variety of symptoms such as weakness or paralysis. Organophosphates are quite toxic to vertebrates and have in some cases been replaced by less toxic carbamates.^{[9]:136-137} Thiocarbamate and dithiocarbamates are subclasses of carbamates. Prominent families of herbicides include phenoxy and benzoic acid herbicides (e.g. 2,4-D), triazines (e.g., atrazine), ureas (e.g., diuron), and Chloroacetanilide (e.g., alachlor). Phenoxy compounds tend to selectively kill broad-leaf weeds rather than grasses. The phenoxy and benzoic acid herbicides function similar to plant growth hormones, and grow cells without normal cell division, crushing the plant's nutrient transport system.^{[9]:300} Triazines interfere with photosynthesis in soil and water of district Ganganagar, Rajasthan.



The application of pest control agents is usually carried out by dispersing the chemical in an (often hydrocarbon-based) solvent-surfactant system to give a homogeneous preparation. A virus lethality study performed in 1977 demonstrated that a particular pesticide did not increase the lethality of the virus. Combinations that included surfactants and the solvent clearly showed that pretreatment with them markedly increased the viral lethality in the test mice.^[10]

Pesticides can be classified based upon their biological mechanism function or application method. Most pesticides work by poisoning pests.^[11] A systemic pesticide moves inside a plant following absorption by the plant. With insecticides and most fungicides, this movement is usually upward (through the xylem) and outward. Increased efficiency may be a result. Systemic insecticides, which poison pollen and nectar in the flowers, may kill bees and other needed pollinators.^[12]

In 2010, the development of a new class of fungicides called paldoxins was announced. These work by taking advantage of natural defense chemicals released by plants called phytoalexins, which fungi then detoxify using enzymes. The paldoxins inhibit the fungi's detoxification enzymes. They are believed to be safer and greener in soil and water of district Ganganagar, Rajasthan[7,8,9]

II. DISCUSSION

Type of pesticide	Target pest group
Algicides or algaecides	Algae
Avicides	Birds
Bactericides	Bacteria
Fungicides	Fungi and oomycetes
Herbicides	Plant
Insecticides	Insects
Lampricides	Lampreys ^[3]
Miticides or acaricides	Mites
Molluscicides	Snails
Nematicides	Nematodes
Rodenticides	Rodents
Slimicides	Algae, Bacteria, Fungi, and Slime molds
Virucides	Viruses

Pesticides are used to control organisms that are considered to be harmful, or pernicious to their surroundings.^[28] For example, they are used to kill mosquitoes that can transmit potentially deadly diseases like West Nile virus, yellow fever, and malaria. They can also kill bees, wasps or ants that can cause allergic reactions. Insecticides can protect animals from illnesses that can be caused by parasites such as fleas.^[28] Pesticides can prevent sickness in humans that could be caused by moldy food or diseased produce. Herbicides can be used to clear roadside weeds, trees, and brush. They can also kill invasive weeds that may cause environmental damage. Herbicides are commonly applied in ponds and lakes to control algae and plants such as water grasses that can interfere with activities like swimming and fishing and cause the water to look or smell unpleasant.^[29] Uncontrolled pests such as termites and mold can damage structures such as houses.^[28] Pesticides are used in grocery stores and food storage facilities to manage rodents and insects that infest food such as grain. Each use of a pesticide carries some associated risk. Proper pesticide use decreases these associated risks to a level deemed acceptable by pesticide regulatory agencies in Sri Ganganagar, Rajasthan

DDT, sprayed on the walls of houses, is an organochlorine that has been used to fight malaria since the 1950s. Recent policy statements by the World Health Organization have given stronger support to this approach.^[30] It and other organochlorine pesticides have been banned in most countries worldwide because of their persistence in the

environment and human toxicity. DDT use is not always effective, as resistance to DDT was identified in Africa as early as 1955, and by 1972 nineteen species of mosquito worldwide were resistant to DDT.^{[31][32]}

III. RESULTS

Pesticides can save farmers' money by preventing crop losses to insects and other pests; in the U.S., farmers get an estimated fourfold return on money they spend on pesticides.^[37] One study found that not using pesticides reduced crop yields by about 10%.^[38] Another study, conducted in 1999, found that a ban on pesticides in the Sri ganganagr may result in a rise of food prices, loss of jobs, and an increase in Rajasthan hunger.^[39]

There are two levels of benefits for pesticide use, primary and secondary. Primary benefits are direct gains from the use of pesticides and secondary benefits are effects that are more long-term.^[40]

Biological

Controlling pests and plant disease vectors in Rajasthan

- Improved crop yields
- Improved crop/livestock quality
- Invasive species controlled

Controlling human/livestock disease vectors and nuisance organisms in Sri ganganar Rajasthan[10,11,12]

- Human lives saved and disease reduced. Diseases controlled include malaria,^[40] with millions of lives having been saved or enhanced with the use of DDT alone.^[41]
- Animal lives saved and disease reduced

Controlling organisms that harm other human activities and structures

- Drivers view unobstructed
- Tree/brush/leaf hazards prevented
- Wooden structures protected^[40]

Monetary

In one study, it was estimated that for every dollar (\$1) that is spent on pesticides for crops can yield up to four dollars (\$4) in crops saved.^[42] This means based that, on the amount of money spent per year on pesticides, \$10 billion, there is an additional \$40 billion savings in crop that would be lost due to damage by insects and weeds. In general, farmers benefit from having an increase in crop yield and from being able to grow a variety of crops throughout the year. Consumers of agricultural products also benefit from being able to afford the vast quantities of produce available year-round.^[40]

Post- WWII conditions caused the pesticide industry to flourish for several reasons including the growing middle class and the invention of cheap tractor-drawn spraying equipment. By the 1980s the demand for pesticides had dropped due to farmers struggling financially and the market for chemicals becoming oversaturated. There were also new costs for producing pesticides due to the strict EPA laws surrounding the chemicals. The modern pesticide market is seven billion dollars and is growing 4% per year due to the invention of the lawn and the stigma surrounding the untamed yard and in soil and water of district Ganganagar, Rajasthan

Costs

On the cost side of pesticide use there can be costs to the environment, costs to human health,^[44] as well as costs of the development and research of new pesticides.[15,17,18]

Health effects



A sign warning about potential pesticide exposure

Pesticides may cause acute and delayed health effects in people who are exposed.^[45] Pesticide exposure can cause a variety of adverse health effects, ranging from simple irritation of the skin and eyes to more severe effects such as affecting the nervous system, hearing,^[46] mimicking hormones causing reproductive problems, and also causing cancer.^[47] A 2007 systematic review found that "most studies on non-Hodgkin lymphoma and leukemia showed positive associations with pesticide exposure" and thus concluded that cosmetic use of pesticides should be decreased.^[48] There is substantial evidence of associations between organophosphate insecticide exposures and neurobehavioral alterations.^{[49][50][51][52]} Limited evidence also exists for other negative outcomes from pesticide exposure including neurological, birth defects, and fetal death.^[53] when added in soil and water of district Ganganagar, Rajasthan

The American Academy of Pediatrics recommends limiting exposure of children to pesticides and using safer alternatives.^[54]

Owing to inadequate regulation and safety precautions, 99% of pesticide-related deaths occur in developing countries that account for only 25% of pesticide usage.^[55]

One study found pesticide self-poisoning the method of choice in one third of suicides worldwide, and recommended, among other things, more restrictions on the types of pesticides that are most harmful to humans when used in soil and water of district Ganganagar, Rajasthan

A 2014 epidemiological review found associations between autism and exposure to certain pesticides, but noted that the available evidence was insufficient to conclude that the relationship was causal.^[57]


Occupational exposure among agricultural workers in soil and water of district Ganganagar, Rajasthan [19,20]

At-home pesticide use, use of unregulated products, and the role of undocumented workers within the agricultural industry makes characterizing true pesticide exposure a challenge. It is estimated that 50–80% of pesticide poisoning cases are unreported.

Underreporting of pesticide poisoning is especially common in areas where agricultural workers are less likely to seek care from a healthcare facility that may be monitoring or tracking the incidence of acute poisoning. The extent of unintentional pesticide poisoning may be much greater than available data suggest, particularly among developing countries. The, agriculture and food production remain one of the largest industries. In Sri ganaganagr, Rajasthan agricultural industry represents one of the largest sectors of the economy, with nearly 80% of its population relying on agriculture for income.^[62] Farmers in these communities rely on pesticide products to maintain high crop yields.

Some East Africa governments are shifting to corporate farming, and opportunities for foreign conglomerates to operate commercial farms have led to more accessible research on pesticide use and exposure among workers. In other areas where large proportions of the population rely on subsistence, small-scale farming, estimating pesticide use and exposure is more difficult.

Pesticide poisoning in Sri Ganganagar Rajasthan



The cholinergic synapse and the breakdown of acetylcholine into choline and acetate by acetylcholinesterase

Pesticides may exhibit toxic effects on humans and other non-target species, the severity of which depends on the frequency and magnitude of exposure. Toxicity also depends on the rate of absorption, distribution within the body, metabolism, and elimination of compounds from the body. Commonly used pesticides like organophosphates and carbamates act by inhibiting acetylcholinesterase activity, which prevents the breakdown of acetylcholine at the neural synapse. Excess acetylcholine can lead to symptoms like muscle cramps or tremors, confusion, dizziness and nausea. Studies show that farm workers in soil and water of district Ganganagar, Rajasthan have decreased concentrations of plasma acetylcholinesterase, the enzyme responsible for breaking down acetylcholine acting on synapses throughout the nervous system.^{[63][64][65]} Other studies have observed reduced respiratory function among farm workers who spray crops with pesticides.^[66] Numerous exposure pathways for farm workers increase the risk of pesticide poisoning, including dermal absorption walking through fields and applying products, as well as inhalation exposure in soil and water of district Ganganagar, Rajasthan

Measuring exposure to pesticides in soil and water of district Ganganagar, Rajasthan

There are multiple approaches to measuring a person's exposure to pesticides, each of which provides an estimate of an individual's internal dose. Two broad approaches include measuring biomarkers and markers of biological effect.^[67] The former involves taking direct measurement of the parent compound or its metabolites in various types of media: urine, blood, serum. Biomarkers may include a direct measurement of the compound in the body before it's been biotransformed during metabolism. Other suitable biomarkers may include the metabolites of the parent compound after they've been biotransformed during metabolism.^[67] Toxicokinetic data can provide more detailed information on how quickly the compound is metabolized and eliminated from the body, and provide insights into the timing of exposure.

Markers of biological effect provide an estimation of exposure based on cellular activities related to the mechanism of action. For example, many studies investigating exposure to pesticides often involve the quantification of the acetylcholinesterase enzyme at the neural synapse to determine the magnitude of the inhibitory effect of organophosphate and carbamate pesticides.^{[67][63][64][65]}

Another method of quantifying exposure involves measuring, at the molecular level, the amount of pesticide interacting with the site of action. These methods are more commonly used for occupational exposures where the mechanism of action is better understood, as described by WHO guidelines published in "Biological Monitoring of Chemical Exposure in the Workplace".^[68] Better understanding of how pesticides elicit their toxic effects is needed before this method of exposure assessment can be applied to occupational exposure of agricultural workers.



Alternative methods to assess exposure include questionnaires to discern from participants whether they are experiencing symptoms associated with pesticide poisoning. Self-reported symptoms may include headaches, dizziness, nausea, joint pain, or respiratory symptoms.^[64]

Challenges in assessing pesticide exposure in soil and water of district Ganganagar, Rajasthan[18,19,20]

Multiple challenges exist in assessing exposure to pesticides in the general population, and many others that are specific to occupational exposures of agricultural workers. Beyond farm workers, estimating exposure to family members and children presents additional challenges, and may occur through "take-home" exposure from pesticide residues collected on clothing or equipment belonging to parent farm workers and inadvertently brought into the home. Children may also be exposed to pesticides prenatally from mothers who are exposed to pesticides during pregnancy.^[69] Characterizing children's exposure resulting from drift of airborne and spray application of pesticides is similarly challenging, yet well documented in developing countries.^[70] Because of critical development periods of the fetus and newborn children, these non-working populations are more vulnerable to the effects of pesticides, and may be at increased risk of developing neurocognitive effects and impaired development.^{[71][69]}

While measuring biomarkers or markers of biological effects may provide more accurate estimates of exposure, collecting these data in the field is often impractical and many methods are not sensitive enough to detect low-level concentrations. Rapid cholinesterase test kits exist to collect blood samples in the field. Conducting large scale assessments of agricultural workers in remote regions of developing countries makes the implementation of these kits a challenge.^[67] The cholinesterase assay is a useful clinical tool to assess individual exposure and acute toxicity. Considerable variability in baseline enzyme activity among individuals makes it difficult to compare field measurements of cholinesterase activity to a reference dose to determine health risk associated with exposure.^[67] Another challenge researchers face in deriving a reference dose is identifying health endpoints that are relevant to exposure. More epidemiological research is needed to identify critical health endpoints, particularly among populations who are occupationally exposed.

Prevention

Minimizing harmful exposure to pesticides can be achieved by proper use of personal protective equipment, adequate reentry times into recently sprayed areas, and effective product labeling for hazardous substances as per FIFRA regulations. Training high-risk populations, including agricultural workers, on the proper use and storage of pesticides, can reduce the incidence of acute pesticide poisoning and potential chronic health effects associated with exposure. Continued research into the human toxic health effects of pesticides serves as a basis for relevant policies and enforceable standards that are health protective to all populations.

Environmental effects

Pesticide use raises a number of environmental concerns. Over 98% of sprayed insecticides and 95% of herbicides reach a destination other than their target species, including non-target species, air, water and soil.^[21] Pesticide drift occurs when pesticides suspended in the air as particles are carried by wind to other areas, potentially contaminating them. Pesticides are one of the causes of water pollution, and some pesticides are persistent organic pollutants and contribute to soil and flower (pollen, nectar) contamination.^[72] Furthermore, pesticide use can adversely impact neighboring agricultural activity, as pests themselves drift to and harm nearby crops that have no pesticide used on them in soil and water of district Ganganagar, Rajasthan

In addition, pesticide use reduces biodiversity, contributes to pollinator decline,^{[74][75][76]} destroys habitat (especially for birds),^[77] and threatens endangered species.^[21] Pests can develop a resistance to the pesticide (pesticide resistance), necessitating a new pesticide. Alternatively a greater dose of the pesticide can be used to counteract the resistance, although this will cause a worsening of the ambient pollution problem.

There are 12 most dangerous and persistent organic chemicals that were (now mostly obsolete) organochlorine pesticides.^{[61][78]} Since chlorinated hydrocarbon pesticides dissolve in fats and are not excreted, organisms tend to retain them almost indefinitely. Biological magnification is the process whereby these chlorinated hydrocarbons (pesticides) are more concentrated at each level of the food chain. Among marine animals, pesticide concentrations are higher in carnivorous fishes, and even more so in the fish-eating birds and mammals at the top of the ecological pyramid.^[79] The distillation is the process whereby pesticides are transported from warmer to colder regions of the Earth, in particular the Poles and mountain tops. Pesticides that evaporate into the atmosphere at relatively high temperature can be carried considerable distances (thousands of kilometers) by the wind to an area of lower temperature, where they condense and are carried back to the ground in rain or snow.^[80]



In order to reduce negative impacts, it is desirable that pesticides be degradable or at least quickly deactivated in the environment. Such loss of activity or toxicity of pesticides is due to both innate chemical properties of the compounds and environmental processes or conditions.[81] For example, the presence of halogens within a chemical structure often slows down degradation in an aerobic environment.[82] Adsorption to soil may retard pesticide movement, but also may reduce bioavailability to microbial degraders in soil and water of district Ganganagar, Rajasthan
Economics of pesticides in soil and water of district Ganganagar, Rajasthan

In one study, the human health and environmental costs due to pesticides in soil and water of district Ganganagar, Rajasthan was estimated to get higher in increased agricultural production.[84]
Resistance against pesticides in soil and water of district Ganganagar, Rajasthan[10,11]

The use of pesticides inherently entails the risk of resistance developing. Various techniques and procedures of pesticide application can slow the development of resistance, as can some natural features of the target population and surrounding environment.[3]
Alternatives uses against pesticides in soil and water of district Ganganagar, Rajasthan

Alternatives to pesticides are available and include methods of cultivation, use of biological pest controls (such as pheromones and microbial pesticides), genetic engineering (mostly of crops), and methods of interfering with insect breeding.[21] Application of composted yard waste has also been used as a way of controlling pests.[85]
These methods are becoming increasingly popular and often are safer than traditional chemical pesticides. In addition, EPA is registering reduced-risk pesticides in increasing numbers in soil and water of district Ganganagar, Rajasthan

Cultivation practices in Ganganagar Rajasthan Cultivation practices include polyculture (growing multiple types of plants), crop rotation, planting crops in areas where the pests that damage them do not live, timing planting according to when pests will be least problematic, and use of trap crops that attract pests away from the real crop.[21] Trap crops have successfully controlled pests in some commercial agricultural systems while reducing pesticide usage.[86] In other systems, trap crops can fail to reduce pest densities at a commercial scale, even when the trap crop works in controlled experiments.[87]
Use of other organisms in soil and water of district Ganganagar, Rajasthan

Release of other organisms that fight the pest is another example of an alternative to pesticide use. These organisms can include natural predators or parasites of the pests.[21] Biological pesticides based on entomopathogenic fungi, bacteria and viruses causing disease in the pest species can also be used.[21]
Biological control engineering in soil and water of district Ganganagar, Rajasthan

Interfering with insects' reproduction can be accomplished by sterilizing males of the target species and releasing them, so that they mate with females but do not produce offspring.[21] This technique was first used on the screwworm fly in 1958 and has since been used with the medfly, the tsetse fly,[88] and the gypsy moth.[89] This is a costly and slow approach that only works on some types of insects.[21]
Other

Other alternatives include "laserweeding" – the use of novel agricultural robots for weed control using lasers.[90]
Push pull strategysused in soil and water of district Ganganagar, Rajasthan

The term "push-pull" was established in 1987 as an approach for integrated pest management (IPM). This strategy uses a mixture of behavior-modifying stimuli to manipulate the distribution and abundance of insects. "Push" means the insects are repelled or deterred away from whatever resource is being protected. "Pull" means that certain stimuli (semiochemical stimuli, pheromones, food additives, visual stimuli, genetically altered plants, etc.) are used to attract



pests to trap crops where they will be killed.[91] There are numerous different components involved in order to implement a Push-Pull Strategy in IPM.

Many case studies testing the effectiveness of the push-pull approach have been done across the world. The most successful push-pull strategy was developed for subsistence farming. Another successful case study was performed on the control of *Helicoverpa* in cotton crops. The push-pull strategies were successfully used in the controlling of *Sitona lineatus* in bean fields.[91]

Some advantages of using the push-pull method are less use of chemical or biological materials and better protection against insect habituation to this control method. Some disadvantages of the push-pull strategy are that if there is a lack of appropriate knowledge of the behavioral and chemical ecology of the host-pest interactions then this method becomes unreliable. Furthermore, because the push-pull method is not a very popular method of IPM operational and registration costs are higher.[20]

Effectiveness in soil and water of district Ganganagar, Rajasthan

Some evidence shows that alternatives to pesticides can be equally effective as the use of chemicals. A study of Maize fields found that the application of composted yard waste with high carbon to nitrogen ratio to agricultural fields was highly effective at reducing the population of plant-parasitic nematodes and increasing crop yield, with yield increases ranging from 10% to 212%; the observed effects were long-term, often not appearing until the third season of the study.^[85] Additional silicon nutrition protects some horticultural crops against fungal diseases almost completely, while insufficient silicon sometimes leads to severe infection even when fungicides are used.^[92]

Pesticide resistance is increasing and that may make alternatives more attractive.

Types

Pesticides are often referred to according to the type of pest they control. Pesticides can also be considered as either biodegradable pesticides, which will be broken down by microbes and other living beings into harmless compounds, or persistent pesticides, which may take months or years before they are broken down: it was the persistence of DDT, for example, which led to its accumulation in the food chain and its killing of birds of prey at the top of the food chain. Another way to think about pesticides is to consider those that are chemical pesticides are derived from a common source or production method.^[93]

Insecticides

Neonicotinoids are a class of neuro-active insecticides chemically similar to nicotine. Imidacloprid, of the neonicotinoid family, is the most widely used insecticide in the world.^[94] In the late 1990s neonicotinoids came under increasing scrutiny over their environmental impact and were linked in a range of studies to adverse ecological effects, including honey-bee colony collapse disorder (CCD) and loss of birds due to a reduction in insect populations. Organophosphate and carbamate insecticides have a similar mode of action. They affect the nervous system of target pests (and non-target organisms) by disrupting acetylcholinesterase activity, the enzyme that regulates acetylcholine, at nerve synapses. This inhibition causes an increase in synaptic acetylcholine and overstimulation of the parasympathetic nervous system.^[102] Many of these insecticides, first developed in the mid 20th century, are very poisonous. Although commonly used in the past, many older chemicals have been removed from the market due to their health and environmental effects (e.g. DDT, chlordane, and toxaphene).^{[103][104][105]} Many organophosphates do not persist in the environment.

Pyrethroid insecticides were developed as a synthetic version of the naturally occurring pesticide pyrethrin, which is found in chrysanthemums. They have been modified to increase their stability in the environment. Some synthetic pyrethroids are toxic to the nervous system.^[106]

Herbicides

A number of sulfonylureas have been commercialized for weed control, including: amidosulfuron, flazasulfuron, metsulfuron-methyl, rimsulfuron, sulfometuron-methyl, terbacil,^[107] nicosulfuron,^[108] and triflurosulfuron-methyl.^[109] These are broad-spectrum herbicides that kill plants weeds or pests by inhibiting the enzyme acetolactate synthase. In the 1960s, more than 1 kg/ha (0.89 lb/acre) crop protection chemical was typically applied, while sulfonylureates allow as little as 1% as much material to achieve the same effect.^[110]

Biopesticides used in soil and water of district Ganganagar, Rajasthan

Biopesticides are certain types of pesticides derived from such natural materials as animals, plants, bacteria, and certain minerals. For example, canola oil and baking soda have pesticidal applications and are considered biopesticides. Biopesticides fall into three major classes:

- Microbial pesticides which consist of bacteria, entomopathogenic fungi or viruses (and sometimes includes the metabolites that bacteria or fungi produce). Entomopathogenic nematodes are also often classed as microbial pesticides, even though they are multi-cellular.^{[111][112]}
- Biochemical pesticides or herbal pesticides^[113] are naturally occurring substances that control (or monitor in the case of pheromones) pests and microbial diseases.
- Plant-incorporated protectants (PIPs) have genetic material from other species incorporated into their genetic material (i.e. GM crops).

IV. CONCLUSION

Pesticide residue refers to the pesticides that may remain on or in food after they are applied to food crops.^[135] The maximum allowable levels of these residues in foods are often stipulated by regulatory bodies in many countries. Regulations such as pre-harvest intervals also often prevent harvest of crop or livestock products if recently treated in order to allow residue concentrations to decrease over time to safe levels before harvest. Exposure of the general population to these residues most commonly occurs through consumption of treated food sources, or being in close contact to areas treated with pesticides such as farms or lawns in soil and water of district Ganganagar, Rajasthan^[19]

Many of these chemical residues, especially derivatives of chlorinated pesticides, exhibit bioaccumulation which could build up to harmful levels in the body as well as in the environment.^[137] The problem is most acute in China, the largest producer of chlorinated pesticides.^[138] Persistent chemicals can be magnified through the food chain and have been detected in products ranging from meat, poultry, and fish, to vegetable oils, nuts, and various fruits and vegetables.^[139]

Pesticide contamination in the environment can be monitored through bioindicators such as bee pollinators.^[72]

There is an ongoing research focused on pesticide residues in farming system^[20]

REFERENCES

1. "Basic Information about Pesticide Ingredients". Environmental Protection Agency. April 2, 2018. Retrieved 2018-12-01.
2. ^ Randall C, et al. (2014). "Pest Management". National Pesticide Applicator Certification Core Manual (2nd ed.). Washington: National Association of State Departments of Agriculture Research Foundation.
3. ^ Dunlop, Erin S.; McLaughlin, Rob; Adams, Jean V.; Jones, Michael; Birceanu, Oana; Christie, Mark R.; Criger, Lori A.; Hinderer, Julia L.M.; Hollingworth, Robert M.; Johnson, Nicholas S.; Lantz, Stephen R.; Li, Weiming; Miller, James; Morrison, Bruce J.; Mota-Sanchez, David; Muir, Andrew; Sepúlveda, Maria S.; Steeves, Todd; Walter, Lisa; Westman, Erin; Wirgin, Isaac; Wilkie, Michael P. (2018). "Rapid evolution meets invasive species control: the potential for pesticide resistance in sea lamprey". *Canadian Journal of Fisheries and Aquatic Sciences*. National Research Council Canada. 75 (1): 152–168. doi:10.1139/cjfas-2017-0015. hdl:1807/78674. ISSN 0706-652X.
4. ^ "Pesticide Industry Sales and Usage" (PDF). EPA. Retrieved 17 February 2020.
5. ^ "International Code of Conduct on the Distribution and Use of Pesticides" (PDF). Food and Agriculture Organization of the United Nations. 2002. Archived from the original (PDF) on 4 April 2013.
6. ^ Golden RC, Huffling K, Sattler B (January 2010). "Pesticides and health risks". *Journal of Obstetric, Gynecologic, and Neonatal Nursing*. 39 (1): 103–10. doi:10.1111/j.1552-6909.2009.01092.x. PMID 20409108.
7. ^ "Educational and Informational Strategies to Reduce Pesticide Risks". *Preventive Medicine*. 26 (2): 191–200. 1997. doi:10.1006/pmed.1996.0122. ISSN 0091-7435. PMID 9085387.
8. ^ "Types of Pesticide Ingredients". US Environmental Protection Agency. Jan 3, 2017. Retrieved Dec 1, 2018.
9. ^ Kamrin MA (1997). *Pesticide Profiles: Toxicity, Environmental Impact, and Fate* (1st ed.). Boca Raton: CRC. ISBN 978-1566701907. OCLC 35262311.
10. ^ Safe S, Plugge H, Crocker JF (1977). "Analysis of an aromatic solvent used in a forest spray program". *Chemosphere*. 6 (10): 641–651. Bibcode:1977Chmsp...6..641S. doi:10.1016/0045-6535(77)90075-3. ISSN 0045-6535.



11. ^ "Pesticide Applicator Core Tutorial: Module 4 - Toxicity of Pesticides". Pesticide Safety Education Program (PSEP). Cornell University. Archived from the original on July 21, 2020. Retrieved Dec 1, 2018.
12. ^ Rortais A, Arnold G, Halm MP, Touffet-Briens F (2005). "Modes of honeybees exposure to systemic insecticides: estimated amounts of contaminated pollen and nectar consumed by different categories of bees". *Apidologie*. 36 (1): 71–83. doi:10.1051/apido:2004071.
13. ^ American Chemical Society (2009). "New 'green' pesticides are first to exploit plant defenses in battle of the fungi". *EurekAlert!*. AAAS. Retrieved Dec 1, 2018.
14. ^ Rao GV, Rupela OP, Rao VR, Reddy YV (2007). "Role of biopesticides in crop protection: present status and future prospects" (PDF). *Indian Journal of Plant Protection*. 35 (1): 1–9. Archived (PDF) from the original on 2018-12-02.
15. ^ Miller GT (2002). *Living in the Environment* (12th ed.). Belmont: Wadsworth/Thomson Learning. ISBN 9780534376970. OCLC 819417923.
16. ^ Ritter SK (2009). "Pinpointing Trends In Pesticide Use In 1939". *Chemical & Engineering News*. Vol. 87, no. 7. ACS. ISSN 0009-2347.
17. ^ Goldman LR (2007). "Managing pesticide chronic health risks: U.S. policies". *Journal of Agromedicine*. 12 (1): 67–75. doi:10.1300/J096v12n02_08. PMID 18032337. S2CID 216149465.
18. ^ Daly HV, Doyen JT, Purcell AH (1998). "Chapter 14". *Introduction to insect biology and diversity* (2nd ed.). Oxford: Oxford University Press. pp. 279–300. ISBN 978-0195100334. OCLC 37211384.
19. ^ Steen-Adams, Michelle (2002). "Russell, E. 2001. War and Nature: Fighting Humans and Insects with Chemicals from World War I to Silent Spring. Cambridge University Press, Cambridge, UK and New York, New York, USA". *Conservation Ecology*. 6 (2). doi:10.5751/es-00407-060201. ISSN 1195-5449.
20. ^ Murphy G (Dec 1, 2005). "Pesticide Rotation". Ontario Ministry of Agriculture, Food and Rural Affairs. Archived from the original on October 13, 2007. Retrieved Sep 15, 2007.



Impact Factor:
7.512



INTERNATIONAL JOURNAL OF INNOVATIVE RESEARCH

IN SCIENCE | ENGINEERING | TECHNOLOGY

 9940 572 462  6381 907 438  ijirset@gmail.com



www.ijirset.com

Scan to save the contact details