



e-ISSN: 2319-8753 | p-ISSN: 2320-6710

# IJRSET

International Journal of Innovative Research in  
**SCIENCE | ENGINEERING | TECHNOLOGY**

# INTERNATIONAL JOURNAL OF INNOVATIVE RESEARCH

IN SCIENCE | ENGINEERING | TECHNOLOGY

Volume 10, Issue 3, March 2021

**ISSN** INTERNATIONAL  
STANDARD  
SERIAL  
NUMBER  
INDIA

**Impact Factor: 7.512**

9940 572 462

6381 907 438

[ijirset@gmail.com](mailto:ijirset@gmail.com)

[www.ijirset.com](http://www.ijirset.com)



# COVID-19 Risk Factors among Health Workers

**D. Navaneetha Krishnan, S. Sheik Sulaiman**

Final year PG Student, Department of Industrial Safety Engineering, Francis Xavier Engineering, College, Tirunelveli, Tamil Nadu, India

Professor, Department of Mechanical Engineering, Francis Xavier Engineering College, Tirunelveli, Tamil Nadu, India

**ABSTRACT:** Coronavirus disease 2019 (Covid-19) poses a crucial activity health risk to medical experts (HWs) that has attracted world scrutiny. To date, many thousand HWs globally are according as infected with the severe acute metabolism syndrome coronavirus a pair of virus that causes the malady. it's thus a public health priority for policymakers to grasp risk factors for this vulnerable cluster to avert activity transmission. A fast review was disbursed on twenty Apr 2020 on Covid-19 risk factors among HWs in PubMed, Google Scholar, and EBSCOHost net (Academic Search Complete, CINAHL Complete, phone system with Full Text, CINAHL with Full Text, APA PsycInfo, Health Source—Consumer Edition, Health Source: Nursing/Academic Edition) and United Nations agency world info. we have a tendency to conjointly hunted for preprints on the medRxiv info. we have a tendency to hunted for reports, reviews, and first data-based studies (case management, case cross-over, cross-sectional, and cohort). The review enclosed studies conducted among HWs with Covid-19 that according risk factors regardless of their sample size. Eleven studies met the inclusion criteria. Lack of private protecting instrumentation, exposure to infected patients, work overload, poor infection management, and pre-existent medical conditions were known as risk factors for Covid-19 among HWs. within the context of Covid-19, HWs face associate unprecedented activity risk of morbidity and mortality. there's want for fast development of property measures that shield HWs from the pandemic. Previous article in issue.

## I. INTRODUCTION

The coronavirus malady (Covid-19) pandemic caused by the severe acute metabolism syndrome coronavirus two (SARS-CoV-2) has challenged the health systems of nations it's affected. As of thirty April 2020, the pandemic has infected quite three, 2 million people with over two 30,000 fatal cases in 187 countries. The person-to-person transmission routes of Covid-19 embody direct transmissions, like cough, sneeze, drop inhalation transmission, and call transmission, like the contact with oral, nasal, and eye mucose membranes. medical examiners (HWs) area unit at the forefront of the Covid-19 happening response and intrinsically area unit exposed to hazards that place them in danger of infection. As early as February 2020, students had established that transmission of the malady among HWs is related to overcrowding, absence of isolation space facilities, and environmental contamination. However, within the context of the chop-chop evolving Covid-19 pandemic, activity health and safety policymakers would like timely access to updated health data on the chance HWs face. This proof is best obtained through a scientific review; but, this review technique needs immense resources, together with time, to finish that don't suit the wants. during this study, we tend to aimed to map obtainable proof on Covid-19 risk factors among HWs to guide future analysis and policy.

## II. METHODS

### Information sources and literature search

We searched PubMed, Google Scholar, and EBSCOHost internet (Academic Search Complete, CINAHL Complete, phone system with Full Text, CINAHL with Full Text, APA PsycInfo, Health Source—Consumer Edition, Health Source: Nursing/Academic Edition) and UN agency world information for relevant studies. we tend to conjointly hunted for preprints on the medRxiv information. The search terms enclosed “Coronavirus,” “Covid-19,” “2019-nCoV,” “SARS-CoV-2,” “risk,” “risk factor,” “infection,” “illness,” “disease,” “health care worker,” “health care professional”, and “occupational health”. Boolean terms, AND and OR, were wont to separate the keywords. The reference lists of all full-text articles screened were hunted for relevant studies.

### Inclusion criteria



We explore for reports, reviews, and first experimental studies (case-control, case-cross-over, cross-sectional, and cohort). The review enclosed studies conducted among HWs that rumored COVID-19 risk factors regardless of their sample size.

#### **Screening process**

We developed a screening criterion a priori for every of the 3 stages, title and abstract and full text. 2 researchers screened the articles severally. variations in screen results at full-text stage were resolved by discussion.

#### **Data abstraction and analysis process**

A data abstraction type was developed, discussed, and revised a priori. the shape extracted data on 1st author, title, sample size, reportable risk factors, and different relevant findings. only 1 reviewer (TD) abstracted knowledge from the enclosed studies. For collating, summarizing, and reportage of the findings, first, the reviewers familiarised themselves with the content of the articles. Second, findings reportable within the articles were classified into classes supported the reportable factors, and a narrative was provided.

### **III. RESULTS**

Our initial keyword information search found one,698 probably eligible articles. Results from every information search area unit bestowed in Supplementary File one. Following title screening, eighty two articles were eligible for inclusion in abstract screening. These articles were foreign into EndNote X9 and eight duplicates removed, going away seventy four articles enclosed in abstract screening. Following abstract screening, thirty eight studies were excluded, going away thirty six articles for full-text screening. Twenty-five articles were excluded once full articles screening, going away eleven articles for knowledge extraction.

#### **Characteristics of included studies**

All enclosed studies according Covid-19 risk factors among HWs. Of these, one was a brief communication, 2 were reports, four were reviews, one retrospective cohort, one retrospective information analysis, one prospective cohort, and one case-control. All enclosed studies were printed in 2020. the full variety of HWs according within the studies was one,619; but, 5 studies didn't specify a sample size. One study was a preprint article unfinished review

#### **Study findings**

Eleven articles were enclosed for information extraction and narrative outline. The reportable Covid-19 risk factors among HWs were classified into five: lack of private protecting instrumentality (PPE), exposure to infected patients, work overload, poor infection management, and pre-existing medical conditions. meagerly PPE or lack therefrom was reportable as a vital risk issue for Covid-19. Exposure to infected patients chiefly through add speculative departments and contaminated fluid/aerosols were reportable as another vital Covid-19 risk issue among HWs. Work overload resulted in lack of rest, long-time exposure to infected patients, and dealing harassed. Longer duty hours were reportable as Covid-19 risk think about a retrospective cohort conducted at a hospital in city, China. A review conducted by Yu et al. discovered restricted or absence of information on infection management as a risk issue for Covid-19 among HWs within the radiology department. Similarly, a retrospective analysis by Tibeto-Burman language et al. and a fast review by Viswanath et al. discovered lack of established practices of infection management poor infection management and direct contact with contaminated surfaces as vital risk factors. astonishingly, suboptimal hand hygiene was reportable as a Covid-19 risk issue among HWs. A review conducted by Ing et al. on medico deaths from Covid-19 reportable older age (physicians fifty seven years more matured or older accounted for three-quarters of Covid-19-related deaths), male gender (90% of deaths were male physicians), and pre-existing conditions (hypertension, diabetes, disorder, chronic respiratory organ sickness, and upset individuals) as risk factors.

### **IV. DISCUSSION**

This review has discovered that analysis on Covid-19 risk factors for HWs continues to be in its infancy. whereas this is often to be expected given the pandemic solely bust enter Gregorian calendar month of 2019, the offered proof shows active analysis interest on the topic. The review discovered lack of PPE, exposure to infected patients, work overload, poor infection management, and antecedent medical conditions as Covid-19 risk factors among HWs. The UN agency free interim steering on HWs risk assessment and management within the context of Covid-19 three. The steering is



obtainable for each the HWs and their facilities. whereas this blanket steering is out there, activity health policymakers would wish to contextualize the procedures supported offered resources and knowledge.

Our review discovered exposure to infected patients and work overload as 2 necessary Covid-19 risk factors among HWs. This corroborates a recent Centers for sickness management and interference report that calculable that of the 9282 cases of Covid-19 cases among HWs within the us of America, fifty fifth had exposure within the attention setting. Further, a mathematical modeling study by Temime et al. found that R0 for Covid-19 was higher for HWs as compared thereupon of the final public as they need prolonged contact with infected people. HWs at greatest risk ar those that ar exposed to the airways and therefore the oral fissure of patients for prolonged periods like dentists. Our review found contact with contaminated fluid as a Covid-19 risk issue. Similarly, in step with Meng et al., dentists accounted for five of the fatalities among physicians exposed to Covid-19. considerations on contamination through suction have conjointly been rumored elsewhere. Covid-19 has overcome health systems globally leading to longer operating hours and inflated work for HWs. Italian physicians have advised a community-centered or home care system for Covid-19 to not solely reduce the work on HWs however conjointly decrease the transmission of sickness and HWs exposure. Further, there's a desire for social support services to assist HWs UN agency ar troubled showing emotion and permit for periodic breaks from work.

The protection of HWs is important despite the inflated demand and international shortage of PPE. to confirm the supply of adequate PPE for all HWs, governments have to be compelled to expedite the acquisition and dev ise strategic use of the offered PPE. as an example, in developing countries, the strategic use of PPE might embrace extended wear and apply of sure PPE like robes, particularly in low-risk hospital departments. With the Covid-19 pandemic, there's conjointly the potential for panic among HWs that will end in the irrational use of PPE. A balance can have to be compelled to be affected between HWs safety and strategic use of obtainable resources. Finally, HWs ought to be trained on correct ways in which of sporting and popping out PPE to forestall contamination.

The Covid-19 pandemic has place a strain on the health systemforce attributable to the overwhelming infection rates in some countries. This has meant that older HWs, principally doctors and nurses, have had to hitch as frontline staff and a few to come back out of retirement to supply expertise, expertise, leadership, and boost the morale of younger professionals. maturity aboard antecedent health conditions like cardiovascular disease, DM, upset, chronic respiratory organ sickness, and immunological disorder were necessary Covid-19 risk factors known during this review. These are touted as major risk factors for severe Covid-19 sickness elsewhere. Health facilities have to be compelled to assign duties for top Covid-19 risk departments supported age and underlying conditions.

This article incorporates a range of strengths and weaknesses. whereas there ar several existing papers handling activity health problems connected with the Covid-19, distinctive the danger factors for infection is a vital opening to guide acceptable measures to cut back these risks. this text provided a cryptic review of obtainable proof on the topic as at twenty Apr 2020, a amount once there was AN upward mechanical phenomenon within the range of cases among HWs globally. fast reviews naturally have multiple limitations. during this study, necessary omissions were created on steps for a full systematic review. These embrace publication language restriction and risk of bias/quality appraisal. However, this review can function a precursor to a bigger, thorough, and systematic review and meta-analysis the authors of this text ar acting on. Secondly, the proof on risk factors ar possible to be underreported given the fast evolution of the pandemic and lack of without delay offered knowledge from different elements of the planet like continent at the time of the review. Third, attributable to the heterogeneousness among enclosed articles, our review findings can not be generalized or clinically involved. However, we have a tendency to believe the review yielded necessary findings to tell activity health policymakers on Covid-19 risk factors among HWs and researchers on gaps in scientific proof.

## V. CONCLUSION

In the context of Covid-19, HWs face associate unexampled activity risk of morbidity and mortality. Lack of and/or inadequate PPE, exposure to infected patients, work overload, poor infection management, and pre-existent medical conditions place HWs in danger for healthcare facility Covid-19 infection. additional studies square measure required to tell the event of efficacious infection management measures. there's a requirement for fast development of property measures that shield HWs from the pandemic.



#### REFERENCES

1. WHO  
**Coronavirus disease 2019 (COVID-19) situation report – 101**(30 April 2020)  
[https://www.who.int/docs/default-source/coronaviruse/situation-reports/20200430-sitrep-101-covid-19.pdf?sfvrsn=2ba4e093\\_2](https://www.who.int/docs/default-source/coronaviruse/situation-reports/20200430-sitrep-101-covid-19.pdf?sfvrsn=2ba4e093_2), Accessed 1st May 2020
2. X. Peng, X. Xu, Y. Li, L. Cheng, X. Zhou, B. Ren  
**Transmission routes of 2019-nCoV and controls in dental practice** Int J Oral Sci, 12 (1) (2020) 9-9
3. WHO  
**Health Workers exposure risk assessment and management in the context of COVID-19 virus (Interim guidance)** (2020)  
<https://apps.who.int/iris/handle/10665/331340>, Accessed 23rd Apr 2020
4. Z. Wu, J.M. McGoogan  
**Characteristics of and important lessons from the coronavirus disease 2019 (COVID-19) outbreak in China: summary of a report of 72 314 cases from the Chinese Center for Disease Control and Prevention** Jama (2020)
5. A.C. Tricco, J. Antony, W. Zarin, *et al.*  
**A scoping review of rapid review methods** BMC Med, 13 (2015), p. 224
6. S. Khangura, K. Konnyu, R. Cushman, J. Grimshaw, D. Moher  
**Evidence summaries: the evolution of a rapid review approach** Syst Rev, 1 (2012) 10-10
7. M. Petticrew, H. Roberts  
**Systematic reviews in the social sciences: a practical guide** John Wiley & Sons (2008)
8. J. Wang, M. Zhou, F. Liu  
**Reasons for healthcare workers becoming infected with novel coronavirus disease 2019 (COVID-19) in China** J Hosp Infect (2020)
9. J. Schwartz, C.-C. King, M.-Y. Yen  
**Protecting healthcare workers during the coronavirus disease 2019 (COVID-19) outbreak: lessons from Taiwan's severe acute respiratory syndrome response** Clin Infect Dis (2020)
10. C.A. Grimm  
**Hospital experiences responding to the COVID-19 pandemic: results of a national pulse survey March 23–27, 2020** (2020)
11. <https://justthenews.com/sites/default/files/2020-04/HHS2020ReportPandemicHospitals.pdf>, Accessed 20th Apr 2020



**INNO**  **SPACE**  
SJIF Scientific Journal Impact Factor

**Impact Factor:  
7.512**

**ISSN** INTERNATIONAL  
STANDARD  
SERIAL  
NUMBER  
**INDIA**



# INTERNATIONAL JOURNAL OF INNOVATIVE RESEARCH

IN SCIENCE | ENGINEERING | TECHNOLOGY

 **9940 572 462**  **6381 907 438**  **ijirset@gmail.com**



[www.ijirset.com](http://www.ijirset.com)

Scan to save the contact details