



e-ISSN: 2319-8753 | p-ISSN: 2347-6710

IJIRSET

International Journal of Innovative Research in
SCIENCE | ENGINEERING | TECHNOLOGY



INTERNATIONAL JOURNAL OF INNOVATIVE RESEARCH

IN SCIENCE | ENGINEERING | TECHNOLOGY

Volume 13, Issue 1, January 2024

ISSN INTERNATIONAL
STANDARD
SERIAL
NUMBER
INDIA

Impact Factor: 8.423

9940 572 462

6381 907 438

ijirset@gmail.com

www.ijirset.com

Heavy Minerals: Key Tool in Sedimentological Research

Vikas Swarnkar, S. D. Deshmukh, Komal Singh Verma, Sana Siddique

Department of Geology, Govt. V. Y. T. PG Autonomous College, Durg, Chhattisgarh, India

ABSTRACT: Heavy minerals, comprising dense mineral grains resistant to weathering and erosion, serve as invaluable indicators for understanding sediment provenance in sedimentological research. This article deals with the properties and significance of heavy minerals as a key tool for interpreting sediment provenance, providing insights into the applications of heavy mineral analysis in deciphering the origin of sedimentary deposits.

KEYWORDS: Heavy minerals, provenance, Sedimentology, separation of heavy minerals.

I. INTRODUCTION

Heavy minerals are dense mineral grains with specific gravities significantly higher than the average density of most common rock-forming minerals. They typically exhibit densities greater than 2.9 g/cm³ and are commonly composed of minerals such as garnet, tourmaline, zircon, rutile, and magnetite. Heavy minerals are often concentrated in sedimentary deposits through various sedimentary processes, such as erosion, transport, and sedimentation.

Heavy minerals encompass a wide variety of mineral species, each with its own unique chemical composition, crystal structure, and physical properties. Heavy minerals have high densities, which make them resistant to weathering and erosion processes. Their density allows them to settle out of suspension more rapidly during sedimentation, leading to their concentration in sedimentary deposits. Many heavy minerals are characterized by high hardness and resistance to abrasion. This property contributes to their durability during transport and preservation in sedimentary rocks. They exhibit a wide range of colours, from transparent to opaque, depending on their mineral composition and crystal structure and distinct cleavage planes and fracture patterns, which may aid in their identification and characterization.

Heavy minerals are transported along with other sedimentary grains by various agents such as rivers, glaciers, wind, and waves. They tend to be preferentially concentrated in sedimentary environments where sorting is controlled by hydraulic sorting mechanisms or density sorting processes.

II. HISTORICAL DEVELOPMENT OF HEAVY MINERAL STUDIES

The recognition of heavy minerals in sedimentary rocks dates back to the early days of geological inquiry. Geologists such as Charles Lyell and William Smith noted the presence of dense mineral grains, including garnet, tourmaline, and zircon, in sedimentary deposits during the 19th century. In the late 19th and early 20th centuries, pioneering geologists like Charles Lapworth and Johann Gruson conducted systematic studies on heavy minerals in sedimentary rocks. Lapworth's work on the Scottish Highlands and Gruson's investigations in Germany laid the foundation for understanding heavy mineral assemblages and their significance in sedimentary provenance. The development of quantitative analytical techniques, such as optical microscopy and electron microscopy, enabled geologists to characterize heavy mineral compositions and abundances more precisely. This allowed for quantitative assessments of heavy mineral assemblages in sedimentary deposits and their implications for sediment provenance. The advent of modern analytical techniques, including X-ray diffraction (XRD), scanning electron microscopy (SEM), and energy-dispersive X-ray spectroscopy (EDS), revolutionized heavy mineral studies. These advanced methods allowed for detailed mineralogical and chemical analyses of heavy mineral grains, facilitating the identification of rare and accessory mineral phases.

III. METHODS OF SEPARATION OF HEAVY MINERALS

Appropriate separation techniques are used to isolate heavy mineral concentrates for further mineralogical, geochemical, and petrological analysis. Several methods are utilized for this purpose, each catering to different grain

size ranges and mineral densities and based on different separation criteria. Here are some commonly used methods of heavy mineral separation:

1. **Heavy Liquid Separation:**

Heavy liquid separation exploits the density contrast between heavy minerals and lighter minerals in a sediment sample. A heavy liquid with a known density (e.g., bromoform, sodium polytungstate) is added to the sample, causing heavy minerals to sink while lighter minerals float. The sediment sample is suspended in the heavy liquid and stirred to ensure thorough mixing. After settling, the denser heavy mineral fraction is collected from the bottom of the container, while the lighter fraction is decanted or skimmed off the surface. Heavy liquid separation is suitable for separating heavy minerals with densities higher than the heavy liquid used. It is commonly used for samples with grain sizes ranging from silt to sand.

2. **Magnetic Separation:**

Magnetic separation utilizes the magnetic properties of certain heavy minerals, such as magnetite and ilmenite, to separate them from non-magnetic minerals. A strong magnetic field is applied to the sample, causing magnetic minerals to be attracted to the magnet and separated from the non-magnetic fraction. The sediment sample is passed through a magnetic separator, which applies a magnetic field to the sample. Magnetic minerals are attracted to the magnetized region and collected separately, while non-magnetic minerals pass through unaffected. Magnetic separation is effective for isolating heavy minerals with magnetic properties, particularly iron-rich minerals. It is commonly used for samples with fine-grained magnetite or ilmenite.

3. **Electrostatic Separation:**

Electrostatic separation utilizes differences in the electrical conductivity and charge properties of minerals to separate them. A high-voltage electric field is applied to the sample, causing charged particles to be attracted or repelled depending on their electrical properties. The sediment sample is passed through an electrostatic separator, where it is subjected to a high-voltage electric field. Minerals with different electrical properties are deflected or attracted to oppositely charged electrodes, allowing for separation based on charge. Electrostatic separation is suitable for separating heavy minerals with contrasting electrical properties, such as conductive (e.g., metallic minerals) and non-conductive minerals (e.g., quartz, feldspar). It is commonly used for samples with fine-grained mineral mixtures.

4. **Gravity Separation:**

Gravity separation relies on the differences in settling velocities of heavy and light mineral grains in a fluid medium. Sediment samples are subjected to gravity forces, allowing denser particles to settle faster than lighter particles. The sediment sample is suspended in a fluid medium (e.g., water) and allowed to settle in a gravity separator or shaking table. Denser heavy minerals settle to the bottom under gravity, while lighter minerals remain suspended or settle more slowly.

Gravity separation is effective for separating heavy minerals based on density differences. It is commonly used for samples with coarse-grained heavy mineral assemblages, such as beach sands or alluvial deposits.

5. **Centrifugal Separation:**

Centrifugal separation utilizes centrifugal forces generated by high-speed rotation to separate particles based on their density and size. Sediment samples are spun in a centrifuge, causing denser particles to migrate outward while lighter particles concentrate toward the center. The sediment sample is placed in a centrifuge tube or rotor and spun at high speeds. Centrifugal forces cause heavier particles to move outward and settle along the tube wall, while lighter particles remain suspended or migrate toward the center. Centrifugal separation is effective for separating heavy minerals based on density and size differences. It is commonly used for samples with fine-grained heavy mineral assemblages or in situations where other separation methods are not feasible.

V. APPLICATIONS OF HEAVY MINERAL STUDIES

Heavy minerals serve as valuable tools in provenance studies in sedimentology, providing insights into the sources, transport pathways, and depositional environments of sedimentary deposits. By combining mineralogical, geochemical, and sedimentological analyses of heavy mineral assemblages, researchers can unravel the complex histories of sedimentary basins and enhance our understanding of Earth's dynamic surface processes. Some of the ways in which heavy minerals are used in provenance studies in sedimentology, are as follows:

1. **Mineralogy Identification:** Heavy minerals have distinct mineralogical compositions that reflect their source rocks. By analysing the types and proportions of heavy minerals present in sedimentary rocks, researchers can infer



the lithology and geological characteristics of the source areas. For example, specific heavy mineral species like garnet, tourmaline, and zircon are associated with different rock types and can be used as diagnostic indicators of sediment provenance.

2. **Mineral Assemblages:** The presence or absence of certain heavy mineral assemblages can provide clues about the depositional environment and sediment transport history. Different depositional environments (e.g., fluvial, marine, glacial) are associated with characteristic heavy mineral suites, reflecting the selective transport and sorting processes acting on sediment grains during transport. Comparing heavy mineral assemblages in sediments from different depositional settings allows researchers to distinguish between proximal and distal sediment sources.
3. **Mineral Chemistry:** Heavy minerals can exhibit variations in chemical composition due to differences in source lithology and diagenetic processes. Geochemical analyses of heavy minerals, such as X-ray fluorescence (XRF) or electron microprobe analysis, can provide insights into the elemental composition and provenance of sedimentary material. Isotopic analyses of certain heavy minerals, such as zircon, can also help constrain the age and origin of source rocks.
4. **Transport Pathways:** Heavy minerals are transported along with other sedimentary grains by various agents such as rivers, glaciers, wind, and waves. The degree of rounding, sorting, and maturity of heavy mineral grains can indicate the distance and duration of sediment transport. Abrasion and attrition during transport tend to round and reduce the size of heavy mineral grains, while sorting processes result in the selective enrichment or depletion of certain mineral types.
5. **Paleoenvironmental Reconstructions:** Changes in heavy mineral assemblages through time can provide insights into paleoenvironmental conditions and sedimentary processes. By analysing heavy mineral records in sedimentary sequences, researchers can reconstruct past climate variations, tectonic events, and sediment routing systems. Shifts in heavy mineral provenance can indicate changes in source area geology, erosion rates, or basin dynamics over geological time scales.
6. **Integration with Other Provenance Indicators:** Heavy mineral data are often integrated with other provenance indicators, such as detrital zircon geochronology, sedimentary petrology, and isotopic analyses, to build comprehensive models of sediment provenance.

VI. COMMON HEAVY MINERALS AND THEIR SIGNIFICANCE

S. No.	Name of heavy mineral	Provenance
01	Zircon	Zircon is abundant in granitic and metamorphic rocks. It is resistant to weathering and can be transported over long distances, making it a common heavy mineral in sedimentary deposits derived from continental sources.
02	Garnet	Garnet is often associated with metamorphic rocks such as schists, gneisses, and amphibolites. It can also occur in some igneous rocks like granites. The presence of garnet in sedimentary deposits suggests derivation from nearby mountainous regions or metamorphic terrains.
03	Tourmaline	Tourmaline is commonly found in granites, pegmatites, and metamorphic rocks such as schists and gneisses. It is resistant to weathering and can be transported by rivers and streams, indicating sedimentary deposits with a predominantly continental provenance.
04	Pyroxenes	Pyroxenes are common minerals in mafic and ultramafic igneous rocks such as basalt and gabbro. They can also occur in metamorphic rocks like amphibolites. The presence of pyroxenes in sedimentary deposits suggests input from volcanic source areas or ophiolitic complexes.
05	Magnetite	Magnetite is often associated with igneous rocks such as basalts and

		iron-rich sedimentary rocks like banded iron formations (BIFs). It can also occur as detrital grains in sedimentary deposits, indicating input from mafic or iron-rich source areas.
06	Rutile	Rutile is commonly found in metamorphic rocks such as schists and gneisses, as well as in some igneous rocks like granites. Its presence in sedimentary deposits suggests derivation from areas with high-grade metamorphic terrains or granitic source rocks.
07	Staurolite	Staurolite is typically associated with high-grade metamorphic rocks such as schists, gneisses, and amphibolites. It forms under conditions of high temperature and pressure during regional metamorphism, indicating input from metamorphic terrains in sedimentary deposits.
08	Sphene (Titanite)	Sphene is commonly found in igneous rocks such as granites and syenites, as well as in some metamorphic rocks like amphibolites and gneisses. Its presence in sedimentary deposits suggests derivation from areas with granitic or metamorphic source rocks.
09	Epidote	Epidote is often associated with low to medium-grade metamorphic rocks such as greenschists and metabasalts. It can also occur in some igneous rocks like diorites and gabbros. The presence of epidote in sedimentary deposits indicates input from regions with metamorphic or igneous source rocks.
10	Monazite	Monazite is commonly found in granitic and metamorphic rocks, including granite pegmatites, schists, and gneisses. Its presence in sedimentary deposits suggests derivation from areas with granitic or metamorphic source rocks, and it can be used for geochronological dating of sedimentary sequences.

VII. HEAVY MINERAL STUDIES IN INDIA

Heavy mineral analysis has been extensively utilized in India for various geological and mineralogical investigations, contributing to the understanding of sedimentary provenance, tectonic evolution, and mineral resource potential across different regions. Here are a few notable case studies showcasing the application of heavy mineral analysis in India:

1. Sedimentary Basins of Western India:

Heavy mineral studies have been conducted in sedimentary basins such as the Cambay Basin, Krishna-Godavari Basin, and Mumbai Offshore Basin to understand the provenance and depositional history of sedimentary sequences. These studies have revealed variations in heavy mineral compositions, indicating changes in sediment sources, transport pathways, and paleoenvironmental conditions during basin evolution.

2. Beach Placer Deposits of Kerala:

Kerala coast along the southwestern tip of India hosts significant beach placer deposits enriched with heavy minerals such as ilmenite, rutile, zircon, and garnet. Heavy mineral studies have been instrumental in characterizing these placer deposits, identifying potential economic resources, and evaluating mineral processing techniques for mineral separation and recovery.

3. Mineral Sands of Odisha and Andhra Pradesh:

Coastal regions of Odisha and Andhra Pradesh are known for their extensive mineral sand deposits, particularly along the beaches and dune systems. Heavy mineral analysis has been employed to assess the mineralogical composition, grain size distribution, and mineralogical variability of these deposits, aiding in resource estimation, exploration targeting, and mining planning.

4. Paleogeographic Reconstructions of Peninsular India:

Heavy mineral studies have contributed to paleogeographic reconstructions of Peninsular India, unraveling the tectonic history and sedimentary evolution of the region. Through the study of heavy mineral signatures in sedimentary rocks

and basin-fill sequences, researchers have inferred past depositional environments, sediment transport patterns, and tectonic events that have shaped the geological framework of the Indian subcontinent.

5. Coastal Erosion Studies:

Heavy mineral analysis has been utilized in coastal erosion studies along India's coastline to assess sediment sources, coastal dynamics, and erosion rates. Heavy mineral assemblages in beach sediments and offshore deposits, helped researchers to trace sediment transport pathways, evaluate coastal vulnerability, and develop strategies for coastal management and erosion mitigation.

VIII.CONCLUSION

Heavy mineral studies represent a valuable tool for sedimentologists and geologists in deciphering the complexities of sedimentary systems and understanding Earth's geological history. By integrating heavy mineral data with geological, geochemical, and geophysical datasets, and interpreting heavy mineral data in a multidisciplinary context, researchers can address fundamental questions about sediment provenance, depositional environments, and tectonic processes, thereby advancing our knowledge of Earth's dynamic surface processes and geological evolution.

REFERENCES

- [1] Andersen T, Kristoffersen M, Elburg MA (2016) How far can we trust provenance and crustal evolution information from detrital zircons? A South African case study. *Gondwana Res* 34:129–148. <https://doi.org/10.1016/j.gr.2016.03.003>
- [2] Andò S. (2020) Gravimetric Separation of Heavy Minerals in Sediments and Rocks. *Minerals*.10(3):273. <https://doi.org/10.3390/min10030273>
- [3] Basu A, Bickford ME, Deasy R (2016) Inferring tectonic provenance of siliciclastic rocks from their chemical compositions: a dissent. *Sediment Geol* 336:26–35
- [4] Bershaw J, Garzzone CN, Schoenbohm L, Gehrels G, Tao L (2012) Cenozoic evolution of the Pamir plateau based on stratigraphy, zircon provenance, and stable isotopes of foreland basin sediments at Oytay (Wuyitake) in the Tarim Basin (west China). *J Asian Earth Sci* 44:136–148.
- [5] Carver, R.E. Heavy-Mineral Separation. Chap. 18 in *Procedures in Sedimentary Petrology*; Carver, R.E., Ed.; John Wiley: Hoboken, NJ, USA, 1971.
- [6] Cawood PA, Hawkesworth CJ, Dhuime B (2012) Detrital zircon record and tectonic setting. *Geology* 40:875–878. <https://doi.org/10.1130/G32945.1>
- [7] Chaudhri, R.S. and Gill, G.T.S. (1980) Heavy Mineral Assemblage of the Siwalik. Group of Nepal Himalayas. *Journal of the Geological Society*, 22, 220-226.
- [8] Duarah, B.P., Das, P.K. and Goswami, D.C. (1998) Heavy Mineral Study from Modern Sediments of a Point Bar in the Puthimari River near Puthimari Village, Kamrup District, Assam. *Journal of the Indian Academy of Geoscience*, 41, 47-52.
- [9] Folk, R.L. (1980) The Provenance of Rutile. *Journal of Sedimentary Research*, 50, 485-488.
- [10] Hutton, C.O. Studies of heavy detrital minerals. *Bull. Geol. Soc. Am.* 1950, 61, 635–710.
- [11] Joshi, K.B., Banerji, U.S., Dubey, C.P. *et al.* Heavy minerals in provenance studies: an overview. *Arab J Geosci* 14, 1330 (2021). <https://doi.org/10.1007/s12517-021-07687-y>
- [12] Joshi KB, Ray S, Ahmad T, Manavalan S, Aradhi KK (2021) Geochemistry of meta-sediments from Neoproterozoic Shimla and Chail Group of Outer Lesser Himalaya: implications for provenance, tectonic setting and paleo-weathering conditions. *Geol J.* <https://doi.org/10.1002/gj.4183>
- [13] Kulm, M.D. (1988) Potential Heavy Mineral and Metal Placers on the Oregon Continental Shelf. *Marine Mining*, 7, 361-395.
- [14] Mange, M.A.; Wright, D.T. *Heavy Minerals in Use; Developments in Sedimentology Series*; Elsevier: Amsterdam, The Netherlands, 2007; Volume 58, 1283p.
- [15] Morton, A.C. (1985). Heavy Minerals in Provenance Studies. In: Zuffa, G.G. (eds) *Provenance of Arenites*. NATO ASI Series, vol 148. Springer, Dordrecht. https://doi.org/10.1007/978-94-017-2809-6_12
- [16] Rezaye, F., Shivanna and Rao, M.J. (2018) Studies on Geochemical and Heavy Mineral Characteristics of Sediments of Gosthani River Estuary. *Open Journal of Geology*, 8, 263-277. doi: 10.4236/ojg.2018.83017.



INTERNATIONAL
STANDARD
SERIAL
NUMBER
INDIA



INTERNATIONAL JOURNAL OF INNOVATIVE RESEARCH

IN SCIENCE | ENGINEERING | TECHNOLOGY

 9940 572 462  6381 907 438  ijirset@gmail.com



www.ijirset.com

Scan to save the contact details