

International Journal of Innovative Research in Science, Engineering and Technology

(A High Impact Factor, Monthly, Peer Reviewed Journal)

Visit: www.ijirset.com

Vol. 7, Issue 8, August 2018

Statistically Studing Average Seed Output and Seed Germination in *Daturastramonium L.*

Dr. Anubhuti Tiwari

Associate Professor, Dept. of Botany, SRKP Government College Kishangarh (Ajmer), India

ABSTRACT: Colonization strategies efficient dispersal and high survival capacity of offspring in new habitat, besides dispersal itself, depends on reproductive characteristics, such as germination, average seed output and vegetative spread as well as ecological demands of species are usually considered to determine colonisation success of species during succession. Colonization and establishment of herbaceous plants in semi-desert area is often a critical stage. Many plants grow along roadside areas of Jaipur during rainy season and make a green belt helping in reducing soil erosion, reducing dust pollution caused by vehicular smoke and also gives aesthetic value to the city. A study was conducted on naturally growing plant *Daturastramonium L.* as coloniser for the barren land. A field study was conducted around three selected sites of suburban area of Jaipur and data was collected. Ecophysiology, agrobotanical traits, seed size-number and reproductive capacity was calculated by taking statistical one way ANOVA F test.

KEYWORDS : Seed size, seed number, colonisation,, average seed output, reproductive capacity

I. INTRODUCTION

In India all sort of stray animals defaecate on roadsides Joshi and Swami (2007). Unfortunately, even children are often seen faecating on roadsides. All this faecal matter which contains harmful microorganisms sooner or later becomes part of sandy soil which gets easily blown into houses and pollutes many eatables and drinkables inside houses. There is also no check on people throwing garbage outside their houses which ultimately become part of the sandy roadsides and other bare areas. Colonization strategies efficient dispersal and high survival capacity of offspring in new habitat, besides dispersal itself, depends on reproductive characteristics, such as germination and vegetative spread as well as ecological demands of species are usually considered to determine colonisation success of species during succession (van der Valk 1992). Some of the traits important for colonisation and establishment of species are average seed output, seed size, plant height, specific leaf area and leaf dry weight Weiher *et al.* (1999). Austrheim *et al.* (2005) pointed out a very important characteristic of colonising species which refers to the introduction and colonisation of certain species into previously uninhabited areas into a country from a foreign country with unusual success in the new physical/biological environment designated as genetic revolution. According to him ecologically, the most stressful condition in the semi-arid region of Rajasthan is drought which is related to scanty rainfall and soil remains sandy for most of the year. Germination or emergence percentage has been shown to be closely related to seed weight (Weis, 1982 and Zhang and Maun, 1991). Higher germination and/or emergence force in natural populations by virtue of heavier seeds is an asset in that seeds can emerge through a covering soil layer relatively more easily than lighter seeds. Bhat (1973) showed an inverse relationship between seed weight and seed coat impermeability, which also seems to be true for *Rhyncosia minima*. Eriksson and Jakobsson (1998) have emphasised seed size as well as seed weight as a key functional trait with respect to colonisation. Most of the ecologists believe that ecological traits do have a strong relationship with plant frequencies and that species richness is related to habitat variation, obviously with a low stress level (Grime, 1979)

International Journal of Innovative Research in Science, Engineering and Technology

(A High Impact Factor, Monthly, Peer Reviewed Journal)

Visit: www.ijirset.com

Vol. 7, Issue 8, August 2018

II. STUDY MATERIAL

*Daturastramonium*L belongs to family Solanaceae is commonly called as Jimson weed, Angel's Trumpet, Devil's trumpet, Jamestown weed, Loco weed, Moonflower, Thorn-apple, Mad apple in English, Dhatura in Hindi.

Distribution-*Daturastramonium L* is found throughout the warm temperate parts of the world. In India *D. stramonium* is common in North Western Himalayas, Central and Western India (Braun *et al.*, 1991). It is usually found growing in disturbed areas, waste lands, burn yards, along road sides, rail tracks, rocky open areas, scrub vegetation and grassland.

Botanical Description - The plant is annual or short-lived perennial erect herb up to 2.0 m tall. It grows in winters and flowering and fruiting occurs till late March. Stem is pseudo- dichotomously branched. It is sparsely hairy to glabrous with purple in colour. Leaves are alternate, simple, minutely hairy. The leaf blade is rhombic to ovate with coarsely doubly lobed margins and acute apex. Inflorescence is axillary with pentamerous flowers. Calyx is angular with five longitudinal ribs while corolla is trumpet-shaped, six to eight cm long and white in colour. The stamens are five in number and arranged adnate to the corolla tube. Filaments are short and thick with three to four mm long anthers. Ovary is superior with slender, 3.5–7.0 cm long style and two lobed stigma. Fruit is egg shaped and many seeded capsule with four valves and rigid spines. Seeds are kidney shaped and flattened with dark brown to black in colour.

Chemical Composition and Medicinal Importance -The concentration of total alkaloids in the leaves of *D. stramonium L* is 0.2–0.5%, hyoscyamine being the major compound and scopolamine (hyoscine), apoatropine, tropine, belladonnine and hyoscyamine compounds occur in minor quantity (Berkovet *et al.*, 2006). Zagari (1992) reported that *D. stramonium*L is very useful in curing asthma, cough, tuberculosis and bronchitis by smoking 'asthma cigarettes' made of the dried leaves, roots or flowers. *Datura* fruit is a specific remedy for phlegmatic and bilious types of malarial fever. This herb is also a valuable remedy for heart disorders, as it relieves cardiac pains, distress, palpitations and aortic disorders (Berne and Levy, 1998 and Heath *et al.*, 1999).

A decoction of leaves is given as a sedative to mental and schizophrenic patients and in treating Parkinson's disease. The herb is highly beneficial in halting the secretion of the breast milk in case of the unfortunate death of a newly born baby. Along with this *Datura* is useful in curing the impotency and promotes health and vigour and absolute fitness of the body (Monseeset *et al.*, 2001). The leaves are applied as a dressing to cure rheumatic pain, swellings, patchy baldness, wounds, gout, burns, ingrown toe-nails, fungal infections, tumours, ulcers and piles (Spring, 1989). The flowers and seeds of *D. stramonium*L are hallucinogenic and known to have poisonous effects like raving, insensibility and even death. Food and Drug Administration (FDA) has determined it to be unfit for human consumption.

III. STUDY SITE AND METHODOLOGY

This study site is natural population of plants growing along roadsides and open land in suburban areas of Jaipur at three sites of different localities were selected for experimental study for statistical analysis of total number of seeds per plant were studied as per the methods given by Misra (1968) and Pandeya, Puri and Singh (1968). Mature seeds were collected carefully before they were shed, if the dispersal is quick, the fruits as well as plants were tied with a cloth to trap the seeds to avoid contamination and mixing with other plant material. The collected seeds were used for calculating seed germination percentage, average seed output and reproductive capacity. Average seed output of plants containing more than one seed was calculated as given by Salisbury (1942).

International Journal of Innovative Research in Science, Engineering and Technology

(A High Impact Factor, Monthly, Peer Reviewed Journal)

Visit: www.ijirset.com

Vol. 7, Issue 8, August 2018

$$\text{Average seed output} = \text{Number of seeds per fruit (mean)} \times \text{number of fruits per plant (mean)} \pm \text{standard error of means}$$

Community structure of an area is expressed by measuring its frequency using Quadrat method given by Oosting (1958). Twenty quadrats (1mx1m) were placed at each site for calculation of frequency class given by Raunkiaer (1934).

$$\text{Frequency \%} = \frac{\text{Number of quadrats in which species occurred}}{\text{Total number of quadrats studies}} \times 100$$

$$\text{Reproductive capacity} = \frac{\text{Average seed output} \times \text{Seed germination\%}}{100}$$

Observations

The number of seeds per fruit of *Daturastranomium L* is highest at site C 476.4±16.92 with one way ANOVA F ratio 0.1527* Average seed output was 38242.2 at site C (highest) of *DaturastranomiumL* and reproductive capacity is highest at site C 25622.27 on all the three sites. The seed germination percentage was recorded in *Daturastranomium L* at 70% at site A (highest) one way ANOVA F ratio 15.0815NS The data of seed characters like seed size and number of seeds/plant, average seed output, seed germination percentage, reproductive capacity and frequency are recorded in (Table 1). Seed length (mm) was 1.36±0.15(highest) at site A with one way ANOVA F ratio 1.6058* and seed width(mm) 0.69±0.15 (highest) at site B with one way ANOVA F ratio 1.0385* The statistical analysis showed that there was no significant difference between and among all the three sites. Flowering and Fruiting is dependent upon the vegetative growth and on the availability and proper balance of mineral nutrients. More vegetative growth affects the reproductive growth of a plant. *DaturastranomiumL* belong to Frequency Class 'E' and is common and abundant plant and could be a good coloniser

Table 1: Showing Variation in Seed Characters and Reproductive Capacity of *Daturastramonium* from all the Three Sites (values are mean of 50 readings).

Parameters	Site A (mean±SD)	Site B (mean±SD)	Site C (mean±SD)
Seed Length(mm)	1.22±0.19	1.34±0.22	1.36±0.15
Seed Width(mm)	0.61±0.13	0.69±0.15	0.62±0.11
Number of Seeds/Fruit	433±33.26	423.8±56.46	476.4±16.92
Average Seed Output	36423.96	35624.63	38242.2
Seed Germination (%)	70	67	67
Reproductive Capacity	25496.77	23868.5	25622.27
Frequency Class	E	E	E

International Journal of Innovative Research in Science, Engineering and Technology

(A High Impact Factor, Monthly, Peer Reviewed Journal)

Visit: www.ijirset.com

Vol. 7, Issue 8, August 2018

NS= Not significant * = Significant

Analysis of variance:

F-ratio:

- i. Seed Length =1.6058*
- ii. Seed Width=1.0385*
- iii. Number of Seeds/Fruit =0.1527*
- iv. Seed Germination =15.0815^{NS}

IV. DISCUSSION AND RESULT

Frequency –dependent or game-theoretical-strategy-mixture models, state that more establishment opportunities must be reached by small-seeded species than by large-seeded species. Hence a negative correlation between seed size and species abundance was noticed by Rees and Westoby, (1997). On the contrary Mitchley and Grubb (1986) found positive correlation between seed size and abundance. Community pattern also shows a negative correlation between seed size and abundance (Pake and Venable, 1996). **Seed Size and Seed Number-** Since, environment resources necessary to reproduce are limited hence, plant experience a tradeoff involving allocation of energy to dispersal (higher seed number) or offspring survival (higher seed size) the so called seed size/number trade off (SSNT) gives rise to a competition-colonization trade-off for annual plants. Both seed size and number are regarded as ecologically important life-history traits and provide increased fitness in various environments (Stamp, 1990 and Saeed and Shaukat, 2000). Seed size seems to be an important character in plant fitness. SSNT explaining the relation of seed size and seed number (Aiken and Springer., 1995). Both seed size and number are regarded as ecologically important life-history traits and provide increased fitness in various environments (Westoby *et al.*, 1996). Bowers *et al.*, (2004), explained that larger seeds have greater recruitment, greater percent germination or emergence success as compared to smaller seeds. Relatively low seed number was sufficient for maximal fruit set. Seeds are well known to be a rich source of plant growth regulators (Hedden and Hoad, 1985).

Reproductive Capacity- Reproductive capacity of a plant is another critical aspect of plant reproduction. According to Salisbury (1942) reproductive capacity of many plants is extremely great and there are large differences between species, these differences are due to selective pressures (Harper and White 1974) These findings agree with our observations that in *Datura stramonium L* have high average seed output and greater reproductive capacity.

Frequency- Mitchley and Grubb (1986) found positive correlation between seed size and abundance. This finding is similar to our observations that *Datura stramonium L* belong to Frequency Class 'E'. is the common and abundant plant and could be a good coloniser.

Industrialization and Urbanization of cities especially metropolitan cities are facing fast growth in automobile number which is the major cause of dust pollution. These dust when blown carries with it large amount of pollutant into our houses and are cause of many air born diseases. Hence, covering the roadsides by vegetation is the need of hour. A comparative account of all the data among all the selected sites for consideration as best coloniser our findings suggest that *Datura stramonium L* is a coloniser species. Hence we can recommend *Datura stramonium L* to be distributed for plantation to cover the barren roadside areas of all the three sites of Jaipur. This will help to reduce soil erosion, noise pollution, dust pollution caused by vehicles along roadsides and will help to enhance the greenery of the area.

International Journal of Innovative Research in Science, Engineering and Technology

(A High Impact Factor, Monthly, Peer Reviewed Journal)

Visit: www.ijirset.com

Vol. 7, Issue 8, August 2018

REFERENCES

1. Aiken, G.E. and Springer, T.L. 1995. Seed size distribution, germination and emergence of six switch grass cultivars. *Jour. Range Manag.* 48: 455-458.
2. Austrheim, G., Evju, M. and Myrsetrud, A. 2005. Herb abundance and life history traits in two contrasting alpine habitats in southern Norway. *Plant Ecol.* 179: 217-229.
3. Bowers, J.E., Turner, R.M. and Burgess, T.L. 2004. Temporal and spatial patterns in emergence and early survival of perennial plants in the Sonoran Desert. *Plant Ecol.* 172:107-119.
4. Bradshaw, A. D. 1987a . Functional Ecology = Comparative Ecology? *Functional Ecology* 1: 71-71.
5. Bullock, J.M. 2000. Gaps and seedling colonization. In : Fenner M. (ed.), *Seeds : The Ecology of Regeneration in Plant Communities*. CABI Publishing, Wallingford. pp. 375-396.
6. Baker, H.G. 1958. Studies in reproduction biology of West African Rubiaceae. *Jour. West Afr. Sci. Assoc.*4: 9-24.
7. Chepil, W. S. 1949. Wind erosion control with shelterbelts in North China. *Agron. Jour.* 41: 127-129.
8. Chopra, R.N., Nayar, S.L. and Chopra, I.C. 1986. *Glossary of Indian Medicinal Plants*. Reprint Edition, Publication and Information Directorate, Council of Scientific and Industrial Research, New Delhi.
9. Elizabeth, P.H. 2005. Phytoremediation. *Ann. Rev. Plant Biol.* 56: 15-39.
10. Eriksson, O. 1997. Colonization dynamics and relative abundance of three plant species *Antennaria dioica*, *Hieracium pilosella* and *Hypochoeris maculata* in semi natural grasslands. *Ecography* 20: 559-568.
11. Eriksson, O. and Jakobsson, A. 1998. Abundance, distribution and life histories of grassland plants: a comparative study of 81 species. *Jour. Ecol.* 86: 922-933.
12. Grime, J.P. 1979. *Plant Strategies and Vegetation Process*. John Wiley and Sons, Chichester.
13. Gross, K. L. and Werner, P.A. 1982. Colonizing abilities of "biennial" plant species in relation to ground cover: implications for their distributions in a successional sere. *Ecology* 63: 921-931.
14. Harper, J.L. and J. White. 1974. Demography of plants. *Ann. Rev. Ecol. Syst.* 5: 419-463.
15. Hendrix, S.D., Nielsen, E., Nielsen, T. and Schutt, M. 1991. Are seedlings from small seeds always inferior to seedlings from large seeds? effects of seed biomass on seedling growth in *Pastinaca sativa* L. *New Phytol.* 119:299-305.
16. Jakobsson, A. and Eriksson, O. 2000. A comparative study of seed number, seed size, seedling size and recruitment in grassland plants. *Oikos* 88: 494-502.
17. Joshi, P.C. and Swami, A. 2007. Physiological responses of some tree species under roadside automobile pollution stress around city of Haridwar, India. *Environmentalist* 27: 365-374.
18. Joshi, P.C. and Swami, A. 2009. Air pollution induced changes in the photosynthetic pigments of selected plant species. *Jour. Environ. Biol.* 30: 295-298.
19. Kirkpatrick, M. 1984. Demographic models based on size, not age, for organisms with indeterminate growth. *Ecology* 65:1874-1884.
20. Kirtikar, K.R. and Basu, B.D. 1980. *Indian Medicinal Plants*. Lalit Mohan Basu, Allahabad, India. 3:1774-1777.
21. Leishman, M. 1999. How well do plant traits correlate with establishment ability? Evidence from a study of 16 calcareous grassland species. *New Phytol.* 141: 487-496.
22. Leishman, M.R. 2001. Does the seed size/number trade-off model determine plant community structure? An assessment of the model mechanisms and their generality. *Oikos* 93: 294-302.
23. Misra, R. 1968. *Ecology Work Book*, Calcutta, Bombay, New Delhi.
24. Mitchley, J. and Grubb, P. J. 1986. Control of relative abundance of perennials in chalk grassland in southern England. *Jour. Ecol.* 74: 1139-1166.



ISSN(Online): 2319-8753
ISSN(Print): 2347-6710

International Journal of Innovative Research in Science, Engineering and Technology

(A High Impact Factor, Monthly, Peer Reviewed Journal)

Visit: www.ijirset.com

Vol. 7, Issue 8, August 2018

25. Oosting, H.J. 1958. The Study of Plant Communities. 2 edition. W.H. Freeman and Co., San Francisco. pp 440.
26. Pandeya, S.C., Puri,G.S. and Singh,J.S.1968. Research Methods in Plant Ecology. Bombay-Calcutta-NewDelhi-Madras-Lucknow-Bangalore- London – NewYork
27. Raunkiaer, C. 1934. The Life Forms of Plants and Statistical Plant Geography. Clarendon Press, Oxford, England.
28. Saeed, S. and Shaukat, S. S. 2000. Effect of seed size on germination, emergence, growth and seedling survival of *Senna occidentalis* Link.Pak. Jour. Biol. Sci. 3: 292-295.
29. Salisbury, E. J. 1942. The Reproductive Capacity of Plants.G.Bell and Sons Ltd., London.pp 244.
30. Schaal, B.A. 1980. Reproductive capacity and seed size in *Lupines texensis*. Amer. Jour. Bot. 67: 703-709.