

# Effect of Cultivar & Weed Management on Weed Dynamics and Yield of Black Gram (*Vigna mungo*)

Dr. Shrish Kumar Singh<sup>1</sup>, Jay Prakash Pandey<sup>2</sup>

Associate Professor & Head of Department-Agronomy, TDPG College, Jaunpur, Uttar Pradesh, India<sup>1</sup>

M.Sc (Ag) Agronomy Student, TDPG College, Jaunpur, Uttar Pradesh, India<sup>2</sup>

**ABSTRACT:** Blackgram [*Vigna mungo*] is an important pulse crop in India. It is grown during summer as well as kharif season. Frequent irrigations during summer season and monsoon rains during kharif season result in lot of weed population and weed growth in this crop. Unchecked weeds have been reported to cause a considerable reduction in the grain yield of blackgram, which in case of summer blackgram could be 46.0- 53.0% (Vaishya et al., 2003; Bhandari et al., 2004; Kumar and Tewari, 2004a), whereas in kharif blackgram the losses could be 43.2-64.1% (Chand et al., 2004; Rathi et al., 2004). The critical period of crop-weed competition in case of summer blackgram is 10-40 days after sowing (Kumar and Tewari, 2004b) which in certain situations could be 25-35 days after sowing (Randhawa et al., 2002). Therefore, removal of weeds at appropriate time using a suitable method is essential to obtain high yields of blackgram. In blackgram, weeds could be controlled by hand weedings (Chand et al., 2004). However, hand weeding is laborious, time consuming, costly and tedious. Moreover, many times labour is not available at the critical period of weed removal. Furthermore, weather conditions (rains) do not permit timely hand weeding due to wet field conditions. Delayed removal of weeds is not as effective in controlling weeds and obtaining higher yields as the timely removal of weeds. Under these conditions, use of herbicides offers an alternative for possible effective control of weeds. Therefore, in the present studies, effect of various herbicides was compared with hand weedings and untreated check for evaluating weed control efficacy obtaining high yields of blackgram grown during summer and kharif seasons.

**KEYWORDS:** *Vigna mungo*, weed, management, dynamics, cultivar, yield, weather, season

## I. INTRODUCTION

Field experiments were conducted in the summer seasons for four years (2002, 2003, 2004 and 2005) and in kharif seasons for three years (2002, 2003 and 2005) at the Punjab Agricultural University, Ludhiana to study the effect of various weed control treatments on weeds, grain yield of blackgram and the economics involved.




Five treatments viz., (i) pendimethalin 0.75 kg/ha (pre-emergence), (ii) pendimethalin 0.45 kg/ha (pre-emergence)+hand weeding (HW) 25 days after sowing (DAS), (iii) fluchloralin 0.675 kg/ha (pre-plant incorporation), (iv) two HW 25 & 40 DAS and (v) weedy check were arranged in a randomized complete block design having three replications in 2002, 2003 and 2004 and four replications in 2005. Herbicides were sprayed with a knap sack sprayer fitted with a flat fan nozzle using 500 litres of water per hectare. Pendimethalin was applied either immediately after sowing or one day after sowing, whereas fluchloralin was sprayed just prior to sowing.



In case of hand weeding, weeding was done using a khurpa. In summer season, variety Mash 414 was sown on 14 April, 2 April, 6 April and 28 March in 2002, 2003, 2004 and 2005, respectively, in rows 22.5 cm apart using a seed rate of 50 kg/ha. In kharif season, variety Mash 338 was sown on 10 July, 10 July and 7 July in 2002, 2003 and 2005, respectively, in rows 30 cm apart using a seed rate of 20 kg/ha. The crop was raised as per the PAU recommendations (PAU, 2001).

### Cultural Weed Control



Data on dry matter of weeds were recorded at harvest. Weed control efficiency (WCE) was calculated. Economics of different weed control treatments was also worked out by taking the selling price of blackgram as Rs. 1400/q.

## II. RESULTS AND DISCUSSION

The major weed flora included *Trianthema portulacastrum*, *Eleusine aegyptiacum*, *Digitaria sanguinalis* and *Cyperus rotundus*. Pendimethalin at both the rates of application showed good control of *T. portulacastrum*, *E. aegyptiacum* and *D. sanguinalis*. Kumar and Tewari (2004a) reported very good control of *Trianthema monogyna* by pendimethalin as this herbicide does not allow the emergence and growth of weed seedlings because of reduced cell division and cell elongation. Fluchloralin was not very effective for the control of *T. portulacastrum*. *C. rotundus* remained uncontrolled with both the herbicides. Weedy check had the highest weed dry matter, which, on an average, was 23.60 and 20.90 q/ha during summer and kharif season, respectively .

All other treatments were very effective in controlling weeds thereby resulting in significantly lower dry matter of weeds compared with the weedy check. Among these, the treatment of two HW 25 & 40 DAS recorded the lowest weed dry matter in summer as well as kharif season. The mean values show that two HW done 25 & 40 DAS had the highest weed control efficiency (85.01 and 84.89%) during both the seasons , which was closely followed by pendimethalin 0.45 kg/ha+HW 25 DAS (84.81 and 78.28%) and pendimethalin 0.75 kg/ha (79.12 and 77.52%). Fluchloralin 0.675 kg/ha recorded lowest weed control efficiency.

## MECHANICAL METHOD OF WEED CONTROL



**Mowing**



**Spudding**



**Soil solarisation**



**Burning**

Bhandari et al. (2004) reported effective control of weeds with pendimethalin 1.0-2.0 kg/ha, fluchloralin 0.5-1.5 kg/ha and pendimethalin 1.0 kg/ha + HW 25 DAS. Two HW done 25 & 40 DAS recorded the highest grain yield in all the years except in 2003 when pendimethalin 0.45 kg/ha+HW 25 DAS produced highest grain yield.

### III. CONCLUSION

On the basis of mean data, two HW 25 & 40 DAS recorded the highest (11.95 and 11.76 q/ha) grain yield in summer and kharif seasons, which was followed by pendimethalin 0.45 kg/ha+HW 25 DAS (11.75 and 10.76 q/ha). Pendimethalin 0.5 kg/ ha+HW 30 DAS provided effective control of weeds and high grain yield of kharif season blackgram (Rathi et al., 2004). In kharif blackgram, two HW done 20 and 40 DAS provided as high grain yield as the weedfree treatment (Chand et al., 2004). Pendimethalin 1.0- 2.0 kg, pendimethalin 1 kg/ha+HW 25 DAS and fluchloralin 0.5-1.5 kg/ha have been reported to provide quite good grain yield of summer blackgram (Bhandari et al., 2004).



Similar results have also been reported by Vaishya et al. (2003). As compared to two HW 25 & 40 DAS, the unchecked weeds caused, on an average, 41.2 and 41.6% reduction in grain yield during summer and kharif seasons, respectively. Similar grain yield losses due to weeds have been reported by other researchers in summer blackgram (Bhandari et al., 2004) and kharif blackgram (Chand et al., 2003; Chand et al., 2004). Gross returns were the highest in case of 2 HW 25 & 40 DAS during both the seasons (Table 4). Net returns were the highest with pendimethalin 0.45 kg/ha + HW 25 DAS, closely followed by pendimethalin 0.75 kg/ha in summer season and with fluchloralin 0.675 kg/ha in kharif season. Weedy check though involved the lowest cost of cultivation yet it provided the lowest net returns

#### REFERENCES

- 1) Bhandari, V., B. Singh, J. S. Randhawa and J. Singh. 2004. Relative efficacy and economics of integrated weed management in blackgram under semi-humid climate of Punjab. *Ind. J. Weed Sci.* 36 : 276-277.
- 2) Chand, R., N. P. Singh and V. K. Singh. 2003. Effect of weed management practices on productivity of late planted urdbean during kharif season. *Ind. J. Pulses Res.* 16 : 163-164.
- 3) Chand, R., N. P. Singh and V. K. Singh. 2004. Effect of weed control treatments on weeds and grain yield of late sown urdbean (*Vigna mungo* L.) during kharif season. *Ind. J. Weed Sci.* 36 : 127-128.
- 4) Kumar, A. and A. N. Tewari. 2004a. Efficacy of pre- and post-emergence herbicides in summer blackgram (*Vigna mungo* L.). *Ind. J. Weed Sci.* 36 : 73-75.
- 5) Kumar, A. and A. N. Tewari. 2004b. Crop-weed competition studies in summer sown blackgram (*Vigna mungo* L.). *Ind. J. Weed Sci.* 36 : 76-78. PAU. 2001.
- 6) Package of Practices for Crops of Punjab, Kharif 2001. Punjab Agricultural University, Ludhiana.
- 7) Randhawa, J. S., J. S. Deol, V. Sardana and J. Singh. 2002. Crop-weed competition studies in summer blackgram (*Phaseolus mungo*). *Ind. J. Weed Sci.* 34 : 299-300.
- 8) Rathi, J. P. S., A. N. Tewari and M. Kumar. 2004. Integrated weed management in blackgram (*Vigna mungo* L.). *Ind. J. Weed Sci.* 36 : 218- 220.
- 9) Vaishya, R. D., B. K. Srivastava and G. Singh. 2003. Integrated weed management in summer urdbean. *Ind. J. Pulses Res.* 16 : 161-162.