

Common fixed point theorems for self maps on a fuzzy metric space involving integral type in equalities

K.P.R.Sastry¹, G.A.Naidu², N.Umadevi³

8-28-8/1, Tamil Street, Chinna Waltair, Visakhapatnam-530017, India¹

Department of Mathematics, Andhra university, Visakhapatnam-530003, India²

Department of Mathematics, Raghu Engineering College, Visakhapatnam-531 162, India³

Abstract: Malhotra [7] proved a common fixed point theorem in fuzzy metric spaces for occasionally weakly compatible mappings with integral type in equality by reducing its minimum value. In this paper we prove a common fixed point theorem in fuzzy metric spaces for occasionally weakly compatible mappings with integral type inequality involving some special type of Lebesgue integrable functions.

Key words: Fuzzy metric space, occasionally weakly compatible (owc) mappings, common fixed point.

Mathematical Subject Classification (2010): 47H10, 54H25

I. INTRODUCTION

Fuzzy set was introduced by Zadeh [15]. Kramosil and Michalek [6] introduced the notion of a fuzzy metric space, George and Veermani [4] modified the notion of fuzzy metric spaces with the help of continuous t-norms. Grabiec [3], Subramanyam [13], Vasuki [14], Pant and Jha [9] obtained some analogous results proved by Balasubramaniam [1] et al. Subsequently, it was developed extensively by many authors and used in various fields. Jungck [5] introduced the notion of compatible maps for a pair of self maps. George and Veermani [4], Sessa [11] initiated the tradition of improving commutative condition in fixed point theorems by introducing the notion of weak commuting property. Further Jungck and Rhoades [5] gave a more generalized condition defined as compatibility in metric spaces. Jungck and Rhoades [5] also introduced the concept of weakly compatible maps. Malhotra [7] proved a common fixed point theorem in fuzzy metric spaces for occasionally weakly compatible mappings with integral type in equality by reducing its minimum value. In this paper we prove a common fixed point theorem for selfmaps on fuzzy metric spaces for occasionally weakly compatible mappings with integral type inequality involving some special type of Lebesgue integrable functions (Definition [2.1]).

Definition 1.1 : (Zadeh.L.A. [14]) A fuzzy set A in a nonempty set X is a function with domain X and values in $[0,1]$.

Definition 1.2: (Schweizer.B. and Sklar. A. [9]) A function $*$: $[0,1] \times [0,1] \rightarrow [0,1]$ is said to be a continuous t-norm if $*$ satisfies the following conditions:

For $a, b, c, d \in [0,1]$

- (i) $*$ is commutative and associative
- (ii) $*$ is continuous
- (iii) $a * 1 = a$ for all $a \in [0,1]$
- (iv) $a * b \leq c * d$ whenever $a \leq c$ and $b \leq d$

Definition 1.3: (Kramosil. I. and Michalek. J. [5]) A triple $(X, M, *)$ is said to be a fuzzy metric space (FM space, briefly) if X is a nonempty set, $*$ is a continuous t-norm and M is a fuzzy set on $X^2 \times [0, \infty)$ satisfying the following conditions:

For $x, y, z \in X$ and $s, t > 0$.

- (i) $M(x, y, t) > 0, M(x, y, 0) = 0$
- (ii) $M(x, y, t) = 1$ if and only if $x = y$.
- (iii) $M(x, y, t) = M(y, x, t)$
- (iv) $M(x, y, t) * M(y, z, s) \leq M(x, z, t + s)$
- (v) $M(x, y, \cdot): [0, \infty) \rightarrow [0, 1]$ is continuous.

Then M is called a fuzzy metric space on X .

The function $M(x, y, t)$ denotes the degree of nearness between x and y with respect to t .

Definition 1.4:(Jungck.G. and Rhoades.B.E. [4]) Two self mappings f and g of a fuzzy metric space $(X, M, *)$ are called compatible if $\lim_{n \rightarrow \infty} M(fg x_n, gf x_n, t) = 1$ whenever $\{x_n\}$ is a sequence in X such that $\lim_{n \rightarrow \infty} f x_n = \lim_{n \rightarrow \infty} g x_n = x$ for some $x \in X$.

Definition 1.5:(Cho.Y.J.[1]) Two self mappings f and g of a fuzzy metric space $(X, M, *)$ are called reciprocally continuous on x if $\lim_{n \rightarrow \infty} f x_n = f x$ and $\lim_{n \rightarrow \infty} g f x_n = g x$ whenever $\{x_n\}$ is a sequence in X such that $\lim_{n \rightarrow \infty} f x_n = \lim_{n \rightarrow \infty} g x_n = x$ for some $x \in X$.

Definition 1.6: (George.A. and Veeramani.P. [3]) Let $(X, M, *)$ be a fuzzy metric space.

Then,

- (i) A sequence $\{x_n\}$ in X is said to be convergent to a point $x \in X$ if $\lim_{n \rightarrow \infty} M(x_n, x, t) = 1 \forall t > 0$.
- (ii) A sequence $\{x_n\}$ in X is called a Cauchy sequence if $\lim_{n \rightarrow \infty} M(x_{n+p}, x_n, t) = 1 \forall t > 0$ and $p = 1, 2, \dots$
- (iii) An FM –space in which every Cauchy sequence is convergent is said to be complete.

Definition 1.7: (Malhotra.S.K.Naveen Verma and Ravindra Sen [6]) Two self maps f and g of a set X are occasionally weakly compatible (owc) iff there is a point x in X which is a coincidence point of f and g at which f and g commute.

Lemma: 1.8:(Mishra.S.N. and Sharma.N.[7]) Suppose $(X, M, *)$ is a fuzzy metric space. If for all $x, y \in X, t > 0$ and for a number $k \in (0, 1), M(x, y, kt) \geq M(x, y, t)$ then $x = y$.

Lemma: 1.9:(Jungck.G. and Rhoades.B.E. [4]) Let X be a set and f, g owc self maps of X . If f & g have a unique point of coincidence, $w = f x = g x$, then w is the unique common fixed point of f and g .

Notation 1.10: Let $\Phi = \{\phi; R^+ \rightarrow R\}$ such that (1) ϕ is Lebesgue integrable

$$(2) \epsilon > 0 \Rightarrow \int_0^\epsilon \phi(t)dt > 0$$

Malhotra et.al.[7] proved a common fixed point theorem for four self maps on a fuzzy metric space which satisfy an inequality involving a function $\phi \in \Phi$.

Theorem 1.11: (Malhotra et.al.[7]) Let $(X, M, *)$ be a complete fuzzy metric space and let F, G, S and T are self-mapping of X . Let the pairs $\{F, S\}$ and $\{G, T\}$ be owc . If there exists $q \in (0,1)$ such that

$$\int_0^{M(Fx, Gy, qt)} \phi(t)dt \geq \int_0^{\min\{M(Sx, Ty, t), M(Sx, Fx, t), M(Gy, Ty, t), M(Fx, Ty, t), M(Gy, Sx, t)\}} \phi(t)dt$$

for all $x, y \in X$ and for all $t > 0$, then there exists a unique point $w \in X$ such that $Fw = Sw = w$ and a unique point $z \in X$ such that $Gz = Tz = z$ Moreover , $z = w$, so that there is a unique common fixed point of F, G, S and T .

II. MAIN RESULT

In this section we introduce a special class of Lebesgue integrable function and use this notion to prove a fixed point theorem.

Definition 2.1: Let $\Psi = \{\psi: [0,1] \rightarrow [0,1]\}$ is Lebesgue integrable such that

$$\int_\alpha^\beta \psi(t)dt > 0 \text{ for } \alpha < \beta.$$

Theorem 2.2: Let $(X, M, *)$ be a complete fuzzy metric space and let F, G, S and T be self-mappings of X . Let the pairs (F, S) and (G, T) be owc . Suppose there exists $\psi \in \Psi$ and $q \in (0,1)$ such that

$$\int_0^{M(Fx, Gy, qt)} \psi(s)ds \geq \int_0^{\min\{M(Sx, Ty, t), M(Sx, Fx, t), M(Gy, Ty, t), M(Fx, Ty, t), M(Gy, Sx, t)\}} \psi(s)ds \quad (2.2.1)$$

for all $x, y \in X$ and for all $t > 0$.

Then F, G, S and T have a unique common fixed point.

Proof:

Since the pairs (F, S) and (G, T) are owc, there exist $x, y \in X$ such that $Fx = Sx$ and $Gy = Ty$. Now we claim that $Fx = Gy$.

By (2.2.1)

$$\begin{aligned} \int_0^{M(Fx, Gy, qt)} \psi(s)ds &\geq \int_0^{\min\{M(Sx, Ty, t), M(Sx, Fx, t), M(Gy, Ty, t), M(Fx, Ty, t), M(Gy, Sx, t)\}} \psi(s)ds \\ &= \int_0^{\min\{M(Fx, Gy, t), M(Fx, Fx, t), M(Gy, Gy, t), M(Fx, Gy, t), M(Gy, Fx, t)\}} \psi(s)ds \\ &= \int_0^{\min\{M(Fx, Gy, t), 1, 1, M(Fx, Gy, t), M(Gy, Fx, t)\}} \psi(s)ds \end{aligned}$$

$$\Rightarrow \int_0^{M(Fx, Gy, qt)} \psi(s)ds \geq \int_0^{M(Fx, Gy, t)} \psi(s)ds$$

$$\Rightarrow \int_0^{M(Fx, Gy, t)} \psi(s)ds \geq \int_0^{M(Fx, Gy, qt)} \psi(s)ds \geq \int_0^{M(Fx, Gy, t)} \psi(s)ds$$

$$\begin{aligned} \Rightarrow \int_0^{M(Fx, Gy, qt)} \psi(s) ds &= \int_0^{M(Fx, Gy, t)} \psi(s) ds = \int_0^{M(Fx, Gy, qt)} \psi(s) ds + \int_{M(Fx, Gy, qt)}^{M(Fx, Gy, t)} \psi(s) ds \\ \Rightarrow \int_{M(Fx, Gy, qt)}^{M(Fx, Gy, t)} \psi(s) ds &= 0 \\ \Rightarrow M(Fx, Gy, qt) &= M(Fx, Gy, t) \quad (\because \psi \in \Psi) \end{aligned}$$

Hence it follows that $M(Fx, Gy, qt) = M(Fx, Gy, t)$ for every $t > 0$.

Consequently $Sx = Fx = Gy = Ty$ (2.2.2)

Suppose that $\exists z \in X \ni Fz = Sz$. Then following the above argument we get that

$$Fz = Sz = Gy = Ty.$$

$$\therefore Fx = Fz$$

$$\text{Write } w = Fx = Sx.$$

$$\text{Since } (F, S) \text{ is owc, } Fx = Sx \Rightarrow FSx = SFx \Rightarrow Fw = Sw$$

Now replacing x by w in (2.1.1) following the argument mentioned above and we get

$$Fw = Sw = Gy = Ty.$$

$$\therefore Fw = Sw = Gy = Ty = Fx = Sx.$$

$$\therefore Fw = Fx = w = Sx = Sw.$$

$$\Rightarrow Fw = w = Sw.$$

Hence w is a common fixed point of F and G .

From (2.2.2) we have $Gy = Ty = w$, following the above argument follows that $Gw = Tw = w$.

Thus w is a common fixed point of F, S, G and T .

Uniqueness : Let $w \& v$ be $\ni Fw = Sw = Gw = Tw = w$ and $Fv = Sv = Gv = Tv = v$.

By (2.2.1)

$$\begin{aligned} \int_0^{M(Fw, Gv, qt)} \psi(s) ds &\geq \int_0^{\min \{M(Sw, Tv, t), M(Sw, Fw, t), M(Gv, Tv, t), M(Fw, Tv, t), M(Gv, Sw, t)\}} \psi(s) ds \\ \int_0^{M(w, v, qt)} \psi(s) ds &\geq \int_0^{\min \{M(w, v, t), 1, M(w, v, t), M(v, w, t)\}} \psi(s) ds = \int_0^{M(w, v, t)} \psi(s) ds \\ \Rightarrow \int_0^{M(w, v, qt)} \psi(s) ds &\geq \int_0^{M(w, v, t)} \psi(s) ds. \\ \Rightarrow \int_0^{M(w, v, qt)} \psi(s) ds &\geq \int_0^{M(w, v, qt)} \psi(s) ds \geq \int_0^{M(w, v, t)} \psi(s) ds. \\ \Rightarrow \int_0^{M(w, v, t)} \psi(s) ds &= \int_0^{M(w, v, qt)} \psi(s) ds \\ \Rightarrow M(w, v, qt) &= M(w, v, t) \quad \forall t > 0. \\ \Rightarrow w &= v. \end{aligned}$$

Hence common fixed point of F, G, S and T is unique.

Theorem 2.3: Let $(X, M, *)$ be a complete fuzzy metric space and let F, G, S and T be self-mapping of X . Let the pairs (F, S) and (G, T) be owc . If there exists $q \in (0, 1)$ such that

$$\int_0^{M(Fx, Gy, qt)} \psi(s) ds \geq \int_0^{\phi(\min \{M(Sx, Ty, t), M(Sx, Fx, t), M(Gy, Ty, t), M(Fx, Ty, t), M(Gy, Sx, t)\})} \psi(s) ds \quad (2.3.1)$$

for all $x, y \in X$ and $\phi: [0, 1] \rightarrow [0, 1]$ such that $\phi(t) \geq t$ for all $0 < t < 1$.

Then F, G, S and T have a unique common fixed point.

Proof : By (2.3.1)

$$\begin{aligned} \int_0^{M(Fx, Gy, qt)} \psi(s) ds &\geq \int_0^{\phi(\min \{M(Sx, Ty, t), M(Sx, Fx, t), M(Gy, Ty, t), M(Fx, Ty, t), M(Gy, Sx, t)\})} \psi(s) ds \\ &= \int_0^{\phi(\min \{M(Fx, Gy, t), M(Fx, Fx, t), M(Gy, Gy, t), M(Fx, Gy, t), M(Gy, Fx, t)\})} \psi(s) ds \\ &= \int_0^{\phi(\min \{M(Fx, Gy, t), 1, 1, M(Fx, Gy, t), M(Gy, Fx, t)\})} \psi(s) ds \\ &= \int_0^{\phi(M(Fx, Gy, t))} \psi(s) ds \\ \Rightarrow \int_0^{M(Fx, Gy, qt)} \psi(s) ds &\geq \int_0^{\phi(M(Fx, Gy, t))} \psi(s) ds \geq \int_0^{M(Fx, Gy, t)} \psi(s) ds \\ \Rightarrow \int_0^{M(Fx, Gy, t)} \psi(s) ds &\geq \int_0^{M(Fx, Gy, qt)} \psi(s) ds \geq \int_0^{M(Fx, Gy, t)} \psi(s) ds \\ \Rightarrow \int_0^{M(Fx, Gy, t)} \psi(s) ds &= \int_0^{M(Fx, Gy, qt)} \psi(s) ds \\ \Rightarrow M(Fx, Gy, qt) &= M(Fx, Gy, t) \\ \therefore Fx &= Gy \end{aligned}$$

Hence result follows from theorem (2.2)

Theorem 2.4: Let $(X, M, *)$ be a complete fuzzy metric space and let F, G, S and T be self-mapping of X . Let the pairs (F, S) and (G, T) be owc . If there exists $q \in (0, 1)$ such that

$$\int_0^{M(Fx, Gy, qt)} \psi(s) ds \geq \int_0^{\phi(M(Sx, Ty, t), M(Sx, Fx, t), M(Gy, Ty, t), M(Fx, Ty, t), M(Gy, Sx, t))} \psi(s) ds \quad (2.4.1)$$

for all $x, y \in X$ and $\phi: [0, 1]^5 \rightarrow [0, 1]$ such that $\phi(t, 1, 1, t, t) > t$ for all $0 < t < 1$. Then F, G, S and T have a unique common fixed point.

Proof : By (2.4.1)

$$\begin{aligned} \int_0^{M(Fx, Gy, qt)} \psi(s) ds &\geq \int_0^{\phi(M(Sx, Ty, t), M(Sx, Fx, t), M(Gy, Ty, t), M(Fx, Ty, t), M(Gy, Sx, t))} \psi(s) ds \\ &= \int_0^{\phi(M(Fx, Gy, t), 1, 1, M(Fx, Gy, t), M(Gy, Fx, t))} \psi(s) ds \\ &> \int_0^{M(Fx, Gy, t)} \psi(s) ds \\ \Rightarrow \int_0^{M(Fx, Gy, t)} \psi(s) ds &\geq \int_0^{M(Fx, Gy, qt)} \psi(s) ds \geq \int_0^{M(Fx, Gy, t)} \psi(s) ds \\ \Rightarrow \int_0^{M(Fx, Gy, t)} \psi(s) ds &= \int_0^{M(Fx, Gy, qt)} \psi(s) ds \\ \Rightarrow M(Fx, Gy, qt) &= M(Fx, Gy, t) \\ \therefore Fx &= Gy \end{aligned}$$

Hence result follows from theorem (2.2)

Theorem 2.5: Let $(X, M, *)$ be a complete fuzzy metric space and let F, G, S and T are self-mapping of X . Let the pairs (F, S) and (G, T) be owc . Suppose there exists $q \in (0, 1)$ such that

$$\int_0^{M(Fx, Gy, qt)} \psi(s) ds \geq \int_0^{\min \{M(Sx, Ty, t) * M(Fx, Sx, t), M(Gy, Ty, t) * M(Fx, Ty, t)\}} \psi(s) ds \quad (2.5.1)$$

for all $x, y \in X$. Then F, G, S and T have a unique common fixed point.

$$\begin{aligned} \text{Proof: } \int_0^{M(Fx, Gy, qt)} \psi(s) ds &\geq \int_0^{\min \{M(Sx, Ty, t) * M(Fx, Sx, t), M(Gy, Ty, t) * M(Fx, Ty, t)\}} \psi(s) ds \\ &= \int_0^{\min \{M(Fx, Gy, t) * 1, 1 * M(Fx, Gy, t)\}} \psi(s) ds \end{aligned}$$

$$= \int_0^{M(Fx,Gy,t)} \psi(s) ds$$

$$\Rightarrow \int_0^{M(Fx,Gy,qt)} \psi(s) ds \geq \int_0^{M(Fx,Gy,t)} \psi(s) ds$$

$$\Rightarrow M(Fx, Gy, qt) = M(Fx, Gy, t)$$

$$\therefore Fx = Gy$$

Hence result follows from theorem (2.2)

Theorem 2.6: Let $(X, M, *)$ be a complete fuzzy metric space where $*$ is the min t – norm and let F, G, S and T be self–mapping of X . Let the pairs (F, S) and (G, T) be owc . Suppose there exists $q \in (0,1)$ such that

$$\int_0^{M(Fx,Gy,qt)} \psi(s) ds \geq \int_0^{\min \{M(Sx,Ty,t)*M(Fx,Sx,t), M(Gy,Ty,t)*M(Gy,Sx,2t)*M(Fx,Ty,t)\}} \psi(s) ds \quad (2.6.1)$$

for all $x, y \in X$. Then F, G, S and T have a unique common fixed point.

Proof: By(2.6.1)

$$\begin{aligned} \int_0^{M(Fx,Gy,qt)} \psi(s) ds &\geq \int_0^{\min \{M(Sx,Ty,t)*M(Fx,Sx,t), M(Gy,Ty,t)*M(Gy,Sx,2t)*M(Fx,Ty,t)\}} \psi(s) ds \\ &= \int_0^{\min \{M(Fx,Gy,t)*M(Fx,Sx,t), M(Gy,Ty,t)*M(Sx,Ty,t)*M(Ty,Gy,t)*M(Fx,Ty,t)\}} \psi(s) ds \\ &= \int_0^{\min \{M(Fx,Gy,t)*1, 1*M(Fx,Gy,t)*1*M(Fx,Gy,t)\}} \psi(s) ds \\ &\geq \int_0^{M(Fx,Gy,t)} \psi(s) ds \end{aligned}$$

$$\Rightarrow \int_0^{M(Fx,Gy,qt)} \psi(s) ds \geq \int_0^{M(Fx,Gy,t)} \psi(s) ds$$

$$\Rightarrow \int_0^{M(Fx,Gy,t)} \psi(s) ds \geq \int_0^{M(Fx,Gy,qt)} \psi(s) ds \geq \int_0^{M(Fx,Gy,t)} \psi(s) ds$$

$$\Rightarrow \int_0^{M(Fx,Gy,t)} \psi(s) ds = \int_0^{M(Fx,Gy,qt)} \psi(s) ds$$

$$\Rightarrow M(Fx, Gy, qt) = M(Fx, Gy, t)$$

$$\therefore Fx = Gy$$

Hence result follows from theorem (2.2)

Theorem 2.7: Let $(X, M, *)$ be a complete fuzzy metric space and let F, G, S and T be self–mappings of X . Let the pairs (F, S) and (G, T) be owc . If there exists $q \in (0,1)$ such that

$$\int_0^{M(Sx,Sy,qt)} \psi(s) ds \geq \int_0^{\alpha M(Fx,Fy,t)+\beta \min \{M(Fx,Fy,t), M(Sx,Fx,t), M(Sy,Fy,t)\}} \psi(s) ds \quad (2.7.1)$$

for all $x, y \in X$ and $t > 0$ where $\alpha, \beta > 0, \alpha + \beta \geq 1$, then F, G, S and T have a unique common fixed point.

Proof : Let the pair (F, S) be owc.

$$\Rightarrow \exists x \in X \ni Fx = Sx.$$

$$\text{Suppose } \exists y \in X \ni Fy = Sy.$$

$$\begin{aligned} \int_0^{M(Sx,Sy,qt)} \psi(s) ds &\geq \int_0^{\alpha M(Fx,Fy,t)+\beta \min \{M(Fx,Fy,t), M(Sx,Fx,t), M(Sy,Fy,t)\}} \psi(s) ds \\ &= \int_0^{\alpha M(Sx,Sy,t)+\beta \min \{M(Sx,Sy,t), 1, M(Sx,Sy,t)\}} \psi(s) ds \\ &= \int_0^{\alpha M(Sx,Sy,t)+\beta M(Sx,Sy,t)} \psi(s) ds \end{aligned}$$

$$= \int_0^{(\alpha+\beta)M(Sx,Sy,t)} \psi(s) ds$$

$$\int_0^{M(Sx,Sy,qt)} \psi(s) ds \geq \int_0^{(\alpha+\beta)M(Sx,Sy,t)} \psi(s) ds > \int_0^{(\alpha+\beta)M(Sx,Sy,t)} \psi(s) ds$$

$$\Rightarrow \int_0^{M(Sx,Sy,qt)} \psi(s) ds \geq \int_0^{M(Sx,Sy,t)} \psi(s) ds$$

$$\Rightarrow M(Sx, Sy, qt) = M(Sx, Sy, t)$$

$$\therefore Sx = Sy$$

$$\therefore Fx = Fy$$

$$\Rightarrow F \text{ and } S \text{ have a unique common fixed point.}$$

ACKNOWLEDGEMENT

The third author (N.U) is grateful to Raghu Engineering College authorities for giving permission and the management of SITAM for giving facilities to carry on this research.

REFERENCES

- [1] Balasubramaniam, P., Muralisankar, S., Pant, R.P., Common 'fixed' points of four mappings in a fuzzy metric space, J. Fuzzy Math., 10(2) (2002), 379-384.
- [2] Cho. Y.J. , H.K. Pathak , S.M. Kang , J.S. Jung "Common Fixed Points of compatible maps of type (A) on fuzzy metric spaces" , Fuzzy Sets and Systems 93 (1998) , 99-111.
- [3] Grabiec .M. "Fixed points in fuzzy metric spaces", Fuzzy Sets and Systems 27 (1988) , 385-389.
- [4] George.A and Veeramani.P, On some results in fuzzy metric spaces, Fuzzy Sets and Systems, 64(1994), 395-399.
- [5] Jungck.G and Rhoades.B.E, Fixed points for set valued functions without continuity, Indian J. Pure Appl. Math., 29(1998), 227-238.
- [6] Kramosil.I and Michalek.J, Fuzzy metric and statistical metric spaces, Ky-bernetika, 11(1975), 336-344.
- [7]Malhotra.S.K,Navin Verma,Ravindra Sen, "Occasionally weakly compatible mappings and fixed point theorem in fuzzy metric spaces satisfying integral type inequality." International journal of statistika and matematika,vol2,issue3,2012,pp21-26.
- [8] Mishra.S.N, Sharma.N and S. L. Singh, Common fixed point of maps on fuzzy metric spaces, Internat. J. Math. Math. Sci., 17(1994), 253-258.
- [9] Pant. R.P., K. Jha, "A remark on common fixed points of four mappings in a fuzzy metric space", J Fuzzy Math. 12(2) (2004) ,433-437.
- [10] Schweizer.B and Sklar.A,Probabilistic Metric Spaces, North Holland, Ams-terdam, 1983.
- [11] Sessa.S, "On weak commutativity condition of mapping in fixed point consideration". Publ. Inst.Math(Beograd) N.S.,32(46),(1982),149-153.
- [12] Singh.B and Jain.S. Weak compatibility and fixed point theorems in fuzzy metric spaces,Ganita, 56(2)(2005), 167-176.
- [13] Subramanayam. P.V., "Common fixed points theorem in fuzzy metric spaces", InformationScience 83(1995) 105-112.
- [14] Vasuki.R. "Common fixed points for Rweakly commuting Maps in fuzzy metric spaces".Indian J .Pure Appl. Math 30 (1999) ,419-423.
- [15] Zadeh.L.A, Fuzzy sets, Infor. and Control, 8(1965), 338-353.