

International Journal of Innovative Research in Science, Engineering and Technology

(An ISO 3297: 2007 Certified Organization)

Vol. 4, Issue 12, December 2015

Hausdorff and Box Dimensions of Cantor Like Sets

Dr. S. M. Padhye¹ , Satish B. Khobragade²

Associate Professor, HOD, Department Of Mathematics, Shri R.L.T. College Of Science, Akola, Maharashtra,
 India¹

Research Student, Department Of Mathematics, Shri R.L.T. College Of Science, Akola, Maharashtra, India²

ABSTRACT: The Cantor like sets S have been constructed in [3], for any sequence $\{ \epsilon_n \}$ with $0 < \epsilon_n < 1$ with the help of sequence of sets $\{ S_n \}$ of subsets of $[0, 1]$ such that $S_n \supset S_{n+1}$, $S = \bigcap S_n$ and $m(S) = \prod_{n=1}^{\infty} (1 - \epsilon_n)$. Further $\sum \epsilon_n = \infty$ iff $m(S) = 0$.

In this paper we determine the estimates for the Hausdorff and upper / lower Box dimension of such Cantor like sets S. In particular it is proved that the Hausdorff dimension of the set S is given by $\frac{-\log 2}{\gamma}$, where $\gamma = \lim_{k \rightarrow \infty} \log \left[\prod_{n=1}^k \frac{(1-\epsilon_n)}{2} \right]^{1/k}$ and $\overline{\lim}_{k \rightarrow \infty} \frac{\log 2}{-\log \left[2 \prod_{n=1}^k \frac{(1-\epsilon_n)}{2} \right]^{1/k}} \geq \overline{\dim}_B(S) \geq \underline{\dim}_B(S) \geq \underline{\lim}_{k \rightarrow \infty} \frac{\log 2}{-\log \left\{ \left[\prod_{n=1}^k \frac{(1-\epsilon_n)}{2} \right]^{1/k} [\epsilon_{k+1}]^{1/k} \right\}}$. If the above upper and lower limits are equal then $\underline{\dim}_B(S) = \overline{\dim}_B(S)$.

KEYWORDS : Cantor set, Cantor like set, Hausdorff dimension, upper Box dimension and lower Box dimension.

I. INTRODUCTION

Definition 1.1 Hausdorff measure

If U is any non-empty subset of n-dimensional Euclidean space, \mathbb{R}^n , then diameter of U is defined as $|U| = \sup\{|x - y| : x, y \in U\}$. If $\{U_i\}$ is a countable (or finite) collection of sets of diameter at most δ that cover F, i.e. $F \subset \bigcup_{i=1}^{\infty} U_i$ with $0 < |U_i| \leq \delta$ for each i, we say that $\{U_i\}$ is a δ - cover of F.

Suppose that F is a subset of \mathbb{R}^n and s is a non-negative number. For any $\delta > 0$ we define

$$H_s^{\delta}(F) = \inf \{ \sum_{i=1}^{\infty} |U_i|^s : \{U_i\} \text{ is a } \delta\text{- cover of } F \}$$

We define $H^s(F) = \lim_{\delta \rightarrow 0} H_s^{\delta}(F)$

This limit exists for any subset F of \mathbb{R}^n , though the limiting value can be 0 or ∞ .

We call $H^s(F)$ the s-dimensional Hausdorff measure of F.

OR $H^s(F) = \inf \{ \sum_{i=1}^{\infty} |U_i|^s : \{U_i\} \text{ is a countable cover of } F \}$

Definition 1.2 Hausdorff dimension

The Hausdorff dimension of F is defined as

$$\dim_H(F) = \inf \{ s : H^s(F) = 0 \}.$$

It is proved that $\inf \{ s : H^s(F) = 0 \} = \sup \{ s : H^s(F) = \infty \}$.

Above definitions and properties are taken from [2].

Lemma 1.3 :- $(a + b)^{\alpha} \leq a^{\alpha} + b^{\alpha}$, $a, b \geq 0$ and $0 < \alpha < 1$.

International Journal of Innovative Research in Science, Engineering and Technology

(An ISO 3297: 2007 Certified Organization)

Vol. 4, Issue 12, December 2015

Proof :-

$$\text{Let } f(c) = 1 + c^\alpha - (1 + c)^\alpha, c \geq 0.$$

We show that f is monotonic increasing. For this we show that its derivative is positive.

$$\begin{aligned} f'(c) &= \alpha c^{\alpha-1} - \alpha(1+c)^{\alpha-1} \\ &= \alpha[c^{\alpha-1} - (1+c)^{\alpha-1}] \\ &= \alpha \left[\frac{1}{c^{1-\alpha}} - \frac{1}{(c+1)^{1-\alpha}} \right] \end{aligned}$$

Since $c < c+1$, $c^{1-\alpha} < (c+1)^{1-\alpha}$. Thus $f'(c) > 0$, $c \geq 0$

This shows that f is monotonically increasing for $c \geq 0$. But $f(0) = 0$ and hence $f(c) \geq 0$ for all $c \geq 0$.

Thus for $c = \frac{a}{b}$, $b > 0$ we get $1 + (\frac{a}{b})^\alpha \geq (1 + \frac{a}{b})^\alpha$ i.e. $(a+b)^\alpha \leq a^\alpha + b^\alpha$.

This proves the result.

Lemma 1.4 : Let S be compact subset of \mathbb{R} and $S \subset \bigcup_{n=1}^{\infty} C_n$ where C'_n s are intervals. Then for given $\epsilon > 0$ and $0 < \alpha < 1$, there exists finitely many open intervals $D_1, D_2, D_3, \dots, D_m$

such that $\bigcup_{n=1}^{\infty} C_n \subseteq \bigcup_{j=1}^m D_j$ and $\sum_{j=1}^m |D_j|^\alpha \leq \sum_{n=1}^{\infty} |C_n|^\alpha + \epsilon$

Proof : Suppose $\epsilon > 0$ is given and $0 < \alpha < 1$. For $S \subset \bigcup_{n=1}^{\infty} C_n$ where C'_n s are interval ,define F_n to be a closed interval with end points same as that of C_n . Then $S \subset \bigcup C_n \subset \bigcup F_n$ where each F_n is a closed interval.

Let $F_n = [a_n, b_n]$. For $\epsilon' > 0$ and $\epsilon' = \epsilon \cdot 2^{1-\alpha}$, consider $E_n = (a_n - (\frac{\epsilon'}{2^{n+1}})^{1/\alpha}, b_n + (\frac{\epsilon'}{2^{n+1}})^{1/\alpha})$

Then $S \subset \bigcup_{n=1}^{\infty} F_n \subset \bigcup_{n=1}^{\infty} E_n$.

Since S is compact. There is a finite subfamily $\{E_{n_1}, E_{n_2}, \dots, E_{n_m}\}$ of $\{E_n\}$ such that $S \subset \bigcup_{j=1}^m E_{n_j}$. Put $D_j = E_{n_j}$, $j = 1, 2, 3, \dots, m$

$$\begin{aligned} \sum_{j=1}^m |D_j|^\alpha &\leq \sum_{n=1}^{\infty} |E_n|^\alpha = \sum_{n=1}^{\infty} \{ (b_n - a_n) + 2(\frac{\epsilon'}{2^{n+1}})^{1/\alpha} \}^\alpha \\ \sum_{j=1}^m |D_j|^\alpha &\leq \sum_{n=1}^{\infty} [(b_n - a_n)^\alpha + 2^\alpha \frac{\epsilon'}{2^{n+1}}] \quad (\because \text{By Lemma 1.3}) \\ &= \sum_{n=1}^{\infty} (b_n - a_n)^\alpha + 2^\alpha \sum_{n=1}^{\infty} \frac{\epsilon'}{2^{n+1}} \\ &= \sum_{n=1}^{\infty} |C_n|^\alpha + 2^\alpha \frac{\epsilon'}{2} \\ &= \sum_{n=1}^{\infty} |C_n|^\alpha + \epsilon \quad (\because \epsilon = 2^\alpha \frac{\epsilon'}{2}) \\ \therefore \sum_{j=1}^m |D_j|^\alpha &\leq \sum_{n=1}^{\infty} |C_n|^\alpha + \epsilon \end{aligned}$$

II. MAIN RESULTS

2.1 Construction of Cantor like sets

In the construction of Cantor like set S at first stage from $I = [0, 1]$ we remove middle open interval $I_{1,1} = (\frac{(1-\epsilon_1)}{2}, \frac{(1+\epsilon_1)}{2})$ of length ϵ_1 . We get two closed intervals denoted by $J_{1,1} = [0, \frac{(1-\epsilon_1)}{2}]$ and $J_{1,2} = [\frac{(1+\epsilon_1)}{2}, 1]$. We get the set $S_1 = J_{1,1} \cup J_{1,2}$ with measure $(1 - \epsilon_1)$ i.e. $m(S_1) = (1 - \epsilon_1)$. In second stage from $J_{1,1}$ and $J_{1,2}$ we remove two

International Journal of Innovative Research in Science, Engineering and Technology

(An ISO 3297: 2007 Certified Organization)

Vol. 4, Issue 12, December 2015

middle open intervals $I_{1,2} = \left(\frac{(1-\epsilon_1)(1-\epsilon_2)}{4}, \frac{(1-\epsilon_1)(1+\epsilon_2)}{4}\right)$ and $I_{2,2} = \left(\frac{4-(1-\epsilon_1)(1+\epsilon_2)}{4}, \frac{4-(1-\epsilon_1)(1-\epsilon_2)}{4}\right)$. The remaining four closed intervals are denoted by

$$J_{2,1} = [0, \frac{(1-\epsilon_1)(1-\epsilon_2)}{4}], J_{2,2} = [\frac{(1-\epsilon_1)(1+\epsilon_2)}{4}, \frac{(1-\epsilon_1)}{2}], J_{2,3} = [\frac{(1+\epsilon_1)}{2}, \frac{4-(1-\epsilon_1)(1+\epsilon_2)}{4}], \\ J_{2,4} = [\frac{4-(1-\epsilon_1)(1-\epsilon_2)}{4}, 1].$$

\therefore Length of removed intervals $= m(I_{1,2}) + m(I_{2,2}) = (1 - \epsilon_1)\epsilon_2 < (1 - \epsilon_1)$

We get the set S_2 union of remaining four closed intervals i.e. $S_2 = J_{2,1} \cup J_{2,2} \cup J_{2,3} \cup J_{2,4}$ with measure $(1 - \epsilon_1)(1 - \epsilon_2)$ i.e. $m(S_2) = (1 - \epsilon_1)(1 - \epsilon_2)$

Continuing in this way we can construct Cantor like set $S = \bigcap_{n=1}^{\infty} S_n$ with $m(S) = \lim_{n \rightarrow \infty} m(S_n) = \prod_{n=1}^{\infty} (1 - \epsilon_n)$. This is from [3].

Theorem 2.2 : Hausdorff Dimension of Cantor like set S defined above is $\frac{-\log 2}{\lim_{k \rightarrow \infty} \log[\prod_{n=1}^k \frac{(1-\epsilon_n)}{2}]^{1/k}}$ i.e. $\frac{-\log 2}{\gamma}$ if the required limit exists and $\inf\{2^j \left[\prod_{n=1}^j \frac{(1-\epsilon_n)}{2} \right]^\alpha : j \geq 1\} > 0$.

Proof : Let $\alpha = \frac{-\log 2}{\gamma}$, where $\gamma = \lim_{k \rightarrow \infty} \log[\prod_{n=1}^k \frac{(1-\epsilon_n)}{2}]^{1/k}$

We first show that if $\beta > \alpha$ then $H^\beta(S) = 0$.

Take $k \geq 1$ and let $J_{1,k}, J_{2,k}, J_{3,k}, \dots, J_{2^k, k}$ be the family of 2^k intervals remaining at k^{th} step whose union is S_k .

$$|J_{j,k}| = \text{Length of each } J_{j,k} = \prod_{n=1}^k \frac{(1-\epsilon_n)}{2}. \therefore |J_{j,k}|^\beta = \left[\prod_{n=1}^k \frac{(1-\epsilon_n)}{2} \right]^\beta \\ \sum_{j=1}^{2^k} |J_{j,k}|^\beta = \sum_{j=1}^{2^k} \left[\prod_{n=1}^k \frac{(1-\epsilon_n)}{2} \right]^\beta = 2^k \left[\prod_{n=1}^k \frac{(1-\epsilon_n)}{2} \right]^\beta \dots \dots \dots (1) \\ \text{Now } \alpha = \frac{-\log 2}{\gamma}, \text{ where } \gamma = \lim_{k \rightarrow \infty} \log[\prod_{n=1}^k \frac{(1-\epsilon_n)}{2}]^{1/k} \\ \gamma \leq \lim_{k \rightarrow \infty} \log\left[\left(\frac{1}{2}\right)^k\right]^{1/k} \quad (\because (1 - \epsilon_n) < 1 \forall n = 1, 2, 3, \dots, k) \\ \gamma \leq \lim_{k \rightarrow \infty} \log\left(\frac{1}{2}\right) = \log\left(\frac{1}{2}\right) = -\log 2 \\ \therefore \gamma \leq -\log 2. \text{ Thus } \gamma < 0$$

Now, since $\beta > \alpha = \frac{-\log 2}{\gamma}$

$$\therefore \beta\gamma < -\log 2 \quad (\because \gamma < 0) \\ \Rightarrow \log 2 + \beta\gamma < 0$$

$$\Rightarrow \log 2 + \beta \lim_{k \rightarrow \infty} \log[\prod_{n=1}^k \frac{(1-\epsilon_n)}{2}]^{1/k} < 0 \dots \dots \dots (2)$$

Now,

$$2^k \left[\prod_{n=1}^k \frac{(1-\epsilon_n)}{2} \right]^\beta = e^{\log \{2^k [\prod_{n=1}^k \frac{(1-\epsilon_n)}{2}]^\beta\}} \\ = e^{\{\log 2^k + \log [\prod_{n=1}^k \frac{(1-\epsilon_n)}{2}]^\beta\}} \\ = e^{\{k \log 2 + \beta \log [\prod_{n=1}^k \frac{(1-\epsilon_n)}{2}]\}} \\ = e^{k \{\log 2 + \frac{\beta}{k} \log [\prod_{n=1}^k \frac{(1-\epsilon_n)}{2}]\}} \\ = e^{k \{\log 2 + \beta \log [\prod_{n=1}^k \frac{(1-\epsilon_n)}{2}]^{1/k}\}}$$

Putting in equation (1)

International Journal of Innovative Research in Science, Engineering and Technology

(An ISO 3297: 2007 Certified Organization)

Vol. 4, Issue 12, December 2015

$$\sum_{j=1}^{2^k} |J_{j,k}|^\beta = e^{k \{\log 2 + \beta \log \left[\prod_{n=1}^k \frac{(1-\epsilon_n)}{2} \right] \}^{1/k}} \quad \dots \dots \dots \quad (3)$$

We claim that $\lim_{k \rightarrow \infty} e^{k \{ \log 2 + \beta \log [\prod_{n=1}^k \frac{(1-\epsilon_n)}{2}]^{1/k} \}} = 0$

Since $\log 2 + \beta\gamma$ is negative by equation (2), we can put $\log 2 + \beta\gamma = -\delta_1$, $\delta_1 > 0$.

$$\begin{aligned}
 & \text{If } \gamma_k = \log \left[\prod_{n=1}^k \frac{(1-\epsilon_n)}{2} \right]^{1/k}, \text{ then } \lim \gamma_k = \gamma \text{ and hence } \exists K_1 \text{ such that } k \geq K_1 \\
 & \Rightarrow |(\log 2 + \beta \gamma_k) + \delta_1| < \frac{\delta_1}{2} \quad \Rightarrow |(\log 2 + \beta \gamma_k) - (-\delta_1)| < \frac{\delta_1}{2} \\
 & \Rightarrow -\delta_1 - \frac{\delta_1}{2} < (\log 2 + \beta \gamma_k) < -\delta_1 + \frac{\delta_1}{2} \\
 & \quad - \frac{3\delta_1}{2} < (\log 2 + \beta \gamma_k) < -\frac{\delta_1}{2} \\
 & \Rightarrow k(\log 2 + \beta \gamma_k) < -\frac{k\delta_1}{2} \therefore e^{k(\log 2 + \beta \gamma_k)} < e^{-\frac{k\delta_1}{2}}, k \geq 1.
 \end{aligned}$$

This proves that $\lim_{k \rightarrow \infty} e^{k\{\log 2 + \beta \log [\prod_{n=1}^k \frac{(1-\epsilon_n)}{2}]^{1/k}\}} = 0$ (4)

Now, for $\delta > 0$, choose $k \geq 1$ such that $|J_{j,k}| = \prod_{n=1}^k \frac{(1-\epsilon_n)}{2} < \delta$, $j = 1, 2, \dots, 2^k$.

This is possible if $\lim_{k \rightarrow \infty} \prod_{n=1}^k (1 - \epsilon_n)$ exists and then $\prod_{n=1}^{\infty} \frac{(1 - \epsilon_n)}{2} = \lim_{k \rightarrow \infty} \frac{\prod_{n=1}^k (1 - \epsilon_n)}{2^k} = \lim_{k \rightarrow \infty} \prod_{n=1}^k (1 - \epsilon_n) \lim_{k \rightarrow \infty} \frac{1}{2^k} = 0$. Then $\{J_{j,k} : j = 1, 2, \dots, 2^k\}$ is a δ -cover of S_k and therefore of S .

Now $H^\beta_\delta(S) = \inf \{ \sum_{i=1}^{\infty} |U_i|^\beta : \{U_i\} \text{ is a } \delta\text{-cover of } S \}$. Since $S = \bigcap_{k=1}^{\infty} S_k$, $S \subset S_k$ and $S_k = \bigcup_{j=1}^{2^k} J_{j,k}$,

$$\therefore H_{-\delta}(S) \leq \sum_{i=1}^{2^k} |J_{i,k}|^\beta = e^{k \{ \log 2 + \beta \log \left[\prod_{n=1}^k \frac{(1-\epsilon_n)}{2} \right]^{1/k} \}}$$

As $\delta \rightarrow 0$, $k \rightarrow \infty$

Now $H^\beta(S) = \lim_{\delta \rightarrow 0} H^\beta_\delta(S) \leq \lim_{k \rightarrow \infty} e^{k \{\log 2 + \beta \log [\prod_{n=1}^k \frac{(1-\epsilon_n)}{2}]^{1/k}\}} = 0$ (By equation (4))

$$\therefore H^\beta(S) = 0$$

Thus $(\alpha, \infty) \subset \{ \beta > 0 : H^\beta(S) = \{ \beta > 0 \}$

$$\therefore \inf\{\beta > 0 : H^\beta$$

$$\therefore \dim_H(S) \leq \alpha$$

Now to show that $\dim_H(S) \geq \alpha$

suppose further that $\eta = \inf$

We prove that $H^\alpha(S) > 0$.

This will imply that $\dim_H(S) \geq \alpha$ as

$$\dim_H(S) = \sup\{\gamma \geq 0 : H^\gamma(S)$$

Now we claim that $H^\alpha(S) \geq \eta$

Suppose $\{C_i\}$ is any countable cover of S by nonempty intervals.

We show that $\sum |C_i|^\alpha \geq \eta$.

$\epsilon > 0$ we choose $\{D_i\}$ a finite collection of open or closed intervals covering S such that

$$\sum_{i=1}^m |D_i|^\alpha < \sum_{i=1}^{\infty} |C_i|^\alpha + \epsilon \quad (\text{By Lemma I.4})$$

By taking intersection of D_i with $[0,1]$ we may

International Journal of Innovative Research in Science, Engineering and Technology

(An ISO 3297: 2007 Certified Organization)

Vol. 4, Issue 12, December 2015

Then D_i can intersect at most one basic interval of S_{k_i} . If D_i intersects two intervals each of length $\prod_{n=1}^{k_i} \frac{(1-\epsilon_n)}{2}$ of S_{k_i} then $|D_i| \geq \left[\prod_{n=1}^{k_i-1} \frac{(1-\epsilon_n)}{2} \right] \epsilon_{k_i}$ which is distance between consecutive intervals of S_{k_i} which is false.

If $j \geq k_i$ then the basic intervals of S_j are $[0, \prod_{n=1}^j \frac{(1-\epsilon_n)}{2}] \cup \dots \cup [\frac{2^j - \prod_{n=1}^j (1-\epsilon_n)}{2^j}, 1]$

These are 2^j basic intervals.

Choose $j \geq k_1, k_2, \dots, k_m$ so that $|D_i| \geq \left[\prod_{n=1}^j \frac{(1-\epsilon_n)}{2} \right] \epsilon_{j+1}$ for all $i = 1, 2, 3, \dots, m$.

For $i = 1, 2, 3, \dots, m$, if N_{k_i} is the number of sets D_i which intersect the basic intervals of S_j . Then $\{D_i\}$ intersects all 2^j basic intervals of length $\prod_{n=1}^j \frac{(1-\epsilon_n)}{2}$ in S_j and $\sum_{i=1}^m N_{k_i} = 2^j$.

$$\begin{aligned}
\text{From equation (8)} \quad & |D_i| \geq \left[\prod_{n=1}^j \frac{(1-\epsilon_n)}{2} \right] \epsilon_{j+1} \\
\therefore \quad & |D_i|^\alpha \geq \left\{ \left[\prod_{n=1}^j \frac{(1-\epsilon_n)}{2} \right] \epsilon_{j+1} \right\}^\alpha \\
& |D_i|^\alpha \geq \left[\prod_{n=1}^j \frac{(1-\epsilon_n)}{2} \right]^\alpha [\epsilon_{j+1}]^\alpha \\
& \sum_{N_{k_i}} |D_i|^\alpha \geq \left[\prod_{n=1}^j \frac{(1-\epsilon_n)}{2} \right]^\alpha [\epsilon_{j+1}]^\alpha N_{k_i} \\
\therefore \sum_{i=1}^m |D_i|^\alpha \geq & \left[\prod_{n=1}^j \frac{(1-\epsilon_n)}{2} \right]^\alpha [\epsilon_{j+1}]^\alpha \sum_{i=1}^m N_{k_i} \\
\sum_{i=1}^m |D_i|^\alpha \geq & \left[\prod_{n=1}^j \frac{(1-\epsilon_n)}{2} \right]^\alpha [\epsilon_{j+1}]^\alpha 2^j \quad (\because \sum_{i=1}^m N_{k_i} = 2^j)
\end{aligned}$$

$$\text{But } \sum_{i=1}^{\infty} |C_i|^\alpha + \epsilon > \sum_{i=1}^m |D_i|^\alpha$$

$$\sum_{i=1}^{\infty} |C_i|^\alpha + \epsilon > 2^j \left[\prod_{n=1}^j \frac{(1-\epsilon_n)}{2} \right]^\alpha [\epsilon_{j+1}]^\alpha \geq \eta \quad (\text{From equation(6)})$$

$\therefore \sum_{i=1}^{\infty} |C_i|^\alpha + \epsilon \geq \eta$. Since $\epsilon > 0$ is arbitrary, $\sum_{i=1}^{\infty} |C_i|^\alpha + \epsilon \geq \eta$
 i.e. $\inf\{\sum_{i=1}^{\infty} |C_i|^\alpha + \epsilon : \{C_i\} \text{ is countable cover of } S\} \geq \eta$

By property of Hausdorff Dimension

By property of Hausdorff Dimension

Since $H^\alpha(S) \geq 0$

(By equation (9))

$\therefore \alpha \in \{ \gamma \geq 0 : H^\gamma(S) \geq 0 \}$

$$\therefore \sup\{\gamma \geq 0 : H^\gamma(S) \geq 0\} \geq \alpha$$

$$\therefore \dim_{\mathbb{C}}(S) \geq \alpha$$

(10)

From equations (5) and (10), we get

From equation

$$\dim_{\mathbb{K}}(S) = \alpha$$

We consider now several particular cases of $\{\epsilon_n\}$ where the condition in equation (6) is satisfied so that the result in Theorem 2.2 holds.

Case I: Cantor ternary set. It is obtained by taking $\epsilon_n = \frac{1}{3^n}$ for all $n \geq 1$.

$$\text{Thus } \prod_{n=1}^k \frac{(1-\epsilon_n)}{3} = \prod_{n=1}^k \left(1 - \frac{1}{3}\right) = \left(\frac{1}{3}\right)^k$$

$$\therefore \left[\prod_{n=1}^k \frac{(1-\epsilon_n)}{2} \right]^{1/k} = \frac{1}{2}$$

International Journal of Innovative Research in Science, Engineering and Technology

(An ISO 3297: 2007 Certified Organization)

Vol. 4, Issue 12, December 2015

$$\begin{aligned} \therefore \log \left[\prod_{n=1}^k \frac{(1-\epsilon_n)}{2} \right]^{1/k} &= \log \left(\frac{1}{3} \right) \\ \alpha &= \frac{\lim_{k \rightarrow \infty} \log \left[\prod_{n=1}^k \frac{(1-\epsilon_n)}{2} \right]^{1/k}}{-\log 2} \\ &= \frac{-\log 2}{\log \left(\frac{1}{3} \right)} \\ \alpha &= \frac{\log 2}{\log 3} \\ \therefore 3^\alpha &= 2 \end{aligned}$$

$$\begin{aligned} \text{Now } 2^j \left[\prod_{n=1}^j \frac{(1-\epsilon_n)}{2} \right]^\alpha [\epsilon_{j+1}]^\alpha &= 2^j \left(\frac{1}{3} \right)^{j\alpha} \left(\frac{1}{3} \right)^\alpha \\ &= 2^j 2^{-j} \frac{1}{2} \\ &= \frac{1}{2} \end{aligned} \quad (\because 3^\alpha = 2)$$

$$\therefore \inf \left\{ 2^j \left[\prod_{n=1}^j \frac{(1-\epsilon_n)}{2} \right]^\alpha [\epsilon_{j+1}]^\alpha : j \geq 1 \right\} = \frac{1}{2}$$

Case II : Consider the case $\epsilon_n = \epsilon$ for all $n \geq 1$, $\epsilon < 1$.

$$\begin{aligned} \text{Thus } \prod_{n=1}^k \frac{(1-\epsilon_n)}{2} &= \prod_{n=1}^k \frac{(1-\epsilon)}{2} = \left(\frac{1-\epsilon}{2} \right)^k \\ \therefore \left[\prod_{n=1}^k \frac{(1-\epsilon_n)}{2} \right]^{1/k} &= \frac{1-\epsilon}{2} \\ \therefore \log \left[\prod_{n=1}^k \frac{(1-\epsilon_n)}{2} \right]^{1/k} &= \log \left(\frac{1-\epsilon}{2} \right) \end{aligned}$$

$$\therefore \lim_{k \rightarrow \infty} \log \left[\prod_{n=1}^k \frac{(1-\epsilon_n)}{2} \right]^{1/k} = \log \left(\frac{1-\epsilon}{2} \right)$$

$$\begin{aligned} \text{Also } \alpha &= \frac{\lim_{k \rightarrow \infty} \log \left[\prod_{n=1}^k \frac{(1-\epsilon_n)}{2} \right]^{1/k}}{-\log 2} \\ &= \frac{-\log 2}{\log \left(\frac{1-\epsilon}{2} \right)} \end{aligned}$$

$$\alpha \log \left(\frac{1-\epsilon}{2} \right) = -\log 2$$

$$\therefore \log \left(\frac{1-\epsilon}{2} \right)^\alpha = \log 2^{-1}$$

$$\therefore \left(\frac{1-\epsilon}{2} \right)^\alpha = 2^{-1}$$

$$\begin{aligned} \text{Now } 2^j \left[\prod_{n=1}^j \frac{(1-\epsilon_n)}{2} \right]^\alpha [\epsilon_{j+1}]^\alpha &= 2^j \left(\frac{1-\epsilon}{2} \right)^{j\alpha} \epsilon^\alpha \\ &= 2^j 2^{-j} \epsilon^{-1} \\ &= \epsilon^\alpha \end{aligned} \quad (\because \left(\frac{1-\epsilon}{2} \right)^\alpha = 2^{-1})$$

$$\therefore \inf \left\{ 2^j \left[\prod_{n=1}^j \frac{(1-\epsilon_n)}{2} \right]^\alpha [\epsilon_{j+1}]^\alpha : j \geq 1 \right\} = \epsilon^\alpha$$

Case III : Consider the case $\epsilon_n = \frac{1}{n+1}$ for all $n \geq 1$

$$\begin{aligned} \text{Thus } \prod_{n=1}^k \frac{(1-\epsilon_n)}{2} &= \prod_{n=1}^k \frac{\left[1 - \frac{1}{(n+1)} \right]}{2} \\ &= \prod_{n=1}^k \frac{\frac{n}{n+1}}{2(n+1)} \\ &= \prod_{n=1}^k \frac{n}{2(n+1)} \\ &= \frac{1}{2^k} \prod_{n=1}^k \frac{n}{n+1} \end{aligned}$$

International Journal of Innovative Research in Science, Engineering and Technology

(An ISO 3297: 2007 Certified Organization)

Vol. 4, Issue 12, December 2015

$$\begin{aligned}
 &= \frac{1}{2^k} \frac{1}{2} \frac{2}{3} \frac{3}{4} \frac{4}{5} \cdots \frac{k}{(k+1)} \\
 \prod_{n=1}^k \frac{(1-\epsilon_n)}{2} &= \frac{1}{2^k} \frac{1}{k+1} \\
 \therefore [\prod_{n=1}^k \frac{(1-\epsilon_n)}{2}]^{1/k} &= \left[\frac{1}{2^k(k+1)} \right]^{1/k} = \frac{1}{2(k+1)^{1/k}} = \frac{1}{2} \left(\frac{1}{k+1} \right)^{1/k} \\
 \therefore \log \left[\prod_{n=1}^k \frac{(1-\epsilon_n)}{2} \right]^{1/k} &= \log \left[\frac{1}{2} \left(\frac{1}{k+1} \right)^{1/k} \right] \\
 \alpha &= \frac{\lim_{k \rightarrow \infty} \log [\prod_{n=1}^k \frac{(1-\epsilon_n)}{2}]^{1/k}}{-\log 2} \\
 &= \frac{\lim_{k \rightarrow \infty} \log \left[\frac{1}{2} \left(\frac{1}{k+1} \right)^{1/k} \right]}{-\log 2} \\
 &= \frac{-\log 2}{\log \left(\frac{1}{2} \right)} \\
 \alpha &= \frac{-\log 2}{-\log 2} = 1 \\
 \therefore \alpha &= 1
 \end{aligned}$$

Now $2^j \left[\prod_{n=1}^j \frac{(1-\epsilon_n)}{2} \right]^\alpha [\epsilon_{j+1}]^\alpha = 2^j \left[\frac{1}{2^{j(j+1)}} \right] \frac{1}{(j+2)} = \frac{1}{(j+1)(j+2)}$
 $\therefore \inf \{ 2^j \left[\prod_{n=1}^j \frac{(1-\epsilon_n)}{2} \right]^\alpha [\epsilon_{j+1}]^\alpha : j \geq 1 \} = 0$

$\alpha = 1$ but $\inf \{ 2^j \left[\prod_{n=1}^j \frac{(1-\epsilon_n)}{2} \right]^\alpha [\epsilon_{j+1}]^\alpha : j \geq 1 \} = 0$ i.e. condition is not satisfied.

Case IV : Consider the case $\epsilon_n = \frac{n}{2(n+1)}$ for all $n \geq 1$.

$$\begin{aligned}
 \text{Thus } \prod_{n=1}^k \frac{(1-\epsilon_n)}{2} &= \prod_{n=1}^k \frac{1 - \frac{n}{2(n+1)}}{2} \\
 &= \prod_{n=1}^k \frac{2(n+1)-n}{4(n+1)} \\
 &= \prod_{n=1}^k \frac{n+2}{4(n+1)} \\
 &= \frac{1}{4^k} \prod_{n=1}^k \frac{n+2}{n+1} \\
 &= \frac{1}{4^k} \frac{3}{2} \frac{4}{3} \frac{5}{4} \frac{6}{5} \cdots \frac{(k+2)}{(k+1)} \\
 \prod_{n=1}^k \frac{(1-\epsilon_n)}{2} &= \frac{1}{4^k} \frac{(k+2)}{2} \\
 \therefore [\prod_{n=1}^k \frac{(1-\epsilon_n)}{2}]^{1/k} &= \left[\frac{(k+2)}{2 \cdot 4^k} \right]^{1/k} = \frac{(k+2)^{1/k}}{4 \cdot 2^{1/k}} = \frac{1}{4} \left(1 + \frac{k}{2} \right)^{1/k} \\
 \therefore \log \left[\prod_{n=1}^k \frac{(1-\epsilon_n)}{2} \right]^{1/k} &= \log \left[\frac{1}{4} \left(1 + \frac{k}{2} \right)^{1/k} \right] \\
 \alpha &= \frac{\lim_{k \rightarrow \infty} \log [\prod_{n=1}^k \frac{(1-\epsilon_n)}{2}]^{1/k}}{-\log 2} \\
 &= \frac{\lim_{k \rightarrow \infty} \log \left[\frac{1}{4} \left(1 + \frac{k}{2} \right)^{1/k} \right]}{-\log 2} \\
 &= \frac{-\log 2}{\log \left(\frac{1}{4} \right)} \\
 \alpha &= \frac{\log 2}{\log 4} = \frac{1}{2}
 \end{aligned}$$

International Journal of Innovative Research in Science, Engineering and Technology

(An ISO 3297: 2007 Certified Organization)

Vol. 4, Issue 12, December 2015

$$\begin{aligned}
 \therefore \alpha &= \frac{1}{2} \\
 \text{Now } 2^j \left[\prod_{n=1}^j \frac{(1-\epsilon_n)}{2} \right]^\alpha [\epsilon_{j+1}]^\alpha &= 2^j \left[\frac{1}{4^j} \left(\frac{j+2}{2} \right) \right]^{1/2} \left[\frac{j+1}{2(j+2)} \right]^{1/2} \\
 &= \frac{2^j}{2^j} \left(\frac{j+2}{2} \right)^{1/2} \left[\frac{j+1}{2(j+2)} \right]^{1/2} \\
 &= \left(\frac{j+1}{4} \right)^{1/2} \\
 &= \frac{1}{2}(j+1)^{1/2} \\
 \therefore \inf \left\{ 2^j \left[\prod_{n=1}^j \frac{(1-\epsilon_n)}{2} \right]^\alpha [\epsilon_{j+1}]^\alpha : j \geq 1 \right\} &= \frac{\sqrt{2}}{2} = \frac{1}{\sqrt{2}}
 \end{aligned}$$

Definition 2.3 Box counting dimension

Let F be any nonempty bounded subset of \mathbb{R}^n and let $N_\delta(F)$ be the smallest number of sets of diameter at most δ which can cover F . The lower and upper box dimensions of F are defined respectively as $\underline{\dim}_B(F) = \liminf_{\delta \rightarrow 0} \frac{\log N_\delta(F)}{-\log \delta}$ and $\overline{\dim}_B(F) = \limsup_{\delta \rightarrow 0} \frac{\log N_\delta(F)}{-\log \delta}$.

If the lower and upper box-counting dimensions of F are equal then the common value is called box-counting dimension of F or box- dimension of F and $\dim_B(F) = \lim_{\delta \rightarrow 0} \frac{\log N_\delta(F)}{-\log \delta}$. Above definitions are taken from [2].

Theorem 2.4: Lower and upper box-dimensions of Cantor like set S associated with $\{\epsilon_n\}$ of positive numbers less than 1 with $\sum \epsilon_j = \infty$ satisfy $\underline{\dim}_{k \rightarrow \infty} \frac{\log 2}{-\log \{ [\prod_{n=1}^k \frac{(1-\epsilon_n)}{2}]^{1/k} [\epsilon_{k+1}]^{1/k} \}}$ and $\overline{\dim}_B(S) \leq \overline{\lim}_{k \rightarrow \infty} \frac{\log 2}{-\log [2 \prod_{n=1}^k \frac{(1-\epsilon_n)}{2}]^{1/k}}$.

Proof : We know that in the construction of Cantor like set S , at k^{th} stage the set S_k consists of union of 2^k disjoint intervals each of length $\prod_{n=1}^k \frac{(1-\epsilon_n)}{2}$. For $0 < \delta < 1$, let U be an interval with $|U| < \delta$. Choose $k \geq 1$ such that $2 \prod_{n=1}^k \frac{(1-\epsilon_n)}{2} \geq \delta > 2 \prod_{n=1}^{k+1} \frac{(1-\epsilon_n)}{2}$. This is possible since $\lim_{k \rightarrow \infty} \prod_{n=1}^k (1 - \epsilon_n)$ exists. Then U can cover at most one interval among 2^k intervals of length $\prod_{n=1}^k \frac{(1-\epsilon_n)}{2}$.

For if U contains more than one interval from $J_{j,k}$ then $\delta > |U| \geq 2 \prod_{n=1}^k \frac{(1-\epsilon_n)}{2}$.

$\therefore \delta > 2 \prod_{n=1}^k \frac{(1-\epsilon_n)}{2}$ which is not possible because $\delta \leq 2 \prod_{n=1}^k \frac{(1-\epsilon_n)}{2}$.

\therefore At most 2^k intervals of diameter less than δ are required to cover S .

$\therefore N_\delta(S) \leq 2^k$

Now $\delta \leq 2 \prod_{n=1}^k \frac{(1-\epsilon_n)}{2}$

$\log \delta \leq \log [2 \prod_{n=1}^k \frac{(1-\epsilon_n)}{2}]$

$-\log \delta \geq -\log [2 \prod_{n=1}^k \frac{(1-\epsilon_n)}{2}]$

$\therefore \frac{1}{-\log \delta} \leq \frac{1}{-\log [2 \prod_{n=1}^k \frac{(1-\epsilon_n)}{2}]} \quad (\because -\log \delta > 0)$

International Journal of Innovative Research in Science, Engineering and Technology

(An ISO 3297: 2007 Certified Organization)

Vol. 4, Issue 12, December 2015

By definition $\overline{\dim}_B(S) = \overline{\lim}_{\delta \rightarrow 0} \frac{\log N_\delta(S)}{-\log \delta}$. Since $\delta \rightarrow 0$ if and only if $k \rightarrow \infty$, we have

$$\begin{aligned}\overline{\dim}_B(S) &\leq \overline{\lim}_{k \rightarrow \infty} \frac{\log 2^k}{-\log [2 \prod_{n=1}^k \frac{(1-\epsilon_n)}{2}]} \\ &\leq \overline{\lim}_{k \rightarrow \infty} \frac{\log 2^k}{-\log [2 \prod_{n=1}^k \frac{(1-\epsilon_n)}{2}]^{k/k}} \\ &\leq \overline{\lim}_{k \rightarrow \infty} \frac{k \log 2}{-\log [2 \prod_{n=1}^k \frac{(1-\epsilon_n)}{2}]^{1/k}} \\ \therefore \quad \overline{\dim}_B(S) &\leq \overline{\lim}_{k \rightarrow \infty} \frac{\log 2}{-\log [2 \prod_{n=1}^k \frac{(1-\epsilon_n)}{2}]^{1/k}}\end{aligned}$$

Now to show that $\underline{\dim}_B(S) \geq \underline{\lim}_{k \rightarrow \infty} \frac{\log 2}{-\log \{ [\prod_{n=1}^k \frac{(1-\epsilon_n)}{2}]^{1/k} [\epsilon_{k+1}]^{1/k} \}}$, consider an interval U of length less than δ , $\delta > 0$ and choose k such that

$[\prod_{n=1}^k \frac{(1-\epsilon_n)}{2}] \epsilon_{k+1} \leq \delta < [\prod_{n=1}^{k-1} \frac{(1-\epsilon_n)}{2}] \epsilon_k$, then U intersects at most one of the basic intervals of length $\prod_{n=1}^k \frac{(1-\epsilon_n)}{2}$ used in the construction of S.

For if U intersects more than one interval of S_k then $|U| \geq$ distance between two disjoint intervals of S_k i.e. $|U| \geq [\prod_{n=1}^{k-1} \frac{(1-\epsilon_n)}{2}] \epsilon_k$ i.e. $\delta > [\prod_{n=1}^{k-1} \frac{(1-\epsilon_n)}{2}] \epsilon_k$ which is false because $\delta < [\prod_{n=1}^{k-1} \frac{(1-\epsilon_n)}{2}] \epsilon_k$. These are 2^k such intervals.

\therefore Atleast 2^k intervals of length δ are required to cover $S_k \supset S$.

$\therefore N_\delta(S) \geq 2^k$

Now $\delta \geq [\prod_{n=1}^k \frac{(1-\epsilon_n)}{2}] \epsilon_{k+1}$

$\log \delta \geq \log \{ [\prod_{n=1}^k \frac{(1-\epsilon_n)}{2}] \epsilon_{k+1} \}$

$-\log \delta \leq -\log \{ [\prod_{n=1}^k \frac{(1-\epsilon_n)}{2}] \epsilon_{k+1} \}$

$\frac{1}{-\log \{ [\prod_{n=1}^k \frac{(1-\epsilon_n)}{2}] \epsilon_{k+1} \}} \leq \frac{1}{-\log \delta}$

$\frac{1}{-\log \delta} \geq \frac{1}{-\log \{ [\prod_{n=1}^k \frac{(1-\epsilon_n)}{2}] \epsilon_{k+1} \}}$

By definition $\underline{\dim}_B(S) = \underline{\lim}_{\delta \rightarrow 0} \frac{\log N_\delta(S)}{-\log \delta}$

$$\begin{aligned}&\geq \underline{\lim}_{k \rightarrow \infty} \frac{\log 2^k}{-\log \{ [\prod_{n=1}^k \frac{(1-\epsilon_n)}{2}] \epsilon_{k+1} \}} \\ &\geq \underline{\lim}_{k \rightarrow \infty} \frac{\log 2^k}{-\log \{ [\prod_{n=1}^k \frac{(1-\epsilon_n)}{2}] \epsilon_{k+1} \}^{k/k}} \\ &\geq \underline{\lim}_{k \rightarrow \infty} \frac{k \log 2}{-\log \{ [\prod_{n=1}^k \frac{(1-\epsilon_n)}{2}] \epsilon_{k+1} \}^{1/k}} \\ &\geq \underline{\lim}_{k \rightarrow \infty} \frac{\log 2}{-\log \{ [\prod_{n=1}^k \frac{(1-\epsilon_n)}{2}] \epsilon_{k+1} \}^{1/k}}\end{aligned}$$

International Journal of Innovative Research in Science, Engineering and Technology

(An ISO 3297: 2007 Certified Organization)

Vol. 4, Issue 12, December 2015

$$\therefore \dim_B(S) \geq \liminf_{k \rightarrow \infty} \frac{\log 2}{-\log \{ [\prod_{n=1}^k \frac{(1-\epsilon_n)}{2}]^{1/k} [\epsilon_{k+1}]^{1/k} \}}$$

Weierstrass theorem :

If $\operatorname{Re} z > 0$ and γ denotes Euler's constant then $\frac{1}{\Gamma(z)} = ze^{\gamma z} \prod_{n=1}^{\infty} \left(1 + \frac{z}{n}\right) e^{-z/n}$. This is theorem, § 4.41, p. 148 of [6].

Lemma 2.5 :

$$\prod_{n=1}^{\infty} \frac{(n-a_1)(n-a_2)\dots(n-a_k)}{(n-b_1)(n-b_2)\dots(n-b_j)} = \prod_{i=1}^k \frac{\Gamma(1-b_i)}{\Gamma(1-a_i)}.$$

Proof : Consider

$$\begin{aligned} \frac{(n-a_1)(n-a_2)\dots(n-a_k)}{(n-b_1)(n-b_2)\dots(n-b_k)} &= \frac{n^k \left(1 - \frac{a_1}{n}\right) \left(1 - \frac{a_2}{n}\right) \dots \left(1 - \frac{a_k}{n}\right)}{n^k \left(1 - \frac{b_1}{n}\right) \left(1 - \frac{b_2}{n}\right) \dots \left(1 - \frac{b_k}{n}\right)} \\ &= \frac{\left(1 - \frac{a_1}{n}\right) \left(1 - \frac{a_2}{n}\right) \dots \left(1 - \frac{a_k}{n}\right)}{\left(1 - \frac{b_1}{n}\right) \left(1 - \frac{b_2}{n}\right) \dots \left(1 - \frac{b_k}{n}\right)} \end{aligned} \quad \dots\dots\dots(1)$$

Since $\sum a_i = \sum b_i$, i = 1, 2, 3, ..., k

$$\therefore \sum a_i - \sum b_i = 0, i = 1, 2, 3, \dots, k$$

$$\therefore a_1 + a_2 + \cdots + a_k - b_1 - b_2 - \cdots - b_k = 0$$

$$\therefore e^{\left(\frac{a_1+a_2+\cdots+a_k-b_1-b_2-\cdots-b_k}{n}\right)} = e^0 = 1$$

∴ Equation (1) can be written as

$$\frac{(n-a_1)(n-a_2)\dots(n-a_k)}{(n-b_1)(n-b_2)\dots(n-b_k)} = \frac{\left(\frac{1-a_1}{n}\right)\left(\frac{1-a_2}{n}\right)\dots\left(\frac{1-a_k}{n}\right)}{\left(\frac{1-b_1}{n}\right)\left(\frac{1-b_2}{n}\right)\dots\left(\frac{1-b_k}{n}\right)} e^{\frac{(a_1+a_2+\dots+a_k-b_1-b_2-\dots-b_k)}{n}}$$

$$= \frac{\left(\frac{1-a_1}{n}\right)\left(\frac{1-a_2}{n}\right)\dots\left(\frac{1-a_k}{n}\right)}{\left(\frac{1-b_1}{n}\right)\left(\frac{1-b_2}{n}\right)\dots\left(\frac{1-b_k}{n}\right)} e^{\frac{(a_1+a_2+\dots+a_k)}{n}}$$

$$= \frac{\left(\frac{1-a_1}{n}\right)\left(\frac{1-a_2}{n}\right)\dots\left(\frac{1-a_k}{n}\right)e^{\frac{a_1}{n}}e^{\frac{a_2}{n}}\dots e^{\frac{a_k}{n}}}{\left(\frac{1-b_1}{n}\right)\left(\frac{1-b_2}{n}\right)\dots\left(\frac{1-b_k}{n}\right)e^{\frac{b_1}{n}}e^{\frac{b_2}{n}}\dots e^{\frac{b_k}{n}}}$$

$$= \frac{\left(\frac{1-a_1}{n}\right)e^{\frac{a_1}{n}}\left(\frac{1-a_2}{n}\right)e^{\frac{a_2}{n}}\dots\left(\frac{1-a_k}{n}\right)e^{\frac{a_k}{n}}}{\left(\frac{1-b_1}{n}\right)e^{\frac{b_1}{n}}\left(\frac{1-b_2}{n}\right)e^{\frac{b_2}{n}}\dots\left(\frac{1-b_k}{n}\right)e^{\frac{b_k}{n}}}$$

By Weierstrass theorem

$$\frac{1}{\Gamma(z)} = z e^{\gamma z} \prod_{n=1}^{\infty} \left(1 + \frac{z}{n}\right) e^{-z/n}$$

$$= ze^{\gamma z} \prod_{n=1}^{\infty} \left(\frac{n+z}{n} \right) e^{-z/n}$$

$$\therefore \Gamma(z) = \frac{1}{ze^{zx}} \prod_{n=1}^{\infty} \left(\frac{n}{n+z} \right) e^{-z/n}$$

$$\Gamma(z) = \frac{1}{ze^{\gamma z}} \prod_{n=1}^{\infty} \left(\frac{n}{n+z} \right) e^{z/n}$$

For $z = 1 - b_i$

$$\Gamma(1 - b_i) = \frac{1}{(1 - b_i)e^{\gamma(1 - b_i)}} \prod_{n=1}^{\infty} \left(\frac{n}{n+1 - b_i} \right) e^{\frac{1 - b_i}{n}}$$

For $z = 1 - a_i$ Equation (3) \Rightarrow

Copyright to IJIRSET

DOI:10.15680/IJIRSET.2015.0412109

12279

International Journal of Innovative Research in Science, Engineering and Technology

(An ISO 3297: 2007 Certified Organization)

Vol. 4, Issue 12, December 2015

$$\Gamma(1 - a_i) = \frac{1}{(1-a_i)e^{\gamma(1-a_i)}} \prod_{n=1}^{\infty} \left(\frac{n}{n+1-a_i} \right) e^{\frac{1-a_i}{n}} \quad \dots \dots \dots (5)$$

Equation (4) ÷ Equation (5)

$$\begin{aligned} \frac{\Gamma(1-b_i)}{\Gamma(1-a_i)} &= \frac{(1-a_i)e^{\gamma(1-a_i)} \prod_{n=1}^{\infty} \left(\frac{n}{n+1-b_i} \right) e^{\frac{1-b_i}{n}}}{(1-b_i)e^{\gamma(1-b_i)} \prod_{n=1}^{\infty} \left(\frac{n}{n+1-a_i} \right) e^{\frac{1-a_i}{n}}} \\ &= \frac{(1-a_i)e^{-\gamma a_i}}{(1-b_i)e^{-\gamma b_i}} \prod_{n=1}^{\infty} \left[\left(\frac{n}{n+1-b_i} \right)^{\frac{1-b_i}{n}} e^{\frac{1-a_i}{n}} \right] \\ &= \frac{(1-a_i)}{(1-b_i)} e^{-\gamma(b_i-a_i)} \prod_{n=1}^{\infty} \left(\frac{n+1-a_i}{n+1-b_i} \right) e^{-\frac{1}{n}(a_i-b_i)} \\ \therefore \prod_{i=1}^k \frac{\Gamma(1-b_i)}{\Gamma(1-a_i)} &= \left[\prod_{i=1}^k \frac{(1-a_i)}{(1-b_i)} e^{-\gamma(b_i-a_i)} \right] \prod_{i=1}^k \left[\prod_{n=1}^{\infty} \left(\frac{n+1-a_i}{n+1-b_i} \right) e^{-\frac{1}{n}(a_i-b_i)} \right] \\ &= \left[\prod_{i=1}^k \frac{(1-a_i)}{(1-b_i)} e^{-\gamma(b_i-a_i)} \right] \prod_{n=1}^{\infty} \left[\prod_{i=1}^k \left(\frac{n+1-a_i}{n+1-b_i} \right) e^{-\frac{1}{n}(a_i-b_i)} \right] \\ &= \frac{(1-a_1)(1-a_2)\dots(1-a_k)}{(1-b_1)(1-b_2)\dots(1-b_k)} e^{-\gamma \sum_{i=1}^k (a_i-b_i)} \prod_{n=1}^{\infty} \left[\frac{(n+1-a_1)(n+1-a_2)\dots(n+1-a_k)}{(n+1-b_1)(n+1-b_2)\dots(n+1-b_k)} \right] e^{-\frac{1}{n} \sum_{i=1}^k (a_i-b_i)} \\ &= \frac{(1-a_1)(1-a_2)\dots(1-a_k)}{(1-b_1)(1-b_2)\dots(1-b_k)} \prod_{n=1}^{\infty} \left[\frac{(n+1-a_1)(n+1-a_2)\dots(n+1-a_k)}{(n+1-b_1)(n+1-b_2)\dots(n+1-b_k)} \right] \end{aligned}$$

(∴ $a_i = b_i$, $i = 1, 2, 3, \dots, k \Rightarrow e^{\sum_{i=1}^k (a_i-b_i)} = e^0 = 1$)

Put $n+1 = n'$ in second product

$$\begin{aligned} &= \frac{(1-a_1)(1-a_2)\dots(1-a_k)}{(1-b_1)(1-b_2)\dots(1-b_k)} \prod_{n=2}^{\infty} \left[\frac{(n'-a_1)(n'-a_2)\dots(n'-a_k)}{(n'-b_1)(n'-b_2)\dots(n'-b_k)} \right] \\ &= \prod_{n=1}^{\infty} \left[\frac{(n'-a_1)(n'-a_2)\dots(n'-a_k)}{(n'-b_1)(n'-b_2)\dots(n'-b_k)} \right] \\ \therefore \prod_{i=1}^k \frac{\Gamma(1-b_i)}{\Gamma(1-a_i)} &= \prod_{n=1}^{\infty} \left[\frac{(n-a_1)(n-a_2)\dots(n-a_k)}{(n-b_1)(n-b_2)\dots(n-b_k)} \right] \\ \therefore \prod_{n=1}^{\infty} \frac{(n-a_1)(n-a_2)\dots(n-a_k)}{(n-b_1)(n-b_2)\dots(n-b_k)} &= \prod_{i=1}^k \frac{\Gamma(1-b_i)}{\Gamma(1-a_i)} \end{aligned}$$

Corollary 2.6 : $\prod_{n=1}^{\infty} \frac{(n+1)^p - 1}{(n+1)^p} = \frac{1}{\Gamma(2-\beta_1)\Gamma(2-\beta_2)\dots\Gamma(2-\beta_p)}$, where $\beta_1, \beta_2, \dots, \beta_p$ are p^{th} root of unity with $\beta_1 = 1$.

Proof : Here $(x+1)^p - 1$ is a polynomial in x of degree p . This polynomial has p solutions say $\alpha_1, \alpha_2, \alpha_3, \dots, \alpha_p$. α_j is a solution of $(x+1)^p - 1 = 0$ iff $\alpha_j + 1 = (1)^{1/p}$

i.e. $(\alpha_j + 1)^p = 1$ iff $\alpha_j + 1 = (1)^{1/p}$

iff $\alpha_j + 1$ is p^{th} root of unity

iff $\alpha_j = -1 + \beta_j$, where β_j is the p^{th} root of unity

$$\begin{aligned} \therefore (x+1)^p - 1 &= (x - \alpha_1)(x - \alpha_2) \dots (x - \alpha_p) \\ &= (x + 1 - \beta_1)(x + 1 - \beta_2) \dots (x + 1 - \beta_p) \end{aligned}$$

$$\begin{aligned} \therefore \frac{(n+1)^p - 1}{(n+1)^p} &= \frac{(n+1 - \beta_1)(n+1 - \beta_2) \dots (n+1 - \beta_p)}{(n+1)(n+1) \dots (n+1)} \\ &= \frac{[n - (\beta_1 - 1)][n - (\beta_2 - 1)] \dots [n - (\beta_p - 1)]}{[n - (-1)][n - (-1)] \dots [n - (-1)]} \quad \dots \dots \dots (1) \end{aligned}$$

Here $a_i = \beta_i - 1$ and $b_i = -1$, $i = 1, 2, 3, \dots, p$

Since $\beta_1, \beta_2, \dots, \beta_p$ are p^{th} root of unity $\Rightarrow \beta_1 + \beta_2 + \dots + \beta_p = 0$ (2)

Now $1 - a_i = 1 - (\beta_i - 1) = 1 - \beta_i + 1 = 2 - \beta_i$

$\therefore \operatorname{Re}(1 - a_i) > 0$, $i = 1, 2, 3, \dots, p$ (3)

International Journal of Innovative Research in Science, Engineering and Technology

(An ISO 3297: 2007 Certified Organization)

Vol. 4, Issue 12, December 2015

$$\text{Now } 1 - b_i = 1 - (-1) = 2, \quad i = 1, 2, 3, \dots, p \\ \therefore \operatorname{Re}(1 - b_i) > 0, \quad i = 1, 2, 3, \dots, p \quad \dots\dots\dots(4)$$

$$\text{Now } \sum_{i=1}^p a_i = \sum_{i=1}^p (\beta_i - 1) = \sum_{i=1}^p \beta_i - \sum_{i=1}^p 1 = 0 - p = -p \quad (\text{From equation (2)})$$

$$\therefore \sum_{i=1}^p a_i = \sum_{i=1}^p b_i \quad \dots \dots \dots (5)$$

From equations (3), (4), (5) and by Corollary 2.5 we get,

$$\therefore \prod_{n=1}^{\infty} \frac{(n+1)^p - 1}{(n+1)^p} = \prod_{i=1}^p \frac{\frac{1}{\Gamma(1 - (-1))}}{\Gamma(1 - (\beta_i - 1))} = \prod_{i=1}^p \frac{\frac{1}{\Gamma(2)}}{\Gamma(2 - \beta_i)} = \prod_{i=1}^p \frac{1}{\Gamma(2 - \beta_i)} = \frac{1}{\Gamma(2 - \beta_1) \Gamma(2 - \beta_2) \dots \Gamma(2 - \beta_p)}$$

$$= \frac{1}{\Gamma(2 - \beta_2) \Gamma(2 - \beta_3) \dots \Gamma(2 - \beta_p)} \quad (\because \beta_1 = 1)$$

$$\therefore \prod_{n=1}^{\infty} \frac{(n+1)^p - 1}{(n+1)^p} = \frac{1}{\Gamma(2 - \beta_2) \Gamma(2 - \beta_3) \dots \Gamma(2 - \beta_p)}$$

Corollary 2.7: If ϵ_n is constant sequence or $\epsilon_n = \frac{1}{(n+1)^p}$ where p is a positive integer then

$$\lim_{k \rightarrow \infty} \frac{\log 2}{-\log \{ [\prod_{n=1}^k \frac{(1-\epsilon_n)}{1-\epsilon_{n+1}}]^{1/k} [\epsilon_{k+1}]^{1/k} \}} = \overline{\lim}_{k \rightarrow \infty} \frac{\log 2}{-\log \{ [2 \prod_{n=1}^k \frac{(1-\epsilon_n)}{1-\epsilon_{n+1}}]^{1/k} \}}.$$

Proof : Let ϵ_n be constant. Let $\epsilon_n = \epsilon \forall n$

Now,

From equations (1) and (2) we get,

$$\therefore \lim_{k \rightarrow \infty} \frac{\log 2}{-\log \{ [\prod_{n=1}^k \frac{(1-\epsilon_n)}{2}]^{1/k} [\epsilon_{k+1}]^{1/k} \}} = \lim_{k \rightarrow \infty} \frac{\log 2}{-\log [2 \prod_{n=1}^k \frac{(1-\epsilon_n)}{2}]^{1/k}}$$

Let $\epsilon_n = \frac{1}{(n+1)^p}$ for all n and p is a positive integer.

$$\therefore 1 - \epsilon_n = 1 - \frac{1}{(n+1)^p} = \frac{(n+1)^p - 1}{(n+1)^p}$$

$$\prod_{n=1}^k \frac{(1-\epsilon_n)}{2} = \prod_{n=1}^k \frac{(n+1)^p - 1}{2(n+1)^p} = \frac{1}{2^k} \prod_{n=1}^k \frac{(n+1)^p - 1}{(n+1)^p}$$

$$\left[\prod_{n=1}^k \frac{(1-\epsilon_n)}{2} \right]^{1/k} = \left[\frac{1}{2^k} \prod_{n=1}^k \frac{(n+1)^p - 1}{(n+1)^p} \right]^{1/k}$$

International Journal of Innovative Research in Science, Engineering and Technology

(An ISO 3297: 2007 Certified Organization)

Vol. 4, Issue 12, December 2015

$$\begin{aligned}
& \therefore \left[\prod_{n=1}^k \frac{(1-\epsilon_n)}{2} \right]^{1/k} = \frac{1}{2} \left[\prod_{n=1}^k \frac{(n+1)^p - 1}{(n+1)^p} \right]^{1/k} \\
& \log \left[\prod_{n=1}^k \frac{(1-\epsilon_n)}{2} \right]^{1/k} = \log \left\{ \frac{1}{2} \left[\prod_{n=1}^k \frac{(n+1)^p - 1}{(n+1)^p} \right]^{1/k} \right\} \\
& \therefore \lim_{k \rightarrow \infty} \log \left[\prod_{n=1}^k \frac{(1-\epsilon_n)}{2} \right]^{1/k} = \lim_{k \rightarrow \infty} \log \left\{ \frac{1}{2} \left[\prod_{n=1}^k \frac{(n+1)^p - 1}{(n+1)^p} \right]^{1/k} \right\} \\
& \quad = -\log 2 + \lim_{k \rightarrow \infty} \log \left\{ \left[\prod_{n=1}^k \frac{(n+1)^p - 1}{(n+1)^p} \right]^{1/k} \right\} \\
& \quad = -\log 2 + \lim_{k \rightarrow \infty} \frac{\log \left[\prod_{n=1}^k \frac{(n+1)^p - 1}{(n+1)^p} \right]}{k} \\
& \quad = -\log 2 + \frac{\log \left[\frac{1}{\Gamma(2-\beta_2)\Gamma(2-\beta_3)\dots\Gamma(2-\beta_p)} \right]}{\infty} \\
& = -\log 2 + 0 \tag{By Corollary 2.6}
\end{aligned}$$

$$\therefore \lim_{k \rightarrow \infty} \log \left[\prod_{n=1}^k \frac{(1-\epsilon_n)}{2} \right]^{1/k} = \lim_{k \rightarrow \infty} \log \left\{ \frac{1}{2} \left[\prod_{n=1}^k \frac{(n+1)^p - 1}{(n+1)^p} \right]^{1/k} \right\} = -\log 2 \quad \dots \dots \dots (3)$$

$$\begin{aligned}
\text{Now } \lim_{k \rightarrow \infty} \frac{\log 2}{-\log \left\{ \left[\prod_{n=1}^k \frac{(1-\epsilon_n)}{2} \right]^{1/k} \left[\epsilon_{k+1} \right]^{1/k} \right\}} &= \lim_{k \rightarrow \infty} \frac{\log 2}{-\log \left\{ \left[\prod_{n=1}^k \frac{\left(1 - \frac{1}{(n+1)^p}\right)}{2} \right]^{1/k} \left[\frac{1}{(k+2)^p} \right]^{1/k} \right\}} \\
&= \lim_{k \rightarrow \infty} \frac{\log 2}{-\log \left\{ \left[\prod_{n=1}^k \frac{\left((n+1)^p - 1\right)}{2(n+1)^p} \right]^{1/k} \left[\frac{1}{(k+2)^p} \right]^{1/k} \right\}} \\
&= \lim_{k \rightarrow \infty} \frac{\log 2}{-\log \left\{ \frac{1}{2} \left[\prod_{n=1}^k \frac{(n+1)^p - 1}{(n+1)^p} \right]^{1/k} \left[\frac{1}{(k+2)^p} \right]^{1/k} \right\}} \\
&= \lim_{k \rightarrow \infty} \frac{\log 2}{-\log \left\{ \frac{1}{2} \left[\prod_{n=1}^k \frac{(n+1)^p - 1}{(n+1)^p} \right]^{1/k} \right\} - \log \left[\frac{1}{(k+2)^p} \right]^{1/k}} \\
&= \frac{\log 2}{\frac{-(-\log 2) - 0}{\log 2}} = 1 \quad (\text{F})
\end{aligned}$$

$$\begin{aligned}
\text{Now, } \overline{\lim}_{k \rightarrow \infty} \frac{\log 2}{-\log [2 \prod_{n=1}^k \frac{(1-\epsilon_n)}{2}]^{1/k}} &= \overline{\lim}_{k \rightarrow \infty} \frac{\log 2}{-\log [2 \prod_{n=1}^k \frac{\left(1 - \frac{1}{(n+1)^p}\right)}{2}]^{1/k}} \\
&= \overline{\lim}_{k \rightarrow \infty} \frac{\log 2}{-\log [2 \prod_{n=1}^k \frac{(n+1)^p - 1}{(n+1)^p}]^{1/k}} \\
&= \overline{\lim}_{k \rightarrow \infty} \frac{\log 2}{-\log (2)^{1/k} - \log \left\{ \frac{1}{2} \left[\prod_{n=1}^k \frac{(n+1)^p - 1}{(n+1)^p} \right]^{1/k} \right\}} \\
&= \frac{\log 2}{0 - (-\log 2)} \\
&= \frac{\log 2}{\log 2} = 1
\end{aligned} \tag{From equation(3)}$$

From equations (4) and (5) we get,

From equations (4) and (5) we get,

International Journal of Innovative Research in Science, Engineering and Technology

(An ISO 3297: 2007 Certified Organization)

Vol. 4, Issue 12, December 2015

Remark 2.8 : According to Corollary 2.7 we observe that $\underline{\lim}_{k \rightarrow \infty} \frac{\log 2}{-\log \{ [\prod_{n=1}^k \frac{(1-\epsilon_n)}{2}]^{1/k} [\epsilon_{k+1}]^{1/k} \}} = \overline{\lim}_{k \rightarrow \infty} \frac{\log 2}{-\log [2 \prod_{n=1}^k \frac{(1-\epsilon_n)}{2}]^{1/k}}$ for $\epsilon_n = \text{constant sequence}$ or $\epsilon_n = \frac{1}{(n+1)^p}$ for p is positive integer. $\overline{\dim}_B(S) = \dim_B(S)$.

III CONCLUSION

We have obtained Hausdorff dimension of Cantor like set S constructed in [3] as $\dim_H(S) = \frac{-\log 2}{\gamma}$, where $\gamma = \lim_{k \rightarrow \infty} \log [\prod_{n=1}^k \frac{(1-\epsilon_n)}{2}]^{1/k}$. Also we have estimated the upper / lower Box dimensions of such Cantor like set S. It is proved that $\overline{\lim}_{k \rightarrow \infty} \frac{\log 2}{-\log [2 \prod_{n=1}^k \frac{(1-\epsilon_n)}{2}]^{1/k}} \geq \overline{\dim}_B(S) \geq \dim_B(S) \geq \underline{\lim}_{k \rightarrow \infty} \frac{\log 2}{-\log \{ [\prod_{n=1}^k \frac{(1-\epsilon_n)}{2}]^{1/k} [\epsilon_{k+1}]^{1/k} \}}$. Also we conclude that $\dim_B(S) = \overline{\dim}_B(S)$ for $\epsilon_n = \epsilon$, $n \geq 1$ or $\epsilon_n = \frac{1}{(n+1)^p}$, $n \geq 1$, positive integer p.

REFERENCES

- [1] G.de. Barra, " Measure Theory And Integration", New Age International (p) Limited, 2003.
- [2] Kenneth Falconer, "Fractal Geometry : Mathematical Foundations and Applications" ,John Willey and Sons.1990.
- [3] Dr. S.M. Padhye, Satish B. Khobragade , " On the construction of Cantor like sets", The IJES, Volume 4 ,Issue 4, pages 49-54, 2015.
- [4] J. Shah. Hausdorff dimension and its application. Manuscript.
<http://math.uchicago.edu/~may/VIGRE/VIGRE2009/REUPapers/Shah.pdf>
- [5] Kannan V. , "Cantor set: From Classical To Modern". The Mathematical Student, Vol 63,No.1-4(1994)P.243-257.
- [6] E. C. Titchmarsh , "The Theory Of Functions", Oxford University Press, Oxford Second .Edition -1987.