

International Journal of Innovative Research in Science, Engineering and Technology

(An ISO 3297: 2007 Certified Organization)

Vol. 4, Issue 1, January 2015

Density of States of an Electron Gas in Lower Dimensions

C.Radhika, R..Suganya

Department of Physics, Bharath Institute Of Higher Education And Research, Chennai, India

ABSTRACT : We discuss the detailed theoretical aspectsof the density of states of an electron gas in lower dimensions. This is one of the important parameter for calculating various electronic properties. From this study it is clear that the density of states of an electron gas in one and three dimensions is proportional to Energy and for two dimensions it remains constant.

KEY WORDS: Density of states , Energy.

I. INTRODUCTION

The density of states plays an important role in the kinetic theory of solids [1,4,5,6]. The product of the density of states and theprobability distribution function is the number of occupied states per unit volume at a given energy for a system in thermal equilibrium. This value is widely used to investigate various physical properties of matter[2,3,7]. The following are examples, using two common distribution functions, of how applying a distribution function to the density of states can give rise to physical properties.

Physics of low dimensional system has been a topic of interest from the theoretical point of view for a long time. This is mainly due to the fact that simple models are amenable for an exact solution from which more complicated results may be interpreted.

Some of the well known 1D models in condensed matter physics are

* Kronig penny model for a bond structure of a crystalline solid.

* Lattice vibrations and Phonon Dispersions

* Linear array of interacting spins

Real electrons are 3D but can be made to behave as though they are free to move in fewer dimensions. This can be achieved by trapping them in a narrow potential well that restricts their motion in one dimension to discrete energy levels. If the separation between these energy level is large enough the electron will appear to be in frozen into the ground state and no motion will be possible in this dimension. The result is a 2D electron gas. The same effect can be achieved with a 2D potential well which leaves the electron free to move in 1D only as a quantum wire. [7,8]

II. DENSITY OF STATES

2.1 Density of states of an electron gas in 3D

Density of states is an important parameter which plays a major role in calculating various properties. It is defined as the number of states or orbital per unit energy range.

If N is the total number of orbital with energy $\leq E$ then the Density of states is ,

 $D_3(E) = \frac{dn}{dE}$ where n is the number of states or orbital per unit volume.

DOI: 10.15680/IJIRSET.2015.0401139

www.ijirset.com



International Journal of Innovative Research in Science, Engineering and Technology

(An ISO 3297: 2007 Certified Organization)

Vol. 4, Issue 1, January 2015

Energy is given by $E = \frac{\hbar^2 K^2}{2m}$ If K_f is the Fermi wave vector then the volume is , $V = \frac{4}{3}\pi K_f^3$ In a volume $(\frac{2\pi}{L})^3$ there is one allowed state. Therefore in unit volume volume there will be $(\frac{L}{2\pi})^3$ states. $N = (\frac{L}{2\pi})^{3\frac{4}{3}} K_{f}^{3} 2$ $K_{f} = (3\pi^{2}n)^{\frac{1}{2}}$ $E_f = \hbar^2 K^2 / 2m$ Substituting the value of k, $E_{f} = \hbar^{2}/2m * (3\pi^{2}n)\frac{1}{2}$ $E_f = C n_{\overline{3}}^1$ $dE_{f}/dn = C*\frac{2}{3}(\frac{n-1}{3}) = \alpha n\frac{1}{3}$ Simplifying weget, Density of states at Efis, $D_3 (E_f) = 1/\alpha * C^{1/2} E_f^{1/2}$ $D_3(E) = \frac{\sqrt{2E}m^{\frac{3}{2}}}{\pi^2\hbar^3}$ This is the equation for Density of states of an electron gas in 3D.

2.2 Density of states of an electron gas in 2d

The expression for the Density of states of an electron gas can be divided in a similar way as the 3D provided the volume is replaced by area.[9,10]

Area of Fermi surface = $4\pi K_f^2$ Radius of Fermi surface = $\frac{2\pi}{l}$

In an area $(\frac{2\pi}{L})^2$ there is only one allowed state. Therefore in unit area the number of states will be $(\frac{L}{2\pi})^2$. Number of states $= (\frac{L}{2\pi})^2 4\pi K_f^2$ $K_f = (n\pi)\frac{1}{2}$ Energy is given by $E = \frac{\hbar^2 K^2}{2m}$ $E = \frac{\hbar^2}{2m} n\pi$ dE/dn $= \frac{\hbar^2}{2m} \pi$ Density of states of an electron gas in 2D is, $D_2(E) = 2m/\pi\hbar^2$(2)

This is the equation for Density of states of an electron gas in 2D.

2.3Density of states of an electron gas in 1D

In one dimension, the area of the Fermi surface in 2D is replaced by the radius of the Fermi sphere[11,12] Radius of Fermi surface = $\frac{2\pi}{L}$ In a length $\frac{2\pi}{L}$ there is an allowed state. Number of states per unit length =2n π Energy is given by $E = \frac{1}{3} \frac{K^2}{2m}$ $E = 2 \frac{1}{3} \frac{\pi^2}{n^2} \frac{n^2}{m}$

DOI: 10.15680/IJIRSET.2015.0401139



International Journal of Innovative Research in Science, **Engineering and Technology**

(An ISO 3297: 2007 Certified Organization)

Vol. 4, Issue 1, January 2015

 $dE/dn = 2 \hbar^2 \pi^2 n^2 / m$

Density of states of an electron gas in 1D is,

 $D_1 (E) = 1/C E^{-1/2}$(3)

This is the equation for Density of states of an electron gas in 1D.

III. RESULTS AND DISCUSSIONS

Density of states is one of the important parameter for calculating various electronic properties. Hence the expression for density of states of an electron gas in one, two and three dimensions have been studied. The main results are $D_1(E) \alpha E^{-1/2}$,

 $D_2(E) = Constant$ and $D_3(E) \alpha E^{1/2}$ respectively.

The density of states of an electron gas in three dimensions is derived from the energy of an electron. To find three dimensions its volume is considered and as a result it shows that it is proportional to the squareroot of energy. For two dimensions the volume is replaced by area and here the density of states remains constant. Then for one dimension its area is replaced by the radius and it is inversely proportional to the squareroot of energy.

REFERENCES

- 1. Elangovan and K. Navaneetha Krishnan, Solid state communication 83,635 (1992).
- H.Haug and Sm. Koch, Quantum theory of optical and electronic properties of 2.
- semiconductors, World Scientific, Singapore, 1993.
- Krishnamoorthy P., Jayalakshmi T., "Preparation, characterization and synthesis of silver nanoparticles by using phyllanthusniruri for the 3 antimicrobial activity and cytotoxic effects", Journal of Chemical and Pharmaceutical Research, ISSN: 0975-7384, 4(11) (2012) pp.4783-4794.
- Lakshmi K., Chitralekha S., Illamani V., Menezes G.A., "Prevalence of bacterial vaginal infections in pre and postmenopausal women", 4. International Journal of Pharma and Bio Sciences, ISSN: 0975-6299, 3(4) (2012) pp.949-956.
- Charles Kittel, Introduction to solid state physics, Wiley, VIII Edition, 2004 5.
- Sharmila S., Jeyanthi Rebecca L., Das M.P., "Production of Biodiesel from Chaetomorpha antennina and Gracilaria corticata", Journal of Chemical 6. and Pharmaceutical Research, ISSN: 0975-7384, 4(11) (2012) pp.4870-4874.
- 7 L. Altshuler and A. G. Aronov, Sov. Phys. JETP 50, 968 (1979).
- 8. Rajkumar B., Vijay Kalimuthu B., Rajkumar R., Santhakumar A.R., "Proportioning of recycled aggregate concrete", Indian Concrete Journal, ISSN : 0019-4565, 79(10) (2005) pp.46-50.
- 9. B. L. Altshuler, A. G. Aronov, and P. A. Lee, Phys. Rev. Lett. 44, 1288 (1980).
- 10. Vijayaprakash S., Langeswaran K., Jagadeesan A.J., Rveathy R., Balasubramanian M.P., "Protective efficacy of Terminalia catappa L. leaves against lead induced nephrotoxicity in experimental rats", International Journal of Pharmacy and Pharmaceutical Sciences, ISSN : 0975 - 1491, 4(S3) (2012) pp.454-458.
- 11. D. V. Khveshchenko and M. Reizer, Phys. Rev. B 57, R4245 (1998).
- 12. E. G. Mishchenko and A. V. Andreev, Phys. Rev. B 65, 235310 (2002).
- Dr.V.Krishnasamy, S.Rosy Christy, SUBSTITUTED AROMATIC ANILS A KINETIC STUDY USING PYRIDINIUM CHLOROCHROMATE (13. PCC), International Journal of Biotech Trends and Technology (IJBTT), ISSN: 2319-8753, pp 495-499, Vol. 2, Issue 2, February 2013
- 14. DR.V.Krishnasamy, S.Rosy Christy, KINETICS OF OXIDATION OF AROMATIC ANIL BY PYRIDINIUM DICHROMATE STUDY OF THE EFFECTS OXIDANTS AND SUBSTRATE, International Journal of Biotech Trends and Technology (IJBTT), ISSN: 2319-8753.pp 298-302, Vol. 2, Issue 1, January 2013
- 15. Mrs.Shanthi &Dr.A.Mukunthan, Nucleation kinetics of L-Alanine acetate, International Journal of Innovative Research in Science, Engineering and Technology, ISSN: 2319-8753,pp 2650-2653, Vol. 2, Issue 7, July 2013
- Mrs.V.G.Vijaya, DESIGN AND FABRICATION OF SEQUENCING CIRCUIT WITH SINGLE DOUBLE ACTING CYLINDER, International 16. Journal of Computer Trends and Technology(IJCTT), ISSN: 2319-8753,pp 6315-6322, Vol. 2, Issue 11, November 2013
- Bharthvaian R. A STUDY ON THE EFFECTIVENESS OF SALES INCENTIVES IN IMPROVING SALES PERFORMANCE. International 17. Journal of Computer & Organization Trends (IJCOT), ISSN: 2319-8753,pp 10931-10933, Vol. 3, Issue 4, April 2014