

ISSN(Online): 2319-8753 ISSN (Print): 2347-6710

## International Journal of Innovative Research in Science, Engineering and Technology

(An ISO 3297: 2007 Certified Organization)

Vol. 4, Issue 7, July 2015

# **Trilinearization of (2+1) Dimensional Long Dispersive Wave Equation Using PTA**

P.Myvizhi, R.Velavan

Department of Physics, Bharath Institute of Higher Education and Research, Chennai, India

Department of Physics, Bharath University, Chennai, India

**ABSTRACT:** <u>U</u>sing the methodology of Painleve Truncation Approach (PTA) ,an infinite integrable non linear dynamical system named long dispersive wave equation (2LDW) has been solved and higher order dromion solution is obtained from generalized periodic solution.

**KEYWORDS:** Painleve, trilinear, solitons

### **I.INTRODUCTION**

The identification of a large number of arbitrary functions in the solutions of (2+1) dimensional integrable models have only added to the richness in the structure of them and hence construction of localized excitations in (2+1) dimensions continues to be challenging and rewarding contemporary problem.

In this equation we developed a very simple and straight forward procedure to generate a rather extended class of generic solutions of physical interest. In this 2LDW equation we utilize the local Laurent expansion of the general solution and truncate it at the constant level term (painleve truncation approach) and obtain solutions in terms of arbitrary functions. Through this procedure we generate various periodic an exponentially localized solutions. The novelty here is that the solution is generated through a simple procedure. But the solution obtained is rich in structure because of the arbitrary functions [9-14].

#### **II. PAINLEVE TRUNCATION APPROACH**

If the solution of a nonlinear differential equation is free from movable (integration constant/initial condition dependent) critical singular points (branch points and essential singularities) it is said to possess the painleve property. In this case the system is expected to be integrable.[16-19]

Painleve singularity structure analysis is used in which the solution of the given equation of motion of a dynamical system is analyzed for the presence or absence of movable critical singular points and it can be used as a powerful test to identify and isolate integrable infinite dimensional nonlinear dynamical system. It has been pointed out that there exists a large number of solitons possessing completely integrable infinite dimensional dynamical systems. [20,21]

**Painleve truncation approach** is a methodology in which we are going to prove that the (2+1) dimensional long dispersive wave equation is a completely integrable infinite dimensional nonlinear dynamical system.

The recent interest in the investigation of integrable models in (2+1) dimensions can be attributed to the identification of dromions in the Davey-Stewartson I equation [3]. These dromions which originate at the cross point of the intersection of two nonparallel ghost solitons decay exponentially in all directions and are driven by lower dimensional boundaries or velocity potentials (arbitrary functions) in the system. It must be mentioned that the presence of these so called boundaries enriches the structure of (2+1) dimensional integrable models leading to the formation of localized solutions like dromions, positions, etc. [5]. Thus a judicious harnessing of these lower dimensional arbitrary functions [9-14] of space and time may give rise to new spatio-temporal patterns in higher dimensions besides throwing more light on their integrability.



ISSN(Online): 2319-8753 ISSN (Print): 2347-6710

## International Journal of Innovative Research in Science, Engineering and Technology

(An ISO 3297: 2007 Certified Organization)

#### Vol. 4, Issue 7, July 2015

In this way, a simple procedure is used to solve the (2+1) dimensional system namely Painleve Truncation Approach (PTA) [5]. This approach converts **the given evolution equation into a multi linear equation in terms of non characteristic manifold. This multi linear equation can be solved in terms of lower dimensional arbitrary functions of space and time.** Using this approach one can generate various generalized periodic and localized solutions. Through PTA (2+1) dimensional long dispersive wave equation can be trilinearized. A four parameter special class of solutions involving three arbitrary functions in the variables naturally arises for this **trilinear** equation. Thus, there exists a possibility of obtaining more general solutions.

#### (2+1) DIMENSIONAL LONG DISPERSIVE WAVE EQUATION

Considering the long dispersive wave equation (2LDW) [15,16]

$\lambda_{qt} + q_{xx} - 2_{qv} = 0$	(1a)
$\lambda_{rt} - r_{xx} + 2_{rv} = 0$	(1b)
$(qr)_x = V_\eta$	(1c)

We now effect a local Laurent expansion in the neighbourhood of a noncharacteristic singular manifold  $\Phi(x, \eta, t)=0$ ,  $\Phi_x \neq 0$ ,  $\Phi_\eta \neq 0$ ,  $\Phi_t \neq 0$ .

Assuming the leading orders of eqn(1), the solution of equation (1) to have the form

$q=q_0\Phi^{\mu}$	(2a)
$r = r_0 \Phi^{\beta}$	(2b)
$V = r_0 \Phi^{\gamma}$	(2c)
tic functions of the (re met) and	a R as and integrand to be determined

where  $q_{0,r_0,V_0}$  are analytic functions of  $\Phi(x, \eta, t)$  and  $\alpha$ ,  $\beta$ ,  $\gamma$  are integers to be determined. With the condition of seed solution

$$q_1 = r_1 = 0, V_2 = V_2(t) \tag{3}$$

Then collecting the coefficients of lower order dimensional terms and converting it into a multilinear equation such as trilinear equation is possible as follows.

 $V_{1\eta} = \Phi_{x\eta}^{2} + (\Phi_{x} \Phi_{x\eta\eta}/\Phi_{\eta}) - (\Phi_{x} \Phi_{x\eta} \Phi_{\eta\eta}/\Phi_{\eta}^{2}) - (\Phi_{xx\eta} \Phi_{x\eta}^{2} \Phi_{\eta}) + \Phi_{x} \Phi_{x\eta\eta} \Phi_{\eta} - \Phi_{x} \Phi_{x\eta} \Phi_{\eta\eta} - \Phi_{\eta}^{2} \Phi_{xx\eta} = 0$ (4)

Equation (4) indicates trilinear form of the equation.

#### **III.CONCLUSION**

In this paper, I have investigated the (2+1) dimensional long dispersive wave equation by using the **Painleve Truncation Approach (PTA)** method. In particular, I have shown that the equation is **trilinearizable**. More general solutions can be generated from the trilinear equation which remains, to be investigated. It will enrich the existence of lower dimensional arbitrary functions, which will help us to construct novel localized solutions such as dromions.

#### ACKNOWLEDGEMENT

I am very thankful to honorable chairman Pro-chancellor, Vice Chancellor, Registrar, Dean R&D and Dean Engineering of Bharath University for their continuous support and encouragement given to carry out this research work



## International Journal of Innovative Research in Science, **Engineering and Technology**

(An ISO 3297: 2007 Certified Organization)

#### Vol. 4, Issue 7, July 2015

#### REFERENCES

[1] Lakshmanan M and Rajasekar S. Nonlinear Dynamics Integrability, Chaos and Patterns 2003

[2] Boiti M. Leon J J P, Martina L and Pempinelli F. Scattering of localized solitons in the plane. 1988 Phys. Lett. A 132:432-9.

[3] Fokas A S and Santini P M. Dromions and a boundary value problem for the Davey-Stewartson I equation. Physica D 1990, 44:99-130

[4] Konopelchenko B G 1993 Solitons in Multidimensions (Singapore: World Scientific)

[5] Radha R, Senthil kumar C, Lakshmanan M, Tang X Y, Lou S Y. Periodic and localized solutions of the long wave-short wave resonance interaction equation. J. Phys A 2005;38:9649-63.

[6] Radha R and Lakshmanan M 1994 J. Math. Phys. 35 4746

[7] Radha R and Lakshmanan M 1995 J. Phys. A: Math. Gen. 28 6977

[8] Radha R and Lakshmanan M 1996 J. Phys. A: Math. Gen. 29 1551

[9] Oikawa M, Okamura M and Funakoshi M 1989 J. Phys.Soc.Japan 58 4416

[10] Lai D W C and Chow K W 1999 J. Phys. Soc. Japan 68 1847

[11] Lou S Y 1995 J. Phys. A: Math. Gen. 28 7227

[12] Shanthi B., Revathy C., Devi A.J.M., Parameshwari P.J., Stephen T., "Serum 25(OH)D and type 2 diabetes mellitus", Journal of Clinical and Diagnostic Research, ISSN : 0973 - 709X, 6(5) (2012) pp.774-776.

[13] Madhubala V., Subhashree A.R., Shanthi B., "Serum carbohydrate deficient transferrin as a sensitive marker in diagnosing alcohol abuse: A case - Control study", Journal of Clinical and Diagnostic Research, ISSN: 0973 - 709X, 7(2) (2013) pp.197-200.

[14] Lou S Y 2000 Phys. Lett. A 277 94

[15] Lou S Y and Ruan H Y 2001 J. Phys. A: Math. Gen. 34 305

[16] Tang X Y, Lou S Y and Zhang Y 2002 Phys. Rev. E 66 046601

[17] Khanaa V., Thooyamani K.P., Saravanan T., "Simulation of an all optical full adder using optical switch", Indian Journal of Science and Technology, ISSN: 0974-6846, 6(S6)(2013) pp.4733-4736.

[18] S Chakravarthy, S L Kent and E T Newman, J. Math. Phys. vol.36, 763 (1995)

[19] Nagarajan C., Madheswaran M., "Stability analysis of series parallel resonant converter with fuzzy logic controller using state space techniques", Electric Power Components and Systems, ISSN: 1532-5008, 39(8) (2011) pp.780-793.

[20] Bhat V., "A close-up on obturators using magnets: Part I - Magnets in dentistry", Journal of Indian Prosthodontist Society, ISSN : 0972-4052, 5(3) (2005) pp.114-118.

[21] Radha R and Lakshmanan M, Pramana - J. Phys, 48 163 (1997)

Dr.V.Krishnasamy, S.Rosy Christy, SUBSTITUTED AROMATIC ANILS - A KINETIC STUDY USING PYRIDINIUM [22] CHLOROCHROMATE ( PCC ), International Journal of Biotech Trends and Technology (IJBTT), ISSN: 2319-8753, pp 495-499 , Vol. 2, Issue 2, February 2013

DR.V.Krishnasamy, S.Rosy Christy, KINETICS OF OXIDATION OF AROMATIC ANIL BY PYRIDINIUM DICHROMATE - STUDY [23] OF THE EFFECTS OXIDANTS AND SUBSTRATE, International Journal of Biotech Trends and Technology (IJBTT), ISSN: 2319-8753.pp 298-302, Vol. 2, Issue 1, January 2013

[24] Mrs.Shanthi &Dr.A.Mukunthan, Nucleation kinetics of L-Alanine acetate, International Journal of Innovative Research in Science, Engineering and Technology, ISSN: 2319-8753, pp 2650-2653, Vol. 2, Issue 7, July 2013 [25] Mrs.V.G.Vijaya, DESIGN AND FABRICATION OF SEQUENCING CIRCUIT WITH SINGLE DOUBLE ACTING CYLINDER,

International Journal of Computer Trends and Technology(IJCTT), ISSN: 2319-8753, pp 6315-6322, Vol. 2, Issue 11, November 2013

[26] Bharthvajan R, A STUDY ON THE EFFECTIVENESS OF SALES INCENTIVES IN IMPROVING SALES PERFORMANCE, International Journal of Computer & Organization Trends (IJCOT), ISSN: 2319-8753,pp 10931-10933, Vol. 3, Issue 4, April 2014