

(An ISO 3297: 2007 Certified Organization)

Vol. 4, Issue 6, June 2015

Application of Grey Possibility Degree in Comparing Poverty

Supratim Mukherjee¹, Banamali Ghosh²

Assistant Professor, Department of Mathematics, Nutangram High School, Murshidabad, West Bengal, India¹

Associate Professor, Department of B.Ed, Union Christian training College, Murshidabad, West Bengal, India²

ABSTRACT: The poverty of a certain region is very much context dependent and is of great importance particularly for the rural areas. The nature of the attributes is complex and not easy to determine mathematically. A number of attempts have been raised towards this problem, but lack of consideration of inherent uncertainty draws back most of them. In this paper, the grey possibility degree has been applied to estimate and compare poverty of some regions. A case study is provided to illustrate the effectiveness of applying grey possibility degree. It can be applied to other decision making approaches too.

KEYWORDS: Poverty, Poverty estimation, Fuzzy Logic, Grey Numbers, Grey Possibility Theory.

I. INTRODUCTION

Poverty in rural areas is a fact of real concern to the policy makers. The traditional way of dealing with poverty is not worthy in this case. The attributes considered in their estimation are represented and aggregated using two valued logical algorithms. Uncertainty is present but not considered at all there. So the necessity of an approach that can deal with the inherent uncertainty must be placed. Otherwise the process remains ineffective. Estimating and comparing poverty in rural areas is really very critical and a concrete model of this problem satisfying all the socio economic parameters is somehow missing in the literature. Several authors represented their views on poverty from different angles both theoretically and practically. The main problem is that the theoretical possibilities face such type of problems that cannot be defined or structured in mathematical framework. So the error is there and we are unable to limit it. This work is a new one from the viewpoint of uncertainty. The various aspects of the status of a rural household are not completely deterministic. For example, consider the attribute: income. Normally in Indian rural areas the total income of an individual household varies in a certain range because of the unavailability of monetary engagements at different times. So when a household is asked to state their status corresponding to this attribute, it is quiet unnatural to have a numeric answer, rather we should seek for a linguistic termed answer, e.g., Very Poor, Poor, Medium Poor, Fair, Medium Good, Good, Very Good. The uncertainty is thus considered for each attribute.

Sen (1976) ([8], [9]) rolled over the problem of poverty index in two ways: poor household identification and aggregation of the attributes of the poor in an overall indicator which quantifies the poverty index. The transfer axiom proposed by Sen stipulates that a transfer of income from a poor to a rich person certainly increases the poverty degree when other attributes are fixed. Another axiom in this context is population monotonicity axiom which states that an addition of a person below the poverty line should increase the poverty degree and the addition of a person above the poverty line should decrease the degree when other attributes are fixed. The introduction of these two axioms leads to a certain anomaly regarding the possibility of any poverty index. The poverty index of a society certainly depends on the individual poverty degree of each household. So this problem has been taken into consideration with great importance. Some poverty models are also proposed on the basis of these methods, e.g., the logit model, conditional income distribution model, Fuzzy set theoretic model, etc.

In section II, some preliminaries on grey system are given. In section III, the methodology is illustrated and in section IV a real life example is demonstrated.



(An ISO 3297: 2007 Certified Organization)

Vol. 4, Issue 6, June 2015

II. GREY NUMBERS AND GREY POSSIBILITY DEGREE

Grey theory was first proposed by Deng in 1982. A grey system is a system presented by grey number and variables and it contains uncertain information.

An interval grey number $\otimes G$ is represented by $[\underline{G}, \overline{G}]$ where \underline{G} and \overline{G} are real numbers and may be infinite valued. According to Moore's dissertation ([6]) we state here some algebraic operational laws on interval grey numbers:

 $\otimes G_1 + \otimes G_2 = [\underline{G}_1 + \underline{G}_2, \overline{G}_1 - \overline{G}_2]$ $\otimes G_1 + \otimes G_2 = [\underline{G}_1 + \underline{G}_2, \overline{G}_1 - \overline{G}_2]$ $\otimes G_1 \times \otimes G_2 = [\min(\underline{G}_1\underline{G}_2, \underline{G}_1\overline{G}_2, \overline{G}_1\underline{G}_2, \overline{G}_1\overline{G}_2), \\ \max(G, G_2, G_1\overline{G}_2, \overline{G}_1\overline{G}_2, \overline{G}_1\overline{G}_2)]$

For two intervals grey numbers $\otimes G_1$ and $\otimes G_2$ the possibility degree of $\otimes G_1 \leq \otimes G_2$ can be expressed as ([6]):

$$P\{\otimes G_1 \leq \otimes G_2\} =$$

$$\frac{\max(0, L(\otimes G_1) + L(\otimes G_2) - \max(0, \overline{G}_1 - \underline{G}_2))}{L(\otimes G_1) + L(\otimes G_2)} \text{ where } L(\otimes G) = \overline{G}_1 - \underline{G}_2.$$

The above mentioned grey possibility degree has the following properties.

- i) If $\otimes G_1 = \otimes G_2$, then $P(\otimes G_1 \le \otimes G_2) = 0.5$,
- ii) If $\underline{G}_2 > \overline{G}_1$, then $P(\otimes G_1 \leq \otimes G_2) = 1$,
- iii) If $\overline{G}_2 < \underline{G}_1$, then $P(\otimes G_1 \le \otimes G_2) = 0$,
- iv) When there is some part in the intersection of $\otimes G_1$ and $\otimes G_2$, then $P(\otimes G_1 \leq \otimes G_2) < 0.5$ implies $\otimes G_2$ is smaller than $\otimes G_1$ and $P(\otimes G_1 \leq \otimes G_2) > 0.5$ implies $\otimes G_1$ is smaller than $\otimes G_2$.

We illustrate this by an example.

Consider a situation when two sets of grey numbers (G₁) and (G₂) are compared where (G₁) = {(1, 3), (2, 4), (5, 6)} and (G₂) = {(2, 3), (3, 5), (7, 9)}.

Here
$$P\{(G_1) \le (G_2)\} = \frac{1}{3} \left[\frac{2}{3} + \frac{3}{4} + 1 \right] = 0.80556.$$

Again consider another situation where two sets of grey numbers (G₁) and (G₃) are compared where (G₁) = {(1, 3), (2, 4), (5, 6)} and (G₃) = {(2, 3), (3, 5), (8, 10)}.

Here
$$P\{(G_1) \le (G_3)\} = \frac{1}{3} \left[\frac{2}{3} + \frac{3}{4} + 1 \right] = 0.80556.$$

Here we clearly observe that $P\{(G_1) \leq (G_2)\} = P\{(G_1) \leq (G_3)\}$, but $(G_2) \neq (G_3)$.

III.COMPARISON OF POVERTY- A CASE STUDY

In this work, a case study is presented to compare the status of poverty in four districts of West Bengal, India. The four districts are Murshidabad (D₁), Malda (D₂), Birbhum (D₃) and Burdwan (D₄). The districts are compared according to the attributes considered. This consideration can be extended to any number of attributes and any suitable number of regions for the comparison. Nevertheless this is somehow equivalent to multi-criteria decision making in uncertain environment. Let us consider a discrete set of m Districts, $D = \{D_1, D_2, ..., D_m\}$ and a set of n attributes, $A = \{A_1, A_2, ..., D_m\}$



(An ISO 3297: 2007 Certified Organization)

Vol. 4, Issue 6, June 2015

 A_n . These attributes are additively independent. Also consider $\bigotimes w = \{\bigotimes w_1, \bigotimes w_2, ..., \bigotimes w_n\}$ as an attribute weight vector of the attribute weights which are realized to be linguistic variables. The attributes we consider here in this example are

A1: Housing Condition

A2: Clothing Status

A3: Educational Status

A₄: Average monthly income per capita

A5: Food Security

Now these linguistic weights and attribute ratings can be expressed in grey numbers as shown in table 3.1.

Linguistic Attribute weights		Attribute ratings		
Linguistic	Grey	Linguistic	Grey	
terms	numbers	terms	numbers	
Very Low	[0.0,0.1]	Very Poor	[0,1]	
(VL)		(VP)		
Low (L)	[0.1,0.3]	Poor (P)	[1,3]	
Medium	[0.3,0.4]	Medium Poor	[3,4]	
Low (ML)		(MP)		
Medium	[0.4,0.5]	Fair (F)	[4,5]	
(M)				
Medium	[0.5,0.6]	Medium	[5,6]	
High (MH)		Good (MG)		
High (H)	[0.6,0.9]	Good (G)	[6,9]	
Very High	[0.9,1.0]	Very Good	[9,10]	
(VH)		(VG)		

TABLE 3.1: EXPRESSION OF LINGUISTIC TERMS IN GREY NUMBERS

Step1: A group of k experts $E = \{E_1, E_2, ..., E_k\}$ is formed. The attribute weight $\bigotimes w_j$ of the jth attribute A_j is calculated as

 $\bigotimes w_j = \frac{1}{k} \left[\bigotimes w_j^1 + \bigotimes w_j^2 + \ldots + \bigotimes w_j^k \right]$ where $\bigotimes w_j^k$ is the attribute weight given by the kth Expert, described by the

grey number [$\underline{w}_j^k, \overline{w}_j^k$].

Here in this case study $E = \{E_1, E_2, E_3\}$, i.e., 3 distinct experts are considered to impose their decisions on different aspects in the decision making process.

Step2: The linguistic variables for the ratings to make an attribute rating value are denoted as $\otimes G_{ij}$, defined by the grey numbers $\left[\otimes \underline{G}_{ij}, \otimes \overline{G}_{ij}\right]$. Thus the Grey Decision Matrix (GDM) is established. This matrix is normalized and the

Normalized Grey Decision Matrix (NGDM) is constituted by the elements $\otimes G_{ij}' = \left\lfloor \frac{\otimes \underline{G}_{ij}}{\overline{G}_{j}^{\max}}, \frac{\otimes \overline{G}_{ij}}{\overline{G}_{j}^{\max}} \right\rfloor$ where

 $G_j^{\max} = \max_{1 \le i \le m} \left\{ \bigotimes \overline{G}_{ij} \right\}$. Table 3.2 displays the status of the 4 districts with respect to the 5 attributes.



(An ISO 3297: 2007 Certified Organization)

Vol. 4, Issue 6, June 2015

TABLE 3.2: GREY DECISION MATRIX (G	OM)
------------------------------------	-----

GDM	A ₁	A_2	A ₃	A_4	A ₅
D ₁	VP	Р	MP	Р	F
D_2	F	MP	MP	F	MG
D_3	Р	VP	Р	VP	MG
D_4	MG	F	G	MG	VP

The weights of the attributes are gathered from the experts and displayed in table 3.3.

$\otimes w_j$	D_1	D_2	D ₃	D_4
A_1	Н	MH	М	MH
A_2	ML	L	ML	ML
A ₃	L	ML	L	L
A_4	VH	VH	VH	VH
A ₅	М	М	MH	М

TABLE 3.3: ATTRIBUTE WEIGHT MATRIX (AWM)

Now the average attribute weights are calculated:

 $\otimes w_1 = [0.345, 0.325], \otimes w_2 = [0.45, 0.525], \otimes w_3 = [0.055, 0.15], \otimes w_4 = [0.87, 1], \otimes w_5 = [0.555, 0.625].$ Using (3.2) and (3.3) we construct the Normalized Grey Decision Matrix as shown in table 3.4.

TABLE 3.4: NORMALIZED GREY DECISION MATRIX					
1		T ₂	T ₃	T_4	

NGDM	T ₁	T ₂	T ₃	T_4	T ₅
D1	[0.11, 0.33]	[0.33, 0.44]	[0, 0.11]	[0.33, 0.44]	[0.33, 0.44]
D ₂	[0.33, 0.44]	[0.44, 0.55]	[0.44, 0.55]	[0.33, 0.44]	[0.55, 0.67]
D ₃	[0, 0.11]	[0.11, 0.33]	[0, 0.11]	[0.11, 0.33]	[0.44, 0.55]
D_4	[0.44, 0.55]	[0.44, 0.55]	[0.55, 0.67]	[0.55, 0.67]	[0.11, 0.33]

Step3: Now each normalized elements are multiplied by their corresponding weights and we retrace the weighted NGDM by the elements $\otimes T_{ij} = \otimes G_{ij}' \times w_j$. The weighted normalized grey decision matrix is shown in table 3.5.

WNGDM	T_1	T_2	T ₃	T_4	T ₅
D_1	[0.047,	[0.140, 0.231]	[0, 0.058]	[0.14, 0.231]	[0.14, 0.231]
	0.173]				
D_2	[0.049,	[0.066, 0.179]	[0.066, 0.179]	[0.049, 0.143]	[0.082, 0.218]
	0.143]				
D ₃	[0, 0.027]	[0.008, 0.082]	[0, 0.027]	[0.008, 0.082]	[0.033, 0.137]
D_4	[0.396, 0.55]	[0.396, 0.55]	[0.495,	[0.495, 0.67]	[0.099, 0.33]
			0.0.67]		

TABLE 3.5: WEIGHTED NORMALIZED GREY DECISION MATRIX



(An ISO 3297: 2007 Certified Organization)

Vol. 4, Issue 6, June 2015

Step4: At the final stage of this method, a pseudo alternative, called Comparatively Richest District (CRD) is

constructed by CRD =
$$\left\{ \bigotimes G_1^{\max} = \left[\max_i \underline{T}_{i1}, \max_i \overline{T}_{i1} \right], \dots, \bigotimes G_n^{\max} = \left[\max_i \underline{T}_{in}, \max_i \overline{T}_{in} \right] \right\}$$
.

We use this pseudo alternative to evaluate the ordering. Each district is compared with the CRD by the grey-possibility

degree
$$P\left\{X_i \leq CRD\right\} = \frac{1}{n} \sum_{j=1}^n P\left\{\otimes T_{ij} \leq \otimes G_j^{\max}\right\}.$$

The entries in the NGDM are multiplied by the corresponding attribute weights. Now using (3.4) we construct the WNGDM as:

From the above matrix, we first construct the 'Comparatively Richest District' CRD by using (3.5) as $CRD = \{[0.297, 0.46], [0.46, 0.57], [0.425, 0.77], [0.515, 0.67], [0.34, 0.331]\}$. Finally we use (3.6) to find the grey-possibility degrees:

$$P\left\{D_1 \le CRD\right\} = 0.87 ,$$

$$P\{D_2 \le CRD\} = 0.943,$$

$$P\{D_3 \leq CRD\} = 1,$$

 $P\{D_4 \leq CRD\} = 0.542$.

The ranking is done according to this possibility value and higher possible alternative gets greater rank.

Based on this rank the Government or the policy makers can establish a poverty line separation algorithm. The next section exemplifies the methodology with an alternative proposition.

Now it is the responsibility of the policy makers to draw the line of separation. We propose here to divide the range [0, 1] of possibilistic values in some intervals like [0, 0.3), [0.3, 0.6), [0.6, 0.8), [0.8, 0.9) and [0.9, 1] and separately economic policies should be applied on them. We call the district individuals of these classes as Comparatively Richest, Comparatively Medium Rich, Comparatively Rich, Comparatively Poor and Comparatively Poorest respectively. In our example, the district D_4 belongs to the Comparatively Richest class and the districts D_1 , D_2 and D_3 belong to the Comparatively Poorest Class.

IV. CONCLUSION

In this work, the comparison of poverty status is surveyed with respect to some attributes using grey possibility degrees. The approach is novel and can be extended to any desired number of attributes. The process of applying grey possibility degree has been taken from Li et al.'s approach. The advantage of using this process lies in the fact that it considers uncertainty to some satisfactory extent. Thus the proposed approach is authentic and can be used for other multi attribute decision making problems also.

ACKNOWLEDGEMENT:

This work has been supported by UGC Minor research project No. RSW/108/13-14

References

- [1] Betti G, Verma VK, Measuring the degree of poverty in a dynamic and comparative context; a multidimensional approach using fuzzy set theory, working paper n. 22, Department of Quantitative Methods, University of Siena, 1998
- [2] Chakravarty SR, Ethically flexible measures of poverty, Canadian Journal of Economics, vol. 16, pp 74-85, 1983a
- [3] Chakravarty SR, A new index of poverty, Mathematical Social Sciences, vol. 6, pp 307-313, 1983b
- [4] Cheli B, Lemmi A, A "Totally" fuzzy and relative approach to the multidimensional analysis of poverty, Economic Notes, vol. 24, pp 115-134, 1995
- [5] Kundu A, Smith TE, An impossibility theorem on poverty indices, International Economic Review, vol. 24, pp 423-434, 1983

- [7] Sen AK, Poverty: on ordinal approach to measurement, Econometrica, vol. 44, pp 219-231, 1976
- [8] Sen AK, Issue in the measurement of poverty, Scandinavian Journal of Economics, vol. 81, pp 285-307, 1979

^[6] Guo-Dong L, Dasiuke Y, Masatake N, A Grey-based decision-making approach to the supplier selection problem, Mathematical and Computer Modelling, vol. 46, pp 573-581, 2007



(An ISO 3297: 2007 Certified Organization)

Vol. 4, Issue 6, June 2015

- [9] D Dutta Majumder, R Bhattacharyya, S. Mukherjee, Methods of Evaluation and Extraction of membership Functions- Review with a new Approach", pp 277-281, ICCTA'07, IEEE Computer Press, London
- [10] S Mukherjee, S Kar, Philosophy of Fuzzy Mathematics, Journal of Computer and Mathematical Sciences, vol. 1, issue 4, pp 497-500, 2010
- [11] Hagenaars AJM, A Class of Poverty Indices. International Economic Review, vol. 28, pp 583-607, 1987
- [12] A de Janvry, E. Sadoulet, World Poverty and the Role of Agricultural Technology: Direct and Indirect Effects, The Journal of Development Studies, vol. 38:4, pp 1-26
- [13] Cerioli A, Zani S, A Fuzzy approach to the Measurement of Poverty, In: Dagum C, Zenga M (eds) Income and Wealth Distribution, Inequality and Poverty, Springer-Verlag, New York, pp 272-284, 1990
- [14] Vero J, Werquin P, Un reexamen de la mesure de la pauvrete. Comment s'en sortent les jeunes en phase d'insertion professionnelle? Economie et Statistique ,308-309-310, pp 143-148, 1997