

(An ISO 3297: 2007 Certified Organization)

Vol. 4, Issue 10, October 2015

Examination of Proper Randomness of the Numbers Generated by Rand Corporation (1955) Random Number Table: t-test

Brajendra Kanta Sarmah¹, Dhritikesh Chakrabarty²,

Associate Professor, Department of Statistics, Bholanath College, Dhubri, Assam, India¹

Associate Professor, Department of Statistics, Handique Girls' College, Assam, India²

ABSTRACT: Proper randomness of the numbers generated by Rand Corporation has been tested by B.K. Sarmah, D. Chakrabarty and N. Barman, IJESM 5 (1) : January – March, 2015, 97-119 by applying the Chi-square test for testing the significance of difference between observed frequency of each of the digit in the table and the corresponding theoretical (expected) frequency .

In this paper, the randomness of the digits have been tested by applying t-test for amount of deviation of the observed number of occurrences and the theoretical (expected) number of occurrences of the respective digits and hence the numbers.

The test shows that the numbers generated by Rand Corporation deviated significantly in most the observations from proper randomness.

KEYWORDS: Random number generated by Rand Corporation, student t-test, testing of randomness.

I. INTRODUCTION

Drawing of random sample has been found to be vital or basic necessity in most of the researches and investigations especially of applied sciences. The convenient practical method of selecting a random sample consists of the use of Table of Random Numbers. Existing tables of random numbers, used commonly, are the ones due to Fisher and Yates (Constructed in 1938), L.H.C. Tippett (Constructed in 1927), Kendall and Babington Smith (Constructed in 1939) and Rand Corporation (constructed in 1955)

The random number tables have been subjected to various statistical tests of randomness. These tests have limitations to decide on proper randomness of the numbers occurring in the corresponding tables. As a Consequence it is not guaranteed that the numbers in each of these tables are properly random. This leads to think of testing the proper randomness of the numbers in the tables. In the present study, an attempt has been made to test this. The study, here, has been made on the testing of randomness of the table of numbers constructed by *Rand Corporation* only. By the existing statistical methods, it is only possible to know whether the randomness of the numbers of a table is proper. It is only possible to know whether the deviation of the degree of its randomness is significant. In order to test the proper randomness of the random numbers table constructed by *Rand Corporation*, t-test has been applied. The test shows that *Rand Corporation*'s Random numbers table deviate significantly from proper randomness.

II. MATERIALS AND METHODS

Rand Corporation random number table consists of a total of one million random digits grouped into 2,00,000 sets of 5 digited random numbers. Here a sample of 50000 digites are considered out of one million digits for testing Randomness.

To know whether the number in random numbers table of **Rand Corporation** are proper or not student's t-test for amount of deviation is applied.

Let 'd' be the variable denoting the measure of the deviation (amount of deviation) of the observed number of occurrences of the respective digit.

Suppose, di(i=0,1....9) are independent observed values of the deviation variables.



(An ISO 3297: 2007 Certified Organization)

Vol. 4, Issue 10, October 2015

If the table of number is random then di=0, for all i, in the ideal situation. However, due to chance error, di may assume non zero value.

Thus the values of di's are due to chance error but not due to any assignable error if the table is random. The chance variables are i.i.d. $N(o,\sigma)$ variables.

Thus testing of randomness is equivalent to testing of the hypothesis H_0

That $E(d_i) = 0$, for all i,

Let us consider the statistic t for testing H_o i.e. $t = \frac{\overline{d} - E(\overline{d})}{S.E.(\overline{d})} \sim t_{n-1}$

where,

 $\overline{d} = \frac{1}{n} \sum_{i=1}^{n} d_i$ We have,

$$E(\overline{d}) = \frac{1}{n} \sum E(d_i) = 0$$

When H_o is true

Also,

 $var(\overline{d}) = \frac{\sigma^2}{n}, \ \sigma^2$ is unknown However unbiased estimate of σ^2 is $s^2 = \frac{1}{n-1} \sum (d_i - \overline{d})^2$

 $= \frac{1}{n-1} \left[\Sigma d_i^2 - \frac{(\Sigma d_i)^2}{n} \right]$ Which implies unbiased estimate of

 $var (\overline{d}) = \frac{s^2}{n}$ and S.D. $(\overline{d}) = \frac{s}{\sqrt{n}}$ Therefore statistic t for testing H_o becomes

 $t = \frac{\overline{d}}{s/\sqrt{n}}$ when H_o is true and this t follows student's t distribution with (n-1) d.f.

III. **STEPS IN THE METHOD**

In order to test the proper randomness of the numbers of Rand Corporation table one is required to proceed with the following steps:

- In the first step, observe the occurrences of the digits 0 to 9 for first 2000 trails, second 2000 trials Step1: up to 25^{th} 2000 trails as shown in the table.
- In the second step, compute the theoretical expected frequencies. This is done by dividing trails i.e. 1st Step2: 2000 and 25th 2000 trails by 10 assuming that the digit 0 to 9 occurs equal number of times.
- Step 3: In the third step, compute the amount of deviation of observed occurrences of digits and expected occurrences of digits.
- Step 4: In the fourth step, compute the value of student's t for each of the trails.
- Step 5: Compare the value of t-statistic with corresponding theoretical (expected) values.
- Step 6: Draw conclusion as per the result obtained in step 5.

IV. RESULTS AND DISCUSSION

The results obtained on operating the steps (Nos. 1 to 5) on the random numbers table constructed by Rand *Corporation* have been observed. It is observed from the table that occurrences of digit 0 to 9 are not equal.

V. CONCLUSION

From the table prepared for observed frequency of occurrence of digits along with respective expected frequency (shown in bracket) it is observed that the calculated value of t is significant in all most of the cases except



(An ISO 3297: 2007 Certified Organization)

Vol. 4, Issue 10, October 2015

 3^{rd} , 7^{th} , 20^{th} , 23^{rd} , 25th 2000 trails. That is calculated value of 't' have been found to be significant on comparing them with the corresponding theoretical values for most of the cases (trails).

Hence it may be concluded that the table of numbers constructed by *Rand Corporation* deviates significantly from proper randomness.

Therefore Rand Corporation Random Numbers Table Cannot be treated as properly random, as per results obtained by applying t-test.

TABLE

Observed frequency of occurrence of digits along with the respective expected frequency (Shown in bracket), amount of deviation (di) and the values of students 't' statistic from *Rand Corporation*.

Digits	0	1	2	3	4	5	6	7	8	9	Value of t
1 st 2000	6 206 (200)	16 184(20 0)	16 184(20 0)	0 200(20 0)	24 224(200)	11 211(200)	3 197(20 0)	4 204(200)	16 184(200)	6 206(20 0)	4.04
2 nd 2000	3 197 (200)	19 219 (200)	15 215 (200)	6 206 (200)	15 185 (200)	19 181 (200)	8 192 (200)	5 205 (200)	0 200(200)	0 200 (200)	4.21
3 rd 2000	2 202 (200)	11 189 (200)	3 203 (200)	15 215 (200)	26 226 (200)	2 198 (200)	17 183 (200)	5 195 (200)	2 202 (200)	13 187 (200)	3.21*
4 th 2000	8 192 (200)	24 224 (200)	18 182 (200)	11 189 (200)	24 224 (200)	4 196 (200)	5 205 (200)	1 199 (200)	8 192 (200)	3 197 (200)	4.00
5 th 2000	7 207 (200)	7 207 (200)	14 186 (200)	25 175 (200)	13 213 (200)	7 193 (200)	10 210 (200)	14 214 (200)	2 202 (200)	7 193 (200)	5.00
6 th 2000	8 208 (200)	7 193 (200)	11 189 (200)	15 215 (200)	16 216 (200)	3 203 (200)	12 188 (200)	12 188 (200)	3 203 (200)	3 197 (200)	6.91
7 th 2000	4 196 (200)	4 196 (200)	8 192 (200)	7 207 (200)	1 199 (200)	0 200 (200)	21 221 (200)	11 211 (200)	21 179 (200)	1 199 (200)	3.28*
`8 th 2000	19 181 (200)	11 211 (200)	19 219 (200)	11 189 (200)	5 195 (200)	17 217 (200)	10 210 (200)	11 189 (200)	7 193 (200)	4 196 (200)	7.27
9 th 2000	9 191 (200)	0 200 (200)	1 199 (200)	10 210 (200)	10 210 (200)	9 191 (200)	19 219 (200)	1 199 (200)	20 180 (200)		3.59
10 th 2000	11 189 (200)	1 199 (200)	19 181 (200)	4 196 (200)	10 190 (200)	16 184 (200)	16 216 (200)	22 222 (200)	15 215 (200)	8 208 (200)	5.53
11 th 2000	31 169 (200)	3 197 (200)	11 189 (200)	8 192 (200)	6 194 (200)	27 227 (200)	13 187 (200)	18 218 (200)	5 205 (200)	22 222 (200)	4.11
12 th 2000	8 108 (200)	0 200 (200)	9 191 (200)	9 209 (200)	9 191 (200)	21 221 (200)	2 198 (200)	28 228 (200)	29 171 (200)	17 183 (200)	3.60
13 th 2000	26 174 (200)	12 188 (200)	23 177 (200)	9 209 (200)	18 218 (200)	15 185 (200)	0 200 (200)	8 192 (200)	34 234 (200)	23 223 (200)	4.65



(An ISO 3297: 2007 Certified Organization)

Vol. 4, Issue 10, October 2015

14^{th}	11	15	9	3	1	2	12	9	8	6	
2000	211	185	209	197	201	198	188	209	200	194	4.85
2000	(200)	(200)	(200)	(200)	(200)	(200)	(200)	(200)	(200)	(200)	
15 th 2000	15	16	29	2	7	16	9	7	1	2	
	215	194	171	202	207	216	191	207	199	198	3.91
	(200)	(200)	(200)	(200)	(200)	(200)	(200)	(200)	(200)	(200)	
16 th 2000	17	12	6	2	3	13	5	8	6	0	
	217	188	194	202	203	187	195	208	206	200	4.80
	(200)	(200)	(200)	(200)	(200)	(200)	(200)	(200)	(200)	(200)	
17 th 2000	14	23	22	17	9	2	7	2	12	0	
	214	177	222	183	209	202	207	198	188	200	4.63
	(200)	(200)	(200)	(200)	(200)	(200)	(200)	(200)	(200)	(200)	
18 th 2000	3	14	8	11	1	15	1	10	9	2	
	203	186	192	211	201	185	201	210	209	202	4.56
	(200)	(200)	(200)	(200)	(200)	(200)	(200)	(200)	(200)	(200)	
19 th 2000	25	15	32	30	3	5	1	9	17	3	
	175	215	232	170	197	205	201	191	217	197	3.91
	(200)	(200)	(200)	(200)	(200)	(200)	(200)	(200)	(200)	(200)	
20 th 2000	2	3	2	6	1	7	2	3	15	13	
	202	203	198	194	199	207	202	197	185	201	3.12*
	(200)	(200)	(200)	(200)	(200)	(200)	(200)	(200)	(200)	(200)	
21 st	25	25	16	8	4	29	9	22	14	20	
21 ^{sr} 2000	175	225	216	192	196	171	209	222	214	180	5.75
	(200)	(200)	(200)	(200)	(200)	(200)	(200)	(200)	(200)	(200)	
22 nd	7	11	6	7	14	2	5	20	6	14	
2000	207	211	194	207	214	202	205	180	194	186	4.32
	(200)	(200)	(200)	(200)	(200)	(200)	(200)	(200)	(200)	(200)	
23 rd 2000	23	6	34	2	2	0	3	0	10	8	
	223	206	166	198	202	200	203	200	210	192	2.25*
	(200)	(200)	(200)	(200)	(200)	(200)	(200)	(200)	(200)	(200)	
24 th 2000	18	19	14	11	1	7	1	8	29	4	
	182	219	214	211	199	193	199	208	171	204	3.96
	(200)	(200)	(200)	(200)	(200)	(200)	(200)	(200)	(200)	(200)	
o ~ th	4	3	9	29	6	3	14	1	14	1	
25 th 2000	196	203	209	171	194	197	214	201	214	201	3.14*
	(200)	(200)	(200)	(200)	(200)	(200)	(200)	(200)	(200)	(200)	
	(-00)	(_00)	(-00)	(-00)	(-00)	(-00)	(-00)	(-00)	(-00)	(-00)	

*Indicates the values which are less than theoretical values of t $_{0.01,8}$ =3.36

REFERENCES

- 1.
- Cochran, W.G. (1940): "Sample survey", Cambridge University press. Chakrabarty, D. (2004): "A theoretical definition of probability based on common sense", Bulletin of pure and applied sciences, 23 E-2, 343-2. 349.
- Chakrabarty, D. (2005): "Probability;Link between classical definition and Empirical definition", J. Ass. Sc. Soc., 45,13-18. 3.
- Chakrabarty, D. (2006): "Non equally likely outcomes: The classical definition of probability", Bulletin of pure and applied sciences, 25, E-2, 4. 471-481.
- 5. Chakrabarty, D. (2007): "Empirical definition of probability : Special case of its Theoretical definations". Int. J. Agricult. Stats. Sci., 3(1), 261-267.
- Chakrabarty, D. (2008): "Bernolli's definition of probability: Special case of its Chakrabarty definition", Int. J. . Agricult. Stats. Sci., 4(1), 23-6. 267.
- 7. Cochran (1940): "A survey of Experimental design", Mimeo, U.S.D.A.
- Fisher, R. A. (1938): Statistical table for biological, agricultural and medical research, 6th edition (1982), Longman group limilted, England, 8. 37-38 & 134-139.
- 9. Kendall, M.G. and B. B. Simith (1938): Randomness and Random sampling numbers, Jour. Roy Sat. Soc., 103, 147-166.



(An ISO 3297: 2007 Certified Organization)

Vol. 4, Issue 10, October 2015

- 10. Rand Corporation (1955): A million random digits free press, Glenoe, III.
- 11. Snedecor, G.W. and W. G. Cochran (1967): "Statistical methods, Lowa state university press, Ames, Lowa, 6th edition.
- 12. Tippet (1927). "Random number tables" tracts of computer, No.-15, Cambridge University press.
- 13. Von Mises R. (1939): "Probability, Statistics and Truth" Macmillian.
- 14. Sarmah, B.K and Chakrabarty, D (2014): "Testing of Randomness of Number generated by Fisher and Yates" (Chi-Square test), IJESRT, 3(11), 632-636.
- 15. Sarmah, B.K and Chakrabarty, D (2014): " Examination of Proper Randomness of the Number Generated by L.H.C Tippett, (Chi-Square test)JESRT, 3(12), 661-668.
- 16. Sarmah, B.K and Chakrabarty, D (2015): Examination of Proper Randomness of the Number Generated by Kendall and Babington Smith (Chi-square test). IJESRT 4(2) Febuary/15, 260-282.
- 17. B.K. Sarmah*, D. Chakrabarty, N. Barman (2015): Examination of Proper Randomness of the Number Generated by Rand Corporation (1955) (Chi-square test), IJESM 5(1): January- March, 2015, 97-119.
- 18. Sarmah, B.K and Chakrabarty, D (2015): Testing of proper Randomness of the numbers generated by Fisher and Yates (Applying t-test) ABJMI Vol-7 issue –I, Jan- June 2015, P 87-90.
- 19. Sarmah, B.K. and Chakrabarty D. (2015): Examination of proper Randomness of the numbers generated by L.H.C. Tippett (1927)(t-test) IOSR Journal of Mathematics, Vol-11, Issue-3 May-June, 2015 PP.35-37