

## International Journal of Innovative Research in Science, Engineering and Technology

*(An ISO 3297: 2007 Certified Organization)*

**Vol. 4, Issue 9, September 2015**

# Some Common Fixed Point Theorems in S-Metric Space

Tanmoy Mitra

Asst. Teacher, Akabpur High School, Akabpur, Nadanghat, Burdwan, West Bengal, India

**ABSTRACT:** In 2012 S. Sedghi, N. Shobe and A. Aliouche have introduced the concept of S-metric space. In this paper some common fixed point results for two mappings has been established in S-metric space.

**KEYWORDS:** S-metric space, Fixed point.

**AMS Subject Classification (2010):** 47H10, 54H25.

### I. INTRODUCTION AND PRELIMINARIES

**Definition 1 [1]:** Let  $X$  be a nonempty set. An S-metric on  $X$  is a function  $S: X^3 \rightarrow [0, \infty)$  that satisfies the following conditions for all  $x, y, z, a \in X$

- (S1)  $S(x, y, z) = 0$  if and only if  $x = y = z$ .
- (S2)  $S(x, y, z) \leq S(x, x, a) + S(y, y, a) + S(z, z, a)$ .

The pair  $(X, S)$  is called an S-metric space.

The concept of S-metric space is a generalization of a G-metric space [11] and a D\*- metric space [14].

**Definition 2 [2]:** Let  $X$  be a nonempty set. A b-metric on  $X$  is a function  $d: X^2 \rightarrow [0, \infty)$  such that, for a real number  $s \geq 1$ , the following conditions hold for all  $x, y, z \in X$ .

- (i)  $d(x, y) = 0$  if and only if  $x = y$ .
- (ii)  $d(x, y) = d(y, x)$ .
- (iii)  $d(x, z) \leq s[d(x, y) + d(y, z)]$ .

The pair  $(X, d)$  is called a b-metric space.

**Definition 3 [1]:** Let  $(X, S)$  be an S-metric space. For  $r > 0$  and  $x \in X$ , we define the open ball  $B_S(x, r)$  and the closed ball  $B_S[x, r]$  with centre  $x$  and radius  $r$  as follows

$$\begin{aligned} B_S(x, r) &= \{y \in X: S(y, y, x) < r\}, \\ B_S[x, r] &= \{y \in X: S(y, y, x) \leq r\}. \end{aligned}$$

**Definition 4 [1]:** Let  $(X, S)$  be an S-metric space.

- (1) A sequence  $\{x_n\}$  in  $X$  is said to converge  $x$ , if  $S(x_n, x_n, x) \rightarrow 0$  as  $n \rightarrow \infty$ . That is for any  $\varepsilon > 0$ , there exists  $N \in \mathbb{N}$  such that for all  $n \geq N$ , we have  $S(x_n, x_n, x) < \varepsilon$ .
- (2) A sequence  $\{x_n\}$  in  $X$  is a Cauchy sequence if  $S(x_n, x_n, x_m) \rightarrow 0$  as  $m, n \rightarrow \infty$ . That is for every  $\varepsilon > 0$ , there exists  $N \in \mathbb{N}$  such that for all  $m, n \geq N$ ,  $S(x_n, x_n, x_m) < \varepsilon$ .
- (3) The S-metric space  $(X, S)$  is complete if every Cauchy sequence is a convergent sequence.

**Lemma 5 [1]:** In an S-metric space, we have  $S(x, x, y) = S(y, y, x)$ ,  $\forall x, y \in X$ .

**Lemma 6 [1]:** Let  $(X, S)$  be an S-metric space. If  $x_n \rightarrow x$  and  $y_n \rightarrow y$  then

$$S(x_n, x_n, y_n) \rightarrow S(x, x, y).$$

# **International Journal of Innovative Research in Science, Engineering and Technology**

*(An ISO 3297: 2007 Certified Organization)*

**Vol. 4, Issue 9, September 2015**

## II. MAIN RESULTS

**Theorem 1:** Let  $U, V$  be two self-maps on a complete S-metric space  $(X, S)$  and

$S(Ux, Ux, Vy) \leq aS(x, x, y) + bS(Ux, Ux, y) + cS(Vy, Vy, x)$ ,  $\forall x, y \in X$  and  $a, b, c$  are non-negative reals with,  $a + b + c < 1$ ,  $a + 3b < 1$ , and  $a + 3c < 1$ .

Then the mappings  $U, V$  have a unique common fixed point in  $X$ .

**Proof:** Let  $x_0 \in X$ .

Let us define a sequence  $\{x_n\}$  in  $X$  as

$$x_{2k+1} = Ux_{2k}, \quad x_{2k+2} = Vx_{2k+1}, \quad k = 0, 1, 2, \dots$$

$$\begin{aligned} \text{Now } S(x_{2k+1}, x_{2k+1}, x_{2k+2}) &= S(Ux_{2k}, Ux_{2k}, Vx_{2k+1}) \\ &\leq aS(x_{2k}, x_{2k}, x_{2k+1}) + bS(Ux_{2k}, Ux_{2k}, x_{2k+1}) + \\ &\quad cS(Vx_{2k+1}, Vx_{2k+1}, x_{2k}) \\ &= aS(x_{2k}, x_{2k}, x_{2k+1}) + bS(x_{2k+1}, x_{2k+1}, x_{2k+1}) + \\ &\quad cS(x_{2k+2}, x_{2k+2}, x_{2k}) \end{aligned}$$

$$\begin{aligned} &\leq \\ &aS(x_{2k}, x_{2k}, x_{2k+1}) + c[2S(x_{2k+2}, x_{2k+2}, x_{2k+1}) + \\ &\quad S(x_{2k}, x_{2k}, x_{2k+1})] \quad [\text{By S2}] \\ &= aS(x_{2k}, x_{2k}, x_{2k+1}) + c[2S(x_{2k+1}, x_{2k+1}, x_{2k+2}) + S(x_{2k}, x_{2k}, x_{2k+1})] \quad [\text{By Lemma 5}] \end{aligned}$$

Thus,

$$S(x_{2k+1}, x_{2k+1}, x_{2k+2}) \leq \frac{a+c}{1-2c} S(x_{2k}, x_{2k}, x_{2k+1}).$$

Let  $\frac{a+c}{1-2c} = h_1$ . Since  $0 \leq a + 3c < 1$ ,  $0 \leq h_1 < 1$ .

Thus

$$S(x_{2k+1}, x_{2k+1}, x_{2k+2}) \leq h_1 S(x_{2k}, x_{2k}, x_{2k+1}).$$

Again,

$$\begin{aligned} S(x_{2k+2}, x_{2k+2}, x_{2k+3}) &= S(Vx_{2k+1}, Vx_{2k+1}, Ux_{2k+2}) \\ &= S(Ux_{2k+2}, Ux_{2k+2}, Vx_{2k+1}) \\ &\leq aS(x_{2k+2}, x_{2k+2}, x_{2k+1}) + bS(Ux_{2k+2}, Ux_{2k+2}, x_{2k+1}) + \\ &\quad cS(Vx_{2k+1}, Vx_{2k+1}, x_{2k+2}) \\ &= aS(x_{2k+2}, x_{2k+2}, x_{2k+1}) + bS(x_{2k+3}, x_{2k+3}, x_{2k+1}) + \\ &\quad cS(x_{2k+2}, x_{2k+2}, x_{2k+2}) \\ &\leq aS(x_{2k+2}, x_{2k+2}, x_{2k+1}) + b[2S(x_{2k+3}, x_{2k+3}, x_{2k+2}) + \\ &\quad S(x_{2k+1}, x_{2k+1}, x_{2k+2})] \\ &= aS(x_{2k+1}, x_{2k+1}, x_{2k+2}) + b[2S(x_{2k+2}, x_{2k+2}, x_{2k+3}) + \\ &\quad S(x_{2k+1}, x_{2k+1}, x_{2k+2})] \end{aligned}$$

Thus,

$$S(x_{2k+2}, x_{2k+2}, x_{2k+3}) \leq \frac{a+b}{1-2b} S(x_{2k+1}, x_{2k+1}, x_{2k+2})$$

Let  $\frac{a+b}{1-2b} = h_2$ . Since  $0 \leq a + 3b < 1$ ,  $0 \leq h_2 < 1$ .

Then  $S(x_{2k+2}, x_{2k+2}, x_{2k+3}) \leq h_2 S(x_{2k+1}, x_{2k+1}, x_{2k+2})$

# **International Journal of Innovative Research in Science, Engineering and Technology**

(An ISO 3297: 2007 Certified Organization)

Vol. 4, Issue 9, September 2015

Take  $h = \max\{h_1, h_2\}$ .

Now

$$S(x_{2k+1}, x_{2k+1}, x_{2k+2}) \leq h S(x_{2k}, x_{2k}, x_{2k+1})$$

And also

$$S(x_{2k+2}, x_{2k+2}, x_{2k+3}) \leq h S(x_{2k+1}, x_{2k+1}, x_{2k+2})$$

Thus

$$S(x_{n+1}, x_{n+1}, x_{n+2}) \leq h S(x_n, x_n, x_{n+1}) \leq h^2 S(x_{n-1}, x_{n-1}, x_n) \leq \dots \dots \dots \leq h^{n+1} S(x_0, x_0, x_1)$$

Now for  $m > n$ ,

$$\begin{aligned}
S(x_n, x_n, x_m) &\leq 2S(x_n, x_n, x_{n+1}) + S(x_{n+1}, x_{n+1}, x_m) \\
&\leq 2S(x_n, x_n, x_{n+1}) + 2S(x_{n+1}, x_{n+1}, x_{n+2}) + S(x_{n+2}, x_{n+2}, x_m) \\
&\leq \dots \\
&\leq 2S(x_n, x_n, x_{n+1}) + 2S(x_{n+1}, x_{n+1}, x_{n+2}) + \dots + \\
&\quad 2S(x_{m-2}, x_{m-2}, x_{m-1}) + S(x_{m-1}, x_{m-1}, x_m) \\
&\leq 2[S(x_n, x_n, x_{n+1}) + S(x_{n+1}, x_{n+1}, x_{n+2}) + \dots + S(x_{m-1}, x_{m-1}, x_m)] \\
&\leq 2[h^n + h^{n+1} + \dots + h^{m-1}]S(x_0, x_0, x_1) \\
&\leq \frac{2h^n}{1-h}S(x_0, x_0, x_1) \\
&\rightarrow 0 \text{ as } n \rightarrow \infty.
\end{aligned}$$

Thus  $\{x_n\}$  is a Cauchy sequence.

Now since  $(X, S)$  is complete, there exists  $t \in X$ , such that  $x_n \rightarrow t$  as  $n \rightarrow \infty$ .

Now,

$$\begin{aligned}
S(Ut, Ut, t) &\leq 2S(Ut, Ut, Vx_{2k+1}) + S(t, t, Vx_{2k+1}) \\
&\leq 2[aS(t, t, x_{2k+1}) + bS(Ut, Ut, x_{2k+1}) + cS(Vx_{2k+1}, Vx_{2k+1}, t)] + S(t, t, x_{2k+2}) \\
&= 2[aS(t, t, x_{2k+1}) + bS(Ut, Ut, x_{2k+1}) + cS(x_{2k+2}, x_{2k+2}, t)] + \\
&\quad S(x_{2k+2}, x_{2k+2}, t)
\end{aligned}$$

Letting  $k \rightarrow \infty$ , we get,

$$S(Ut, Ut, t) \leq 2bS(Ut, Ut, t)$$

Thus,  $(1 - b)S(Ut, Ut, t) < 0$ .

Since  $0 \leq a + 3b < 1$ ,  $S(Ut, Ut, t) = 0$ .

Thus  $Ut = t$ .

$$\begin{aligned} \text{Also, } S(t, t, Vt) &= S(Ut, Ut, Vt) \\ &\leq aS(t, t, t) + bS(Ut, Ut, t) + cS(Vt, Vt, t) \\ &\quad - cS(t, t, Vt) \end{aligned}$$

Thus  $(1 - c)S(t, t; Vt) \leq 0$

Since  $0 \leq a + 3c \leq 1$ ,  $S(t, t, Vt) = 0$ .

Thus  $Vt = t$

Therefore  $UIt = Vt = t$

i.e.  $t$  is a common fixed point of  $U$  and  $V$ .

For uniqueness let  $t^* \in X$  such that

$$U t^* = V t^* = t^*$$

## **International Journal of Innovative Research in Science, Engineering and Technology**

*(An ISO 3297: 2007 Certified Organization)*

**Vol. 4, Issue 9, September 2015**

$$\begin{aligned}
\text{Now, } S(t, t, t^*) &= S(Ut, Ut, Vt^*) \\
&\leq aS(t, t, t^*) + bS(Ut, Ut, t^*) + cS(Vt^*, Vt^*, t) \\
&= aS(t, t, t^*) + bS(t, t, t^*) + cS(t^*, t^*, t) \\
&= aS(t, t, t^*) + bS(t, t, t^*) + cS(t, t, t^*)
\end{aligned}$$

Thus,  $(1 - a - b - c)S(t, t, t^*) \leq 0$ .

Since  $0 \leq a + b + c < 1$ ,  $S(t, t, t^*) = 0$ .

Thus  $t = t^*$ .

Thus  $t$  is the unique common fixed point of  $U$  and  $V$ .

**Corollary 1.1:** Let  $T$  be a self-maps on a complete S-metric space  $(X, S)$  and

$S(Tx, Tx, Ty) \leq aS(x, x, y) + bS(Tx, Tx, y) + cS(Ty, Ty, x)$ ,  $\forall x, y \in X$ , and  
 $a, b, c$  are non-negative reals with  $a + b + c < 1$ ,  $a + 3b < 1$ ,  $a + 3c < 1$ .

Then the mapping  $T$  has a unique fixed point in  $X$ .

**Proof:** Setting  $U = V = T$  in the above theorem, we get the result.

**Theorem 2:** Let  $(X, S)$  be a complete S-metric space. Let  $U, V$  be two self-maps on  $X$ , satisfying

$$S(Ux, Ux, Vy) \leq aS(x, x, y) + \frac{bS(Ux, Ux, y)S(Vy, Vy, x)}{1+S(x, x, y)}, \text{ where } 0 \leq a + b < 1.$$

Then  $U, V$  have a unique common fixed point in  $X$ .

**Proof:** Let  $x_0 \in X$ .

Let us define a sequence  $\{x_n\}$  in  $X$  as

$$x_{2k+1} = Ux_{2k}, \quad x_{2k+2} = Vx_{2k+1}, \quad k = 0, 1, 2, \dots \dots \dots$$

$$\begin{aligned}
\text{Now } S(x_{2k+1}, x_{2k+1}, x_{2k+2}) &= S(Ux_{2k}, Ux_{2k}, Vx_{2k+1}) \\
&\leq aS(x_{2k}, x_{2k}, x_{2k+1}) + \frac{bS(Ux_{2k}, Ux_{2k}, x_{2k+1})S(Vx_{2k+1}, Vx_{2k+1}, x_{2k})}{1+S(x_{2k}, x_{2k}, x_{2k+1})} \\
&= aS(x_{2k}, x_{2k}, x_{2k+1}) + \frac{bS(x_{2k+1}, x_{2k+1}, x_{2k+1})S(x_{2k+2}, x_{2k+2}, x_{2k})}{1+S(x_{2k}, x_{2k}, x_{2k+1})} \\
&= aS(x_{2k}, x_{2k}, x_{2k+1})
\end{aligned}$$

Again,

$$\begin{aligned}
S(x_{2k+2}, x_{2k+2}, x_{2k+3}) &= S(Vx_{2k+1}, Vx_{2k+1}, Ux_{2k+2}) \\
&= S(Ux_{2k+2}, Ux_{2k+2}, Vx_{2k+1}) \\
&\leq aS(x_{2k+2}, x_{2k+2}, x_{2k+1}) + \frac{bS(Ux_{2k+2}, Ux_{2k+2}, x_{2k+1})S(Vx_{2k+1}, Vx_{2k+1}, x_{2k+2})}{1+S(x_{2k+2}, x_{2k+2}, x_{2k+1})} \\
&= aS(x_{2k+1}, x_{2k+1}, x_{2k+2}) + \frac{bS(x_{2k+3}, x_{2k+3}, x_{2k+1})S(x_{2k+2}, x_{2k+2}, x_{2k+2})}{1+S(x_{2k+2}, x_{2k+2}, x_{2k+1})} \\
&= aS(x_{2k+1}, x_{2k+1}, x_{2k+2})
\end{aligned}$$

Now,

$$S(x_{n+1}, x_{n+1}, x_{n+2}) \leq aS(x_n, x_n, x_{n+1}) \leq a^2S(x_{n-1}, x_{n-1}, x_n) \leq \dots \dots \dots \leq a^{n+1}S(x_0, x_0, x_1).$$

Now,

$$S(x_{n+1}, x_{n+1}, x_{n+2}) \leq aS(x_n, x_n, x_{n+1}) \leq a^2S(x_{n-1}, x_{n-1}, x_n) \leq \dots \dots \dots \leq a^{n+1}S(x_0, x_0, x_1).$$

Now, for  $m > n$ ,

$$S(x_n, x_n, x_m) \leq 2S(x_n, x_n, x_{n+1}) + S(x_m, x_m, x_{n+1})$$

## International Journal of Innovative Research in Science, Engineering and Technology

(An ISO 3297: 2007 Certified Organization)

**Vol. 4, Issue 9, September 2015**

$$\begin{aligned}
 &\leq 2[S(x_n, x_n, x_{n+1}) + S(x_{n+1}, x_{n+1}, x_{n+2}) + \dots + S(x_{m-1}, x_{m-1}, x_m)] \\
 &\leq 2[a^n + a^{n+1} + \dots + a^{m-1}]S(x_0, x_0, x_1) \\
 &\leq \frac{2a^n}{1-a}S(x_0, x_0, x_1) \\
 &\rightarrow 0, \text{ as } n \rightarrow \infty \quad [\text{Since } 0 \leq a + b < 1.]
 \end{aligned}$$

Thus  $\{x_n\}$  is a Cauchy sequence.

Since  $(X, S)$  is complete, there exists  $t \in X$ , such that  $x_n \rightarrow t$  as  $n \rightarrow \infty$ .

Now,

$$\begin{aligned}
 S(Ut, Ut, t) &\leq 2S(Ut, Ut, x_{2k+2}) + S(t, t, x_{2k+2}) \\
 &= 2S(Ut, Ut, Vx_{2k+1}) + S(t, t, x_{2k+2}) \\
 &\leq 2 \left[ aS(t, t, x_{2k+1}) + \frac{bS(Ut, Ut, x_{2k+1})S(Vx_{2k+1}, Vx_{2k+1}, t)}{1+S(t, t, x_{2k+1})} \right] + S(t, t, x_{2k+2}) \\
 &= 2 \left[ aS(t, t, x_{2k+1}) + \frac{bS(Ut, Ut, x_{2k+1})S(x_{2k+2}, x_{2k+2}, t)}{1+S(t, t, x_{2k+1})} \right] + S(t, t, x_{2k+2})
 \end{aligned}$$

Letting  $k \rightarrow \infty$  and using the lemma 5 and lemma 6, we get,

$$S(Ut, Ut, t) \leq 0$$

Hence  $Ut = t$ .

Again,

$$\begin{aligned}
 S(t, t, Vt) &= S(Ut, Ut, Vt) \\
 &\leq aS(t, t, t) + \frac{bS(Ut, Ut, t)S(Vt, Vt, t)}{1+S(t, t, t)} \\
 &= 0
 \end{aligned}$$

Thus,  $S(t, t, Vt) = 0$  and hence  $Vt = t$ .

Therefore,  $Ut = Vt = t$ .

i.e.  $t$  is a common fixed point of  $U$  and  $V$ .

For Uniqueness, let  $t^* \in X$ , such that  $Ut^* = Vt^* = t^*$ .

Thus,

## International Journal of Innovative Research in Science, Engineering and Technology

(An ISO 3297: 2007 Certified Organization)

Vol. 4, Issue 9, September 2015

$$\begin{aligned} S(t, t, t^*) &= S(Ut, Ut, Vt^*) \\ &\leq aS(t, t, t^*) + \frac{bS(Ut, Ut, t^*)S(Vt^*, Vt^*, t)}{1+S(t, t, t^*)} \\ &= aS(t, t, t^*) + \frac{bS(t, t, t^*)S(t^*, t^*, t)}{1+S(t, t, t^*)} \\ &\leq aS(t, t, t^*) + bS(t, t, t^*) \end{aligned}$$

Thus,  $(1 - a - b)S(t, t, t^*) \leq 0$ .

Since  $0 \leq a + b < 1$ ,  $S(t, t, t^*) = 0$ .

Thus  $t = t^*$ .

Hence  $t$  is the unique common fixed point of  $U$  and  $V$ .

### III. CONCLUSION

Many other common fixed point results for two mappings, satisfying different types of rational inequalities in S-metric space can also be proved similarly.

### REFERENCES

- [1] S. Sedghi, N. Shobe, A. Aliouche, A generalization of fixed point theorem in S-metric spaces, Mat. Vesnik 64 (2012), 258 – 266.
- [2] I. A. Bakhtin, The contraction principle in quasimetric spaces, Func. An., Ulianowsk, Gos. Ped. Ins. 30 (1989), 26 – 37.
- [3] S. Sedghi, N. V. Dzung, Fixed Point Theorems on S-metric Spaces, Mat. Vesnik (01.11.2012).
- [4] R. Kannan, Some results on fixed points II, Amer. Math. Monthly 76 (1969), 405 – 408.
- [5] S. Sedghi, N. Shobe, H. Zhou, A Common fixed point theorem in \* metric spaces, Fixed Point Theory Appl. 2007 (2007), 1 – 13.