

International Journal of Innovative Research in Science, Engineering and Technology

(A High Impact Factor, Monthly, Peer Reviewed Journal)

Visit: www.ijirset.com

Vol. 7, Issue 12, December 2018

Summing Up Water Characteristics with Software-Analysed Trophic Interrelations in a Pond

Rekha Kumari¹, Nitu Kumari¹, Jainendra Kumar²

M.Sc Scholar, Department of Biotechnology, College of Commerce, Arts & Science, Patna

(Patliputra University, Patna), India¹

Professor & Former Head, Department of Biotechnology, College of Commerce, Arts & Science, Patna

(Patliputra University, Patna), India²

ABSTRACT: All the living creatures in the world and their health depend on the availability of quality water for drinking and other purposes. This report is based on the studies which aimed to evaluate the water quality of a model pond in Bodh Gaya (Bihar), India that has been used by the population for different uses apart from aquaculture since long. Investigation of pond water samples from the body at varying seasons was taken up regarding parameters such as P^H, temperature, conductivity, dissolved oxygen, B.O.D., Turbidity, T.D.S, Transparency, Total hardness, Calcium, Magnesium, Phosphate, Nitrate and Chlorine contents to generate data for EwE (*Ecopath with Ecosim*) based analysis of the ecosystem network, growth of components, fish growth potentiality and population dynamics.

I. INTRODUCTION

Water plays an important role in the development of healthy society. Ecological network analysis is a system based methodology to analyze within the system and its interacting components used to identify holistic properties that are otherwise not evident from the direct observations. Like any analytical technique, the accuracy of the results is as good as the data available but the additional challenge is that the data need to characterize an entire ecosystem's flows and storages (Fath *et al*, 2007).

While 70% surface of the earth is covered by water, majority of water available on the earth is saline and only 3% of the total only exists as fresh water. Fresh water has become a scarce commodity due to over exploitation and pollution (Ghosh and Bose 1968 : Gupta and Shukla 2006, Patil and Tijare 2001, Singh and Mathur, 2005). Since the International Biological programs (IBP) emphasis on ecosystem research around 1970s, the focus of many ecological studies has been on describing flows in the ecosystems and a number of methodologies were developed for measuring trophic interactions between groups of organisms in different types of ecosystem (Edmondson and Winberg, 1971, Holme and McIntyre, 1971, Bagenal, 1978; Fashan, 1984; Christensen and Pauly, 1992). A highly useful software Ecopath II was developed by Christensen and Pauly (1992) for balancing steady state ecosystem model and calculating network characteristics. It was further refined year after year for application in various ecosystems especially for aquatic ecosystems as Ecopath with Ecosim (EwE) (Zhang *et al*; 2011).

Before we go for EwE application for network analysis, it is imperative to assess and analyze water characteristics to correlate the pressure dynamics with the network interrelations.

II. MATERIALS AND METHOD


Sample collection – Water samples were collected from the *Dibrahi* Pond in plastic bottles in winter, summer and rainy seasons. Samples were taken during 07:00 to 08:00 am in the morning from the depth of 30 cm below the surface, brought to the laboratory and stored at 4⁰ c for further analysis.

International Journal of Innovative Research in Science, Engineering and Technology

(A High Impact Factor, Monthly, Peer Reviewed Journal)

Visit: www.ijirset.com

Vol. 7, Issue 12, December 2018



III. RESULT AND DISCUSSION

P^H measurement – The P^H was measured by P^H meter. The P^H values varied from 7 to 8 in all the three season i.e winter, summer and rainy season.

IV. TEMPERATURE

Temperature varies in the range of between 19^0C to 32^0C in all the three season i.e winter, summer and rainy season. The temperature of water directly influence some chemical reactions in aquatic ecosystem the significant corresponding . Temperature is basically an important factor for its effects on chemical and biological reactions in water. Taken temp record on the spot, immediately by dipping the thermometer for about one minute.

Table1 presents the quality parameters of the pond.

Table no. 1

Sl. No.	Water quality parameter	Season		
		Winter	Summer	Rainy
1.	Temp (^0C)	19^0C	32^0C	26^0C
2.	P^H	7	8	8
3.	Conductivity (mohs/cm)	0.5	0.7	0.9
4.	D.O	3.0	3.5	5.8
5.	B.O.D. (mg/l)	7.0	7.5	5.4
6.	Transparency (cm)	33.5	42.3	29.6
7.	Turbidity (NTU)	44	35	52
8.	Total hardness (mg/l)	420	490	350
9.	Calcium (mg/l)	100.2	140.5	65.4
10.	Magnesium (mg/l)	221	146.4	140
11.	Phosphate (mg/l)	1	0	1
12.	Nitrate (mg/l)	1	1	1
13.	T.D.S (mg/l)	275	233	320
14.	Chlorine (mg/l)	68	76	42.4

International Journal of Innovative Research in Science, Engineering and Technology

(A High Impact Factor, Monthly, Peer Reviewed Journal)

Visit: www.ijirset.com

Vol. 7, Issue 12, December 2018

CONDUCTIVITY

Conductivity is the measure of the capacity of a substance or solution to conduct electric current. Conductivity is reciprocal of the resistance conductivity if measured of a cub with each side of 1 cm at 25⁰ C is called specific conductance. The conductivity varies in the range of 0.5 to 2.1 in season wise.

DISSOLVED OXYGEN

Dissolved oxygen is one of the most important parameters in water quality assessment and coordinate the physical and biological processes prevailing in the water. The obtained range of D.O was found 3 – 5.8. It was highest in the rainy season and lowest in winter season. Higher D.O could be increased due to the solubility of oxygen at the lower temperatures.

B.O.D

B.O.D – The amount of oxygen consumed during microbial utilization of organics is called the Biochemical oxygen demand (Howard S. Peavy). B.O.D is a measure of the dissolved oxygen consumed by micro organisms during the oxidation of reduced substances in water. An inverse relationship was found between the dissolved oxygen concentrations and B.O.D value (coscun .I 1989). The obtained range of B.O.D was found 5.4 – 7.5. Highest range was found in summer and lowest in rainy season.

TRANSPARENCY

Transparency is a measure of the depth of light penetration into the water. It depends on the amount of particles in water. Hence, it directly and visually indicates the pollution level. The obtained range was found 29 – 42.3, highest in summer and lowest in rainy season.

TURBIDITY

Suspension of particles of water interfering with the passage of light is turbidity (Hemlata Mahobe, 2013). The range of turbidity varied between 35 to 52. NTU. The highest range was observed in rainy and lowest during summer.

TOTAL HARDNESS

Total hardness is a measure of the calcium and magnesium concentrations in water. Its range was found in the range of 350 to 490. The maximum range was found in summer and the minimum range in rainy season respectively. Upadhyaya, 2013 was reported total hardness was high during summer similar result were observed in the present study.

CALCIUM

It is very important for the vertebrates of the pond likes fishes, crabs etc. The obtained range was 65 to 140.5. The lowest range was found in rainy and the highest range was found in summer.

MAGNESIUM

Plants may loose colour due to lack of magnesium. It was found in the range of 140 to 221 lowest in rainy and highest in winter.

International Journal of Innovative Research in Science, Engineering and Technology

(A High Impact Factor, Monthly, Peer Reviewed Journal)

Visit: www.ijirset.com

Vol. 7, Issue 12, December 2018

PHOSPHATE

It is essential for plant life but too much addition of phosphate can speed up eutrophication. The maximum value was 1 and minimum value was 0.

NITRATE

Nitrate value was 1 in all winter, summer and rainy seasons. TDS is an aggregate indication of the presence of a broad array of chemical contaminants. Its range was highest in rainy season *i.e.* 320 and lowest in winter *i.e.* 275 respectively.

CHLORINE CONTENT

Chloride is one of the most vital inorganic ions in water as it is highly soluble. Its range was 42 – 76 found lowest during rainy and highest in summer season.

V. CONCLUSION

Conclusion – A healthy society can developed by the use of water. Pond water is very essential part since there is lack of water on earth. Range of all water quality parameters I suitable for the survival of aquatic plants and animal. Chlorine content may affect the life under pond but P^H and temperature favours them. Hence, the water supports a healthy environment for this ecosystem.

ACKNOWLEDGEMENT

Acknowledgement - It is very grateful for the another to get the facilities and materials from University Grant Commission, Govt. of Bihar, Collage of Commerce Arts and Science Patna, H.O.D of Botany Department, M.U. Bodhgaya, Lab In charge of water sewage Treatment plant, Anishabad, Patna and Angel Scientific Store, Patna. My investigation has been done successfully by their support under the guidance of Dr. Jainendra Kumar.

REFERENCES

1. B.B. Ghose and A.K. Basu 1968 observation on estuarine pollution of the Hooghly by the effluents from a chemical factory complex at Rashasa , West Bengal Env. Health 10:29-218.
2. S. Gupta and D.N. Shukla (2006), physicochemical analysis of sewage water and its effect on seed germination and seedling growth of sesamum indicum. J Matras Development 1:5-19.
3. Bagenal T. 1978 Method for assessment of fish production in fresh waters. Third edition IBP handbook No.3 Black well scientific publication oxford 365 p.
4. Christensen, V. and Pauly, D. 1992. Eco-Path II- a software for balancing steady state models and calculating network characteristics . ecol Modeling 61:169-185.
5. Edmondson, W.T. and winberg, G. [Eds.] 1971 A manual on methods for the assessment of secondary productivity in fresh water.
6. Zhang, T. Su Y. and Ma, S. 2011 preliminary study of ecopath with ecosim to the shrimp pond ecosystem.
7. Brian D. fath, B.D. scharler, V.M, Ulanowicz, R.E. and Hannon, B.2007 Ecological network analysis network construction.
8. K. Coscum.I, Yurteris. ,Mirat. T. and Gural. D(1989). Removal of dissolved organic concentration by ozonation, Env. Progress, 6(4), 240-244.
9. Howard S. Peavy, Donald R. Rowe, Environmental Engineering, Mcgro-Hill International edition.
10. Hemlata Mahobe 2013, study of Physico-chemical characteristics of water pond (2013) Journal of scientist.
11. M. Upadhyay & Vijay Laxmi Gupta (2013). Analysis of water quality using physic-chemical parameters.