



e-ISSN: 2319-8753 | p-ISSN: 2347-6710

IJIRSET

International Journal of Innovative Research in
SCIENCE | ENGINEERING | TECHNOLOGY



INTERNATIONAL JOURNAL OF INNOVATIVE RESEARCH

IN SCIENCE | ENGINEERING | TECHNOLOGY

Volume 10, Issue 7, July 2021

ISSN INTERNATIONAL
STANDARD
SERIAL
NUMBER
INDIA

Impact Factor: 7.569



9940 572 462



6381 907 438



ijirset@gmail.com



www.ijirset.com

Review on Epilepsy and Its Management

KM Manisha¹, Ravi Verma², Dr. Kaushal Kishore Chandrul³

Student of B.Pharm 4th yr, Department of Pharmacy, Mewar University, Chittaurgarh, India¹

Assistant Professor, Department of Pharmacy, Mewar University, Chittaurgarh, India²

Principal, Department of Pharmacy, Mewar University, Chittaurgarh, India³

ABSTRACT : By and by a day, people are standing up to various kinds of stress in the fast step by step life and most of the people on earth are encountering a diverse neurological issue. Epilepsy is one of the most broadly perceived neurologic issues of the cerebrum, impacting around 50 million individuals worldwide and 90% of them are from making countries. Inherited calculates similarly as malady cerebrum, stroke, tumor, and high fever cause epilepsy. It powers an immense monetary load on social protection structures of countries related to disrespect and abuses the patient and even his/her family in the system, working condition, and school, and home. Various patients with epilepsy persevere through genuine enthusiastic difficulty, direct issues, and unprecedented social separation. There are numerous seizure types and various instruments by which the mind creates seizures. The goal of the audit is to talk about history, the study of disease transmission, etiology, pathophysiology, an order of epilepsy, symptoms, analysis, the executives of epilepsy, and its future patterns.

KEYWORDS : Epilepsy, Seizures, epidemiology, Anti-epileptic drugs, pathophysiology, hypersynchrony

I. INTRODUCTION

About 1% of the all-out masses are affected by Epilepsy, making it second to stroke as one of the most broadly perceived real neurological issues. 90% of the impacted region from making countries which counts around 50 million people the world over. In the past a serious drawn-out period, our understanding of epilepsy has extended in a couple of respects. Epilepsy is the lopsidedness between cerebral affectability and limitation is tipped toward uncontrolled tenseness and portrayed by discontinuous seizures. Epilepsy is comprehensive with no age, sex, geographical or racial boundaries. Some kinds of epilepsy are restricted to explicit periods of youthfulness, the turmoil isn't enduring. Drugs join treatment with anticonvulsants. Anyway, the best open medications don't control seizures of over 30% of the people. The reason for this review is to give general contemplations and the board of epilepsy [1-3].

II. CLASSIFICATION OF EPILEPTIC SEIZURES[4-9]

Based on commission on classification and terminology of the international league against epilepsy table, 1 represents the international classification of epileptic seizures and then it got approved in Sept 1981

During the seizures based on both a clinical expression of the seizures and the electroencephalographic picture the epileptic seizure classification is made Table [1-5]

I. Generalised Seizures	II. Partial Seizures
A. Generalised tonic-clonic seizures. B. Absence Seizures. C. Atonic Seizures. D. Myoclonic seizures. E. Infantile spasms (Hypsarrhythmia).	A. Simple partial Seizures. B. Complex partial seizures. C. Simple partial or complex partial seizures.

III.EPIDEMIOLOGY

Epilepsy is a serious neurological disorder.[8-9].It is estimated about 55 lacks is affected in India 20 lacks in USA and 3 lacks in the UK. At least 8% of the total population will have one seizure but not epilepsy the age of incidence is 44 per 1 lacks people in a year 10% of the patient take at least one antiepileptic drug (ADE)[10-11].

The National Sentinel and it of epilepsy-related death led by epilepsy Bereaved drew attention to this problem[12].

I. RISK FACTOR[13]

- Start of seizures after the age of 12
- Family account of seizures
- Irrepressible seizures for 2-6 years
- Raised periodical of seizures episode(>30)
- Head injuries
- Record of absence seiuires

CAUSE OF EPILEPSY

The reason for epilepsy is unknown. The phrase epilepsy does not suggest whatever about the cause or severity of the character's seizures, some cases of epilepsy is triggered with the aid of genetic factors, but it may additionally result from mind accidents because of blows to the head, stroke, infections, high fever or tumor[14] It has been observed that heredity (genetics) play a critical position in several cases of epilepsy in very young children, however, it can be an issue for humans of any age. For instances, not every person who has extreme head damage (a clean cause of seizures) will increase epilepsy[15].

PATHOPHYSIOLOGY

Seizure results when an unexpected awkwardness happens among excitatory and inhibitory synapses inside the system of cortical neurons. The fundamental physiology of a seizure scene is an unsteady cell film or its encompassing cells. The seizure begins from the dark matter of any cortical or subcortical region. At first, few neurons fire anomalous. Typical film conductance and the inhibitory synaptic current breakdown and expanded edginess spread either locally to deliver a central seizure or all the more broadly to create a summed up seizure. This beginning spreads by physiologic pathways towards nearby areas[16]. Seizures happen from transient irregular synchronization of neurons in the mind which upsets typical examples of neuronal correspondence and results in electrical releases in the EEG. This interruption can create different indications and signs that rely upon the site of the cause of the seizure and its connections. Within the epileptic center, seizures originate from redoubled excitation or shriveled forbiddance, based on a hypothesis that involves acts between two neurons where the activity of the second neuron has a measurable outcome[13/17]. During an acquiring, responsibility for slaying flow to the wit increases to take off CO₂ and to fetch the substrate for the metabolous manifestation of the neurons. As the seizures prolong, the intelligence suffers statesman from anemia that resulting neuronal finish and intelligence damage[3/18].

DIAGNOSIS

Variables of the study contains dependent and independent variable. The study used pre-specified method for the selection of variables. The study used the Stock returns are as dependent variable. From the share price of the firm the Stock returns are calculated. Rate of a stock salable at stock market is known as stock price Neuroimaging examines assume a necessary job in the evaluation of seizures for the assurance of the auxiliary and utilitarian etiology of seizures. Various analytic devices are used to distinguish and order the seizure type and etiology, including electroencephalogram (EEG), Magnetic Resonance Imaging (MRI), and Positron Emission Tomography (PET), single-photon discharge processed tomography (SPECT), Magneto Encephalogram (MEG), and neuropsychiatric testing[19] The EEG regularly is basic for recognizing explicit seizure types. CT output may help in assessing newly analyzed patients, however, MRI is liked. An X-ray may find mind sores or anatomic deformities that are missed by regular radiographs or CT filters[20] EEG is mostly useful in the diagnosis of various types of seizures.

MANAGEMENT

The terms anticonvulsant and antiepileptic are utilized to bury variably. An anticonvulsant is a specialist that squares tentatively created seizures in research center creatures and the antiepileptic drug is a drug utilized restoratively to control the epilepsies.[21]

.PRINCIPLES OF MANAGEMENT[22]

Epilepsy is a perpetual condition requiring a mindful long stretch of the board. The treatment is puzzling, including game plan and end, assurance, and see if the best possible antiepileptic administrator and assessment of the picked drug's side effects and medicine coordinated efforts. Since these side effects increase when meds are joined, monotherapy is recommended. Long stretch the officials' issues and perfect medicine decision shift from patient to tenacious. In case seizures are not compelled by sedate, the patient may have been misdiagnosed or misclassified. Resistance, a huge explanation behind clear dormancy to treatment, should moreover be blocked. Seeing that current treatment isn't ideal for certain patients, new pharmacologic and cautious medicines are immediately discussed.[23].

The antiepileptic drugs have been classified as follows in Table 2[23-25]

Seizure type	Conventional anti-seizure drug
I. Partial seizures	
(i) Simple partial	Carbamazepine Phenytoin Phenobarbital
(ii) Complex partial	Phenobarbital , Phenytoin Primidone, Carbamazepine Valproate, Carbamazepine Phenobarbital, Phenytoin
(iii) Partial with secondly generalized tonic clonic seizure	
II. Generalized seizure	
(i) Absence Seizures	Clonazepam Ethosuximide Valproate Valproate
	Carbamazepine, Phenobarbital, Phenytoin, Primidone
(ii) Myoclonic Seizure	
(iii) Tonic-clonic Seizure	

Mechanism of action of antiepileptic drugs[26]

Antiepileptic drugs may act mainly by one of three main mechanisms

1. Blockade of Na⁺ channel and prolongation of their inactive state and delaying their recovery Example- Phenytoin, carbamazepine, Lamotrigine
2. Blockade of low threshold Ca⁺⁺ current in thiamine neuron Absence seizure
3. Enhancing GABA mediated Inhibition
 - (I) By acting on GABA receptor- Benzodioropine
 - (ii) By inhibiting GABA Metabolism- Valproic acid
 - (iii) By blocking excitatory glutamate receptor

Pregnancy With Epilepsy

When pregnant, a lady with epilepsy who is being treated with AEDs ought to be trailed by an obstetrician who is alright with her ailment and treatment. A high-hazard obstetrician or maternal-fetal medication master is ideal, although this may not generally be conceivable should check circa 5-6 week 10 week and then once per trimester should be check after any seizure during pregnancy may present problems for both the mother and fetus Screening for malformation should be offered

- > AFT(either material serum or amniotic fluid at 14-16 week
- > Ultrasonography at 18-20 week
- > Drug level and appropriate dosage adjustment
- > patients should be advised for taking folic acid supplement
- > Vitamin K supplementation (10-20mg/D) is recommended for women on AED in the final month of pregnancy to reduce the risk of postnatal bleeding in a child[27].

Breast feeding

Breastfeeding an infant has numerous points of interest, including the maternal exchange of antibodies, just as advancing holding among mother and youngster. Likewise, for some, ladies breastfeeding is a significant piece of the experience of motherhood, In low concentration and breastfeeding American Academy of Pediatrics considers with epilepsy taking AEDs are safe except for the barbiturates and ethosuximide. Problems of neonatal sedation may occur with benzodiazepine and barbiturate and ethosuximide is distributed in significant amounts into breast milk and therefore breastfeeding should be avoided.[28-29].

EPILEPSY IN CHILDREN[30-31]

Fits in kids are treated as in grown-ups, yet youngsters may react diversely and because bad-tempered for example with sodium valproate or phenobarbitone. If febrile seizures have happened a medication utilized for significant epilepsy might be given constantly (for example phenobarbitone 3-4 mg/kg/day) until the youngster is 5 years of age. However, drawn-out medication treatment for example with phenytoin or phenobarbitone may meddle with the intellectual turn of events, the medication is pulled back

IV. CONCLUSION

A gigantic measure of research is centered around the conclusion, avoidance, treatment, and fix of epilepsy. Avoidance of injury is in every case superior to treatment. Accordingly, the determination of an anticonvulsant specialist depends basically on its viability for explicit kinds of seizures and epilepsy. Although seizure control is commonly accepted in many patients, a critical extent of patients with epilepsy experience the ill effects of obstinate or drug-resistant epilepsy, notwithstanding early treatment and an ideal day by day measurement of a sufficient anticonvulsant agents.

REFERENCES

1. P.A. Dekker, M.D. Epilepsy – A Manual to Medical & Clinical Officers in Africa. Revised 2nd edition. World Health Organization, Geneva 2002.
2. Technology Appraisal 76. Newer drugs for epilepsy in adults. ISBN: 1-84257-530-9. National Institute of Clinical Excellence (NICE), UK. March 2004.
3. Browne T.R. & Holmes G.L., Philadelphia, PA, Lippincott Williams & Wilkins. Handbook of Epilepsy. 2000; 2nd Ed. 42-55, 91-94.
4. Commission on Classification and Terminology of the International League against Epilepsy. Proposal for revised clinical and electroencephalographic classification of epileptic seizures. *Epilepsia*. 1981; 22:489– 501.
5. Engel Jr. Classifications of the International League against Epilepsy: time for reappraisal. *Epilepsia*. 1998; 39:1014–1017.
6. Manford, M., Hart, Y.M., Sander, J.W., Shorvon, S.D., 1992. The National General Practice Study of Epilepsy. The Syndromic Classification of the International League against Epilepsy applied to epilepsy in a general population. *Archives of Neurology*. 49:801–808.
7. K.P. Sampath Kumar et al. Recent Advances in Epilepsy drug therapy and management disease. *International Journal of PharmTech Research*. 2010, 2(1).
8. James W Y Chen, Claude G Wasterlain. Status Epilepticus: pathophysiology & management in adults. *The Lancet Neurology*. 2006; 5: 246–56.



9. Robert S. Fisher, M.D., Ph.D., Maslah Saul MD Professor of Neurology, What Causes Epilepsy? Stanford, Editor-in-Chief, epilepsy.com
10. Hirtz D, Thurman DJ, Gwinn-Hardy K, Mohamed M, Chaudhury AR, Zalutsky R. How common are the 'Common' neurologic disorders? Neurology 2007;68 (5):326-37.
11. Gidal BE, Garnett WR. Epilepsy. In: Dipiro J T, Talbert RL, Yee GC, Matzke GR, Wells BG, Posey LM, eds. Pharmacotherapy: a pathophysiologic approach. USA: McGraw-Hill Companies Inc, 6th ed. 2005:1023-46.
12. Hanna NJ, Black M, Sander J W et al. The national sentinel audit of epilepsy related death. The Stationary Office, London. 2002:1- 135
13. Mary Anne Koda-Kimble, Lloyd Yee Young, Brian K. Alldredge, Robin L. Corelli, B. Joseph Guglielmo, Wayne A. Kradjan, Bradely B. Williams, Applied therapeutics a clinical use of drugs 9th edition
14. Anthony Murro, M.D, Department of Neurology, Medical College of Georgia. Seizure Disorders. Available at URL: <http://www.georgiahealth.edu>
15. Causes of Epilepsy. Maslah Saul MD Professor, Stanford 2010. Available at URL: http://www.epilepsy.com/epilepsy/causes_s_eizures
16. HIRAK KUMAR MUKHOPADHYAY, CHANDI CHARAN KANDAR, SANJAY KUMAR DAS, LAKSHMIKANTA GHOSH, BIJAN KUMAR GUPTA, Epilepsy and its Management: A Review, Journal of PharmaSci Tech 2012; 1(2):2026.
17. Russell J Greene Norman D Harris, Pathology and therapeutics for pharmacy, third edition.
18. HIRAK KUMAR MUKHOPADHYAY, CHANDI CHARAN KANDAR, SANJAY KUMAR DAS, LAKSHMIKANTA GHOSH, BIJAN KUMAR GUPTA, Epilepsy and its Management: A Review, Journal of PharmaSci Tech 2012; 1(2):2026.
19. NG, Koehler AN, McGhee B, et al. The ketogenic diet in refractory epilepsy: the experience of Children's Hospital of Pittsburgh. Clinical Pediatrics (Phila) 2000; 39:153-159.
20. Marla J. Friedman, DO. Ghazala Q. Sharieff, MD. Seizures in Epilepsy. Paediatric Clinics of North America. 53. 2006; 257 – 277
21. Sudha K. Kessler et al. Dietary therapies for epilepsy: Future research. Epilepsy & behaviour 2011; 22:17-22
22. Sinha, S.R. and E.H. Kossoff (2005). The ketogenic diet. Neurologist, 11(3):161-70.
23. Freeman JM, Kelly MT, Freeman JB. The ketogenic diet: a treatment for epilepsy. 3rd edn. New York: Demos, 2000.
24. La Vega-Talbott M, Solomon GE, Mast J, Green NS, Hosain SA. Ketogenic diet in paediatric epilepsy patients with gastrostomy feeding (abstract). Annals of Neurology. 2001; 50 (suppl 1): S102.
25. [No authors listed] (1998) Practice parameter: management issues for women with epilepsy (summary statement). Report of the Quality Standards Subcommittee of the American Academy of Neurology. Epilepsia 39: 1226-1231
26. Schlanger S, Shinitzky M, Yam D. Diet enriched with omega-3 fatty acids alleviates convulsion symptoms in epilepsy patients. Epilepsia 2002; 43:103-104.
27. Pennell PB et al. (2004) The impact of pregnancy and childbirth on the metabolism of lamotrigine. Neurology 62: 292-295
28. [No authors listed] (1998) Practice parameter: management issues for women with epilepsy (summary statement). Report of the Quality Standards Subcommittee of the American Academy of Neurology. Epilepsia 39: 1226-1231
29. Pennell PB (2003) Antiepileptic drug pharmacokinetics during pregnancy and lactation. Neurology 61 (Suppl 2): S35-S42
30. Bennett PN, Brown MJ. Clinical pharmacology. New Delhi: Elsevier a division of Read Elsevier India (P) Ltd, 9th ed. 2006:413-22
31. NG, Koehler AN, McGhee B, et al. The ketogenic diet in refractory epilepsy: the experience of Children's Hospital of Pittsburgh. Clinical Pediatrics (Phila) 2000; 39:153-159.



**Impact Factor:
7.569**



INTERNATIONAL JOURNAL OF INNOVATIVE RESEARCH

IN SCIENCE | ENGINEERING | TECHNOLOGY

 **9940 572 462**  **6381 907 438**  **ijirset@gmail.com**



www.ijirset.com

Scan to save the contact details