



IJIRSET

International Journal of Innovative Research in
SCIENCE | ENGINEERING | TECHNOLOGY

e-ISSN: 2319-8753 | p-ISSN: 2320-6710



INTERNATIONAL JOURNAL OF INNOVATIVE RESEARCH

IN SCIENCE | ENGINEERING | TECHNOLOGY

Volume 10, Issue 5, May 2021

ISSN INTERNATIONAL
STANDARD
SERIAL
NUMBER
INDIA

Impact Factor: 7.512



9940 572 462



6381 907 438



ijirset@gmail.com



www.ijirset.com

Complex Some of the Results of the Analysis in Higher Mathematics

Sulaymonov Ilyosxo'ja Abdirasulo'g'li

2nd year student of the National University of Uzbekistan named after MirzoUlubek

ANNOTATION: This article is about the application complex analysis in real analysis, which gives us the opportunity to calculate several important integrals in a simple and convenient way and get some additional results. It is also stated that we can get some results even from another important integral Eyler-Puaason integral.

KEYWORDS: application complex analysis, another important integral

$$F(t) = \int_a^t f(x)dx, \text{ integral, Eyler formulation,}$$

$$e^{i\varphi}, \int_0^\infty e^{-x^2} dx, \text{ Eyler-Puasson integral}$$

1. Specific integrals depending on the parameter $f(x)$ function $[a, +\infty)$ in the range ($a \in R$)

granted as voluntary $[a, t]$ in ($a \leq t < +\infty$) let it be integrative. We enter the following

$$F(t) = \int_a^t f(x)dx$$

It is known to us that $t \rightarrow +\infty$ da $F(t)$ this is the limit if the function has a limit $f(x)$ function $[a, +\infty)$ on the infinite interval, the singular is called an integral and

$$\int_a^{+\infty} f(x)dx$$

as defined.

2. Eyler-Poassan integral

$$I = \int_0^\infty e^{-x^2} dx$$

Let's look at the function below $g(t) = (1+t)e^{-t}$

$t = \pm x^2$ we perform the replacement and get the following

$$t = \pm x^2 \rightarrow \begin{cases} 0 < (1-x^2)e^{x^2} < 1 \\ 0 < (1+x^2)e^{-x^2} < 1 \end{cases} \rightarrow \begin{cases} 0 < (1-x^2) < e^{-x^2} \\ 0 < e^{-x^2} < \frac{1}{1+x^2} \end{cases}$$

from this we can have the following

$$1) (1-x^2)^n < e^{-nx^2}, \quad 2) e^{-nx^2} < \frac{1}{(1+x^2)^n}$$

We integrate 1-inequality (0,1) in the range, 2-inequality (0,∞) in the range

$$\begin{aligned} \int_0^1 (1-x^2)^n dx &< \int_0^1 e^{-nx^2} dx < \int_0^{+\infty} e^{-nx^2} dx < \int_0^{+\infty} \frac{dx}{(1+x^2)^n} \Rightarrow \int_0^1 (1-x^2)^n dx < \int_0^{+\infty} e^{-nx^2} dx < \\ &< \int_0^{+\infty} \frac{dx}{(1+x^2)^n} \\ \int_0^{+\infty} e^{-nx^2} dx &= \left[\begin{array}{l} p = \sqrt{nx} \\ \frac{dp}{\sqrt{n}} = dx \end{array} \right] = \frac{1}{\sqrt{n}} \int_0^{+\infty} e^{-p^2} dp = \frac{1}{\sqrt{n}} I \Rightarrow \int_0^1 (1-x^2)^n dx < \frac{1}{\sqrt{n}} I < \int_0^{+\infty} \frac{dx}{(1+x^2)^n} \end{aligned}$$

Now we calculate these two integers

$$\begin{aligned} \int_0^1 (1-x^2)^n dx &= \left[\begin{array}{l} x = \cos t \\ dx = -\sin t dt \\ x = 0 \rightarrow t = \frac{\pi}{2} \\ x = 1 \rightarrow t = 0 \end{array} \right] = - \int_{\frac{\pi}{2}}^0 \sin^{2n+1} t dt = \int_0^{\frac{\pi}{2}} \sin^{2n+1} t dt = \frac{(2n)!!}{(2n+1)!!} \\ \int_0^{+\infty} \frac{dx}{(1+x^2)^n} &= \left[\begin{array}{l} x = ctgt \\ dx = -\frac{1}{\sin^2 t} dt \\ x = 0 \rightarrow t = \frac{\pi}{2} \\ x = +\infty \rightarrow t = 0 \end{array} \right] = - \int_{\frac{\pi}{2}}^0 \sin^{2n-2} t dt = \int_0^{\frac{\pi}{2}} \sin^{2n-2} t dt = \frac{(2n-3)!! \pi}{(2n-2)!! 2} \\ \frac{(2n)!!}{(2n-1)!!} &< \frac{I}{\sqrt{n}} < \frac{(2n-3)!! \pi}{(2n-2)!! 2} \rightarrow \sqrt{n} \frac{(2n)!!}{(2n+1)!!} < I < \frac{(2n-3)!! \pi}{(2n-2)!! 2} \sqrt{n} \\ \frac{n}{2n+1} \left[\frac{(2n)!!}{(2n-1)!!} \right]^2 \frac{1}{2n+1} &< I^2 < \frac{n}{2n-1} \left[\frac{(2n-3)!!}{(2n-2)!!} \right]^2 (2n-1) \frac{\pi^2}{4} \end{aligned}$$

From this, according to the vallis formula, we get the following

$$\lim_{n \rightarrow \infty} \frac{n}{2n+1} \left[\frac{(2n)!!}{(2n-1)!!} \right]^2 \frac{1}{2n+1} = \frac{\pi}{4}$$

$$\lim_{n \rightarrow \infty} \frac{n}{2n-1} \left[\frac{(2n-3)!!}{(2n-2)!!} \right]^2 (2n-1) \frac{\pi^2}{4} = \frac{\pi}{4}$$

All in all

$$I^2 = \frac{\pi}{4} \Rightarrow I = \frac{\sqrt{\pi}}{2}$$

A little $z \in C$ Complex number this $z = r(\cos \varphi + i \sin \varphi)$ let it be given in appearance. Complex is one of the following important formulas known from the course of analysis $e^{i\varphi} = \cos \varphi + i \sin \varphi$

Eyler is the formula. Now we calculate these integers with the help of the properties Complex numbers and the Eyler-Puasson integral the following multiple integers without the presence of the following integers[2].

1-example. Calculate the following specific integrals

$$I_1 = \int_0^\infty e^{-ax} \cos bx dx, \quad I_2 = \int_0^\infty e^{-ax} \sin bx dx, \quad (a > 0)$$

Solution We write the above two integers as follows

$$I_1 + iI_2 = \int_0^A e^{-ax} \cos bx dx + i \int_0^A e^{-ax} \sin bx dx \quad (1)$$

We look at the integral below.

$$\int_0^A e^{-ax} e^{ibx} dx = \int_0^A e^{(-a+ib)x} dx = \frac{e^{(-a+ib)x}}{-a+ib} \Big|_0^A = -\frac{a+ib}{a^2+b^2} e^{(-a+ib)x} \Big|_0^A = -\frac{a+ib}{a^2+b^2} e^{-ax} e^{ibx} \Big|_0^A$$

Now according to Eyler's formulation $e^{ibx} = \cos(bx) + i \sin(bx)$ we can write the above example as follows

$$\begin{aligned} -\frac{(a+ib)(\cos(bx) + i \sin(bx))}{a^2+b^2} e^{-ax} &= \frac{b \sin(bx) - a \cos(bx)}{a^2+b^2} e^{-ax} + \\ &+ i \frac{-a \sin(bx) - b \cos(bx)}{a^2+b^2} e^{-ax} \end{aligned} \quad (2)$$

As a result of the equation of (1) and (2) expressions according to the above theorem, we get the following

$$I_1 = \int_0^A e^{-ax} \cos bx dx = \frac{b \sin(bx) - a \cos(bx)}{a^2+b^2} e^{-x^2} \Big|_0^A$$

$$I_2 = \int_0^A e^{-ax} \sin bx dx = \frac{-a \sin(bx) - b \cos(bx)}{a^2+b^2} e^{-x^2} \Big|_0^A$$

I_1 and I_2 value of $A \rightarrow \infty$ although will be as follows:

$$I_1 = \frac{a}{a^2+b^2}, \quad I_2 = \frac{b}{a^2+b^2}$$

2-example. Calculate the following specific integrals:

$$I_1 = \int_0^\infty \cos(x^2) dx, \quad I_2 = \int_0^\infty \sin(x^2) dx$$

Taking into account the presence of these integers, we are dealing with their calculation

Solution

We write the given integers as follows

$$I_1 - iI_2 = \int_0^\infty \cos(x^2) dx - i \int_0^\infty \sin(x^2) dx = \int_0^\infty (\cos(-x^2) + i\sin(-x^2)) dx = \int_0^\infty e^{-ix^2} dx$$

Now this $\int_0^\infty e^{-ix^2} dx$ we calculate the integral

$$\int_0^\infty e^{-ix^2} dx = \begin{bmatrix} \sqrt{ix} = t \\ ix^2 = t^2 \\ dx = \frac{1}{\sqrt{i}} dt \end{bmatrix} = \frac{1}{\sqrt{i}} \int_0^\infty e^{-t^2} dt = \frac{1}{\sqrt{i}} \frac{\sqrt{\pi}}{2}$$

$\frac{1}{\sqrt{i}}$ let's calculate. $\frac{1}{\sqrt{i}} = -i^{\frac{3}{2}} = -\left(\cos\left(\frac{\pi}{2}\right) + i\sin\left(\frac{\pi}{2}\right)\right)^{\frac{3}{2}}$ According to the Muavr formulation
 $-\left(\cos\left(\frac{\pi}{2}\right) + i\sin\left(\frac{\pi}{2}\right)\right)^{\frac{3}{2}} = -\cos\left(\frac{3\pi}{2}\right) - i\sin\left(\frac{3\pi}{2}\right) = \frac{\sqrt{2}}{2} - i\frac{\sqrt{2}}{2}$

$$\text{So } I_1 - iI_2 = \frac{\sqrt{2}}{2} \frac{\sqrt{\pi}}{2} - i \frac{\sqrt{2}}{2} \frac{\sqrt{\pi}}{2} = \frac{1}{2} \sqrt{\frac{\pi}{2}} - i \frac{1}{2} \sqrt{\frac{\pi}{2}}$$

Complex according to the theorem on the equality of numbers

$$I_1 = \int_0^\infty \cos(x^2) dx = \frac{1}{2} \sqrt{\frac{\pi}{2}}$$

$$I_2 = \int_0^\infty \sin(x^2) dx = \frac{1}{2} \sqrt{\frac{\pi}{2}}$$

REFERENCES

1. Godberganov G., Verisov A., Mansurov H. Eksexanalysis. Tashkent, University, 1998
2. Fichtenholz G. Course of differential and integral calculus. Vol. II. Moscow, ФизFizmat-lit», 2001
3. Demidovich B. Collection of problems and exercises in mathematical analysis. Moscow, Hayka Nauka», 1990



INTERNATIONAL JOURNAL OF INNOVATIVE RESEARCH

IN SCIENCE | ENGINEERING | TECHNOLOGY

9940 572 462 6381 907 438 ijirset@gmail.com

