



IJIRSET

International Journal of Innovative Research in
SCIENCE | ENGINEERING | TECHNOLOGY

e-ISSN: 2319-8753 | p-ISSN: 2347-6710



INTERNATIONAL JOURNAL OF INNOVATIVE RESEARCH

IN SCIENCE | ENGINEERING | TECHNOLOGY

Volume 10, Issue 11, November 2021

ISSN INTERNATIONAL
STANDARD
SERIAL
NUMBER
INDIA

Impact Factor: 7.569



9940 572 462



6381 907 438



ijirset@gmail.com



www.ijirset.com

Modifying the Gut Microbiome to Make Cancer Immunotherapy Effective

Dr.Nida Jamil Khan

Assistant Professor, Department of Biosciences, Jamia Millia Islamia University, Delhi, India

Abstract:

Background: A complex and diverse ecosystem residing at all epithelial barriers of host body constitutes the mammalian microbiome. When the gut microbiome is disrupted it is associated with a variety of inflammatory disorders and malignancies.

Microbiota-host interactions may affect the host response to immunotherapies by modulating innate and adaptive immunity.

Conclusion: Our goal should be development of precision immunotherapies by identification of specific causative bacteria and modulation of cancer progression.

Keywords: microbiome,immunotherapy,dysbiosis

I. BACKGROUND

A healthy individual possesses a good microbial ecosystem. In humans the microbiome comprise bacteria, fungi, archaea, protozoa and virus that reside on body's epithelial barrier(1). The gut microbiome contains approximately 3×10^{13} bacteria, most of which are symbiotic with the host(2,3). For maintaining the intestinal epithelial homeostasis the gut microbiome ,immune cells and mucosal barrier act together. The gut microbiome play a key regulatory role. They regulate barrier function, maintain mucosal immune homeostasis and host-microbial symbiosis, prevent pathogen infection, control overgrowth of pathological organisms, and regulate metabolism, including vitamin synthesis and metabolism of indigestible dietary fiber (4-8).

Among the human microbiome certain bacteria have been known to play an important role in the development of cancer(9). These days immune checkpoint inhibitors(ICIs) are used in cancer immunotherapy.(10) However in certain patients resistance to immunotherapy is developed. Recent studies have shown the role of gut microbiome in improving the efficacy of ICIs against cancer(11-15). In this article we will be analyzing the role of gut microbiome in increasing the efficiency of cancer immunotherapy and the challenges faced by them with possible solutions.

II. ROLE OF GUT MICROBIOME IN CANCER IMMUNOTHERAPY

Immune checkpoint inhibitors play a major role in cancer immunotherapy. They inhibit the interactions between immune checkpoint molecules and inhibitory T cell receptors and stimulate T cell-mediated antitumor immunity. A wide range of metastatic malignancies have been effectively treated using PD-1 blockade as immunotherapy. Varied responses are shown by patients and some have become resistant to it(16-18).

Antibiotic-related dysbiosis was associated with a lower response to PD-1 blockade which means that disruption of microbial networks and loss of specific bacterial species can affect the efficacy of immunotherapy(19). One of the important bacterial species found in fecal microbiota of the patients was *A. muciniphila*. Oral supplementation with *A. muciniphila* or fecal microbiota transplantation (FMT) from responders restored the efficacy of PD-1 blockade in murine models. Cytokines induced by microbiota-host interactions increase the efficacy of PD-1 blockade. But this is not applicable universally in all human races and the response vary from individual to individual(20).

III. CONCLUSION

Thus our goal should be the development of precision immunotherapies. The identification of specific causative bacteria may allow modulation of cancer progression, raising the possibility of precision immunotherapies.

REFERENCES

- (1) Costello EK, Stagaman K, Dethlefsen L, Bohannan BJ, Relman DA. The application of ecological theory toward an understanding of the human microbiome. *Science* 336 (2012) 1255–1262.
- (2) Cho I, Blaser MJ. The human microbiome: at the interface of health and disease. *Nat. Rev. Genet.* 13 (2012) 260–270.
- (3) Sender R, Fuchs S, Milo R. Revised estimates for the number of human and bacteria cells in the body. *PLoS Biol.* 14 (2016) e1002533.
- (4) De Santis S, Cavalcanti E, Mastronardi M, Jirillo E, Chieppa M. Nutritional keys for intestinal barrier modulation. *Front. Immunol.* 6 (2015) 612.
- (5) Vaishnavi S, Behrendt CL, Ismail AS, Eckmann L, Hooper LV. Paneth cells directly sense gut commensals and maintain homeostasis at the intestinal host-microbial interface. *Proc. Natl. Acad. Sci. U.S.A.* 105 (2008) 20858–20863.
- (6) Peterson LW, Artis D. Intestinal epithelial cells: regulators of barrier function and immune homeostasis. *Nat. Rev. Immunol.* 14 (2014) 141–153.
- (7) Chung WS, Walker AW, Louis P, Parkhill J, Vermeiren J, Bosscher D, Duncan SH, Flint HJ. Modulation of the human gut microbiota by dietary fibers occurs at the species level. *BMC Biol.* 14 (2016) 3.
- (8) Sommer F, Backhed F. The gut microbiota engages different signaling pathways to induce Duox2 expression in the ileum and colon epithelium. *Mucosal Immunol.* 8(2015) 372–379.
- (9) Zhang H, Sun L. When human cells meet bacteria: precision medicine for cancers using the microbiota. *Am J Cancer Res.* 2018;8(7):1157–75.
- (10) Weintraub Li, Yu Deng, Qian Chu, Peng Zhang. Gut microbiome and cancer immunotherapy. *Cancer Letters* 447 (2019) 41–47.
- (11) Chaput N, Lepage P, Coutzac C, Soularue E, Le Roux K, Monot C, Boselli L, Routier E, Cassard L, Collins M, Vaysse T, Marthey L, Eggermont A, Asvatourian V, Lanoy E, Mateus C, Robert C, Carbonnel F. Baseline gut microbiota predicts clinical response and colitis in metastatic melanoma patients treated with ipilimumab. *Ann. Oncol.* 28 (2017) 1368–1379.
- (12) Frankel AE, Coughlin LA, Kim J, Froehlich TW, Xie Y, Frenkel EP, Koh AY. Metagenomic shotgun sequencing and unbiased metabolomic profiling identify specific human gut microbiota and metabolites associated with immune checkpoint therapy efficacy in melanoma patients. *Neoplasia* 19 (2017) 848–855.
- (13) Matson V, Fessler J, Bao R, Chongsuwat T, Zha Y, Alegre ML, Luke JJ, Gajewski TF. The commensal microbiome is associated with anti-PD-1 efficacy in metastatic melanoma patients. *Science* 359 (2018) 104–108.
- (14) Gopalakrishnan V, Spencer CN, Nezi L, Reuben A, Andrews MC, Karpinets TV, Prieto PA, Vicente D, Hoffman K, Wei SC, Cogdill AP, Zhao L, Hudgens CW, Hutchinson DS, Manzo T, Petaccia de Macedo M, Cotechini T, Kumar T, Chen WS, Reddy SM, Szczepaniak Sloane R, Galloway-Pena J, Jiang H, Chen PL, Shpall EJ, Rezvani K, Alousi AM, Chemaly RF, Shelburne S, Vence LM, Okhuysen PC, Jensen VB, Swennes AG, McAllister F, Marcelo Riquelme Sanchez E, Zhang Y, Le Chatelier E, Zitvogel L, Pons N, Austin-Breneman JL, Haydu LE, Burton EM, Gardner JM, Sirmans E, Hu J, Lazar AJ, Tsujikawa T, Diab A, Tawbi H, Glitzka IC, Hwu WJ, Patel SP, Woodman SE, Amaria RN, Davies MA, Gershenson JE, Hwu P, Lee JE, Zhang J, Coussens LM, Cooper ZA, Futreal PA, Daniel CR, Ajami NJ, Petrosino JF, Tetzlaff MT, Sharma P, Allison JP, Jenq RR, Wargo JA. Gut microbiome modulates response to anti-PD-1 immunotherapy in melanoma patients. *Science* 359 (2018) 97–103.
- (15) Routy B, Le Chatelier E, Derosa L, Duong CPM, Alou MT, Daillyere R, Fluckiger A, Messaoudene M, Rauber C, Roberti MP, Fidelle M, Flament C, Poirier-Colame V, Opolon P, Klein C, Iribarren K, Mondragon L, Jacquemet N, Qu B, Ferrere G, Clemenson C, Mezquita L, Masip JR, Naltet C, Brosseau S, Kaderbhai C, Richard C, Rizvi H, Levenez F, Galleron N, Quinquis B, Pons N, Ryffel B, Minard-Colin V, Gonin P, Soria JC, Deutsch E, Loriot Y, Ghiringhelli F, Zalcman G, Goldwasser F, Escudier B, Hellmann MD, Eggermont A, Raoult D, Albiges L, Kroemer G, Zitvogel L. Gut microbiome influences efficacy of PD-1-based immunotherapy against epithelial tumors. *Science* 359 (2018) 91–97.
- (16) Borghaei H, Paz-Ares L, Horn L, et al. Nivolumab versus Docetaxel in Advanced Non-Squamous Non-Small-Cell Lung Cancer. *N Engl J Med* 2015;373:1627–39.
- (17) Motzer RJ, Escudier B, McDermott DF, et al. Nivolumab versus Everolimus in Advanced Renal-Cell Carcinoma. *N Engl J Med* 2015;373:1803–13.
- (18) Topalian SL, Hodi FS, Brahmer JR, et al. Safety, activity, and immune correlates of anti-PD-1 antibody in cancer. *N Engl J Med* 2012;366:2443–54.
- (19) Routy B, Le Chatelier E, Derosa L, et al. Gut microbiome influences efficacy of PD-1-based immunotherapy against epithelial tumors. *Science* 2018;359:91–7.
- (20) Shigematsu Y, Inamura K. Gut microbiome: a key player in cancer immunotherapy. *HepatoBiliary Surg Nutr* 2018;7(6):479–480



Impact Factor:
7.569



INTERNATIONAL JOURNAL OF INNOVATIVE RESEARCH

IN SCIENCE | ENGINEERING | TECHNOLOGY

9940 572 462 **6381 907 438** **ijirset@gmail.com**

www.ijirset.com



Scan to save the contact details