



e-ISSN: 2319-8753 | p-ISSN: 2347-6710

IJIRSET

International Journal of Innovative Research in
SCIENCE | ENGINEERING | TECHNOLOGY



INTERNATIONAL JOURNAL OF INNOVATIVE RESEARCH

IN SCIENCE | ENGINEERING | TECHNOLOGY

Volume 11, Issue 6, June 2022

ISSN INTERNATIONAL
STANDARD
SERIAL
NUMBER
INDIA

Impact Factor: 8.118



9940 572 462



6381 907 438



ijirset@gmail.com



www.ijirset.com

Usage of Computational Fluid Dynamics (CFD) to Test the Efficiency of Cancer Treatment Methodologies

Swayambhik Mukherjee¹

Post-Graduate Department of Biotechnology, St. Xavier's College (Autonomous), Kolkata, India¹

ABSTRACT: Computational fluid dynamics (CFD) is a field of fluid mechanics that employs numerical evaluation and interpretation of statistical structures to examine and determine solutions to issues involving fluid flow. Computer systems are programmed to make calculations suited for simulating the free flow of fluids, as well as the interaction of fluids with areas restricted by certain boundary situations. This review is on the usage of CFD to test the efficiency and accuracy of certain cancer treatment procedures. The aim of this paper is to propose the usage of CFD as an alternative for physical experimentation as it provides cost-effective and precise results.

KEYWORDS: Computational Fluid Dynamics (CFD) , Pharmacokinetics , Cancer Treatment ;

I. INTRODUCTION

Cancer is an abnormal condition in which certain cells of the body grow uncontrollably and rapidly spread to other parts of the body. Cells present in any part of the human body can turn out to become cancerous. Normally, human cells grow and multiply (by a process called cell division) to form new cells as the body needs them. When the cells get old or damaged, they die, and new cells take their place. These cells can form tumors, which are tissues. They may or may not be cancerous (benign). Cancer cells spread, or invade, to nearby tissues and can travel to distant parts of the body to form new tumors (a process called metastasis). Contact inhibition is the process of stopping cell growth when cells come into contact. As a result, normal cells stop multiplying when they form a monolayer in a culture dish. Contact inhibition is a powerful way to prevent cancer, this property is absent in cancer cells. Cancer tumors are known to be 'malignant'. Many cancers form solid tumors, but blood cancers, such as leukemia, usually do not. Benign tumors do not spread, or invade, nearby tissues, also after being removed, the benign tumors usually do not grow back unlike the malignant ones which have a possibility of reforming. However, Benign tumors can be quite large at times. Some can cause serious symptoms or even life-threatening symptoms, such as damage to the brain.

Cancer is the leading cause of death worldwide, accounting for about 9.9 million deaths by 2020, or about one in six deaths. Approximately, 33% of the deaths from cancer are due to tobacco use, high body mass index (BMI calculated as the ratio between weight in kg to the square of height in m), alcohol consumption, low intake of fruits and vegetables, and lack of exercise. Cancer-causing diseases, such as human papillomavirus (HPV) and hepatitis, are the cause of almost 30% of cancer cases in low and lower-middle income nations. Most cancers can be cured if they are diagnosed early and treated successfully.

Globally, 18,094,716 cancer cases were diagnosed by 2020. The Age-Standardized Rate (ASR) for all cancers (except melanoma-free skin cancer) in men and women combined was 190 out of 100,000 by 2020. The rate was higher for men (206.99) per 100,000 than women (178.1 per 100,000).

Cases of Incidence	Global Data (Number)	ASR / 100,000
	18,094,716	190.0

Table 1 : Cases of cancer occurrence globally.

Highest Incidence of Cancer					
Men		Women		Both sexes	
Country, Number	ASR / 100,000	Country, Number	ASR / 100,000	Country, Number	ASR / 100,000
Hungary, 31,496	371. 0	Denmark, 19,267	328. 3	Denmark, 39,996	334. 9

Table 2 : Countries that report highest occurrence of cancer along with the number of cases observed in men ,women and both.

Cases of Mortality	Global Data (Number)	ASR / 100,000
	9,894,402	100. 1

Table 3: Cases of mortality globally.

Mortality due to Cancer					
Men		Women		Both sexes	
Country, Number	ASR / 100,000	Country, Number	ASR / 100,000	Country, Number	ASR / 100,000
Mongolia, 2,525	224. 3	Zimbabwe, 6,628	142. 9	Mongolia, 4,465	175. 9

Table 4 : Mortality in terms of number in different countries , in men , women and both.

II. METHODS AND EXPERIMENTAL RESULTS

1. CFD study of intra-arterial chemotherapy for oral cancer

Intra-arterial chemotherapy (IAC), also termed as as super-selective ophthalmic artery chemotherapy or chemosurgery, is now considered as one of the most important therapies for intraocular retinoblastoma across the world. Retinoblastoma (RB) is one of the most commonly treated childhood diseases. Targeted RB treatment with direct delivery of chemotherapeutic agents to the coronary artery (Ophthalmic Artery - OA) has dramatically altered the way we treat this fatal, but curable eye cancer. Intra-arterial chemotherapy (IAC) through the ophthalmic artery has the advantage of high concentration of chemotherapy reaching the tumor, with systemic side effects being overlooked compared to systemic intravenous chemotherapy (IVC). Over the past decade, we have seen growing IAC indicators of tumor control and eye rescue in an advanced and resistant retinoblastoma. Intra-arterial chemotherapy (IAC) is effective in controlling retinoblastoma as a primary and secondary treatment. OAC treatment uses less chemotherapy than when chemotherapy is administered intravenously. It avoids complications and adverse events associated with

toxicity from the systemic, rather than local chemotherapy. The role of IAC in underdeveloped tumors is yet to be established. IAC and IVC continue to be important treatment options for eye rescue with advanced RB. The IAC offers the benefit of reducing the risks of chemotherapy-related complications such as neutropenia, anemia, and secondary neoplasms. Problems in the IAC may be due to the injection strategy itself; and from the use of high doses of chemotherapeutic agents that can lead to ocular and systemic toxicity. Intra-arterial chemotherapy (IAC) has been used to treat malignant neoplasms locally in patients with head and neck cancer for over 50 years. IAC had advantages in the eyes with advanced disease, including groups D, IV, and V, but not in groups A-C, E, and I-III eyes. The overall success rate is higher for IAC than IVC.

We numerically investigated the flow distribution of anticancer agents into the branches of the external carotid artery in IAC models. The CFD simulation results of IAC showed that a curved catheter tip may be placed both below and toward the target artery to increase the distribution rate of anticancer agent into the tumor-feeding artery. Although super-selective IAC can reliably supply anticancer agents to the target artery, high wall shear stress at the target artery can occur, depending on vessel geometry of the patient, and this may cause serious complications during treatment. In future study, analysis of the fluid structure interaction is necessary to account for the elasticity of the vessel walls and catheter movement in many patient models.




Figure 1 : Retrograde Intra-Arterial Chemotherapy for oral cancer. (a) Super-selective intra-arterial chemotherapy, (b) Facial Artery and Lingual Artery diverge from the common trunk, and (c) Traditional intra-arterial chemotherapy. (Adapted from an Open Access Article and used under [Creative Common CC BY license](https://creativecommons.org/licenses/by/4.0/) with proper citation). [21]

2. Multiscale CFD model for Personalized Liver Cancer Radioembolization Dosimetry

The liver is a cone-shaped organ located in the upper right part of the human abdomen, below the diaphragm and above the abdomen. Various types of cancer can affect the liver. Hepatocellular carcinoma is the most widely seen type of liver cancer, it initiates in the hepatocyte, which is the main type of liver cell (occurs in liver that has been damaged by birth defects, alcohol abuse, or chronic illnesses such as hepatitis B and C, hemochromatosis and cirrhosis of the liver). Intrahepatic cholangiocarcinoma and hepatoblastoma which constitute the other types of liver cancer are much less seen.

The liver can be affected by primary liver cancer (which initiates from the liver), or cancer from other parts of the body that spread to the liver. Most liver cancers are secondary or metastatic, meaning they started somewhere else in the body (secondary liver cancer).

The most common benign tumors in the liver include Hemangioma, Liver adenoma, Focal nodular hyperplasia, Cysts, Lipoma, Fibroma, Leiomyoma. None of these tumors are treated as liver cancer. They may need to be surgically removed if they cause pain or bleeding.

Liver cancer includes:

- Hepatocellular carcinoma (HCC)
- Cholangiocarcinoma : These are actually gall bladder cancers.

Secondary liver cancer (metastatic liver cancer): Most of the time, the cancer found in the liver did not start there, but had spread (metastasized) from another part of the body, such as the pancreas, stomach, breast, lung or colon. Because this cancer has spread from its (primary) location, it is called secondary liver cancer.

The noninvasive flow in the patient's hepatic artery can be statistically evaluated using CFD, as part of a new dosimetry method for radioembolization of liver cancer. The number of microsphere delivered to different liver segments was calculated at different injection sites based on clinical data. The results showed a varied distribution of 90Y microspheres in liver segments, indicating the importance of CFD patient-specific mimicry for personal dosimetry. Compared with current dosimetry methods, personal dosimetry has the potential to increase the volume brought while reducing the number of microspheres embedded in other healthy tissues.




Figure 2 : Volume of liver subjected to radiation determined by arteries which received radioactive microsphere injections. (Licenses and permissions reserved, Used with Copyright permission : [View permission](#)). [22]

3. Use of 3D CFD model in intraperitoneal chemotherapy

The three-dimensional CFD model for mass drug delivery to tumor nodules during Intraperitoneal (IP) chemotherapy to incorporate actual tumor geometry and flexible vascular properties. Dynamic Contrast-Enhanced Magnetic Resonance Imaging (DCE-MRI) imaging has made it possible to differentiate between tumor tissues, healthy surrounding tissues and necrotic areas based on differential vascular structure. We found that the effect of interstitial pressure profiles within tumors was highly dependent on abnormal geometry and location. The depth of the cisplatin penetration of a particular tumor ranged from 0.32 mm to 0.50 mm. It can be stated that the positive relationship between tumor size and the Interstitial Fluid Pressure (IFP) is no longer maintained where there are areas with different vascular conditions, at a same time, a positive relationship has been noted between the percentage of active tumor and the main IFP.




Figure 3 : Illustration of the work plan described in this task. (a) Subperitoneal injection of SKOV3-Luc-IP1 cells is suspended from Matrigel in athymic mice, nude-foxn1nu in the growth phase. (b) In vivo in situ (IVIS) scan is performed 14 days after injection to assess tumor growth. (c) Scanning protocol as defined in the MRI protocol section which includes anatomical scans to detect geometric tumors, FLASH sequences for traditional tissue rest periods and DCE-MRI from which tumor tissue parameters were obtained. The red-circular anatomical MRI piece has the same function as the DCE-MRI plane. (d) Mice that approved for tumor growth received 30 min of intraperitoneal chemotherapy (IPC) with cisplatin. The treatment parameters (dose, duration) used in the test were used in setting the model as boundary conditions. (e) Shortly after the IPC, the Interstitial Fluid Pressure (IFP) was measured using a pressure tip catheter. The resulting pressures were used to confirm the pressure of the simulated tumor in the future. (f) Your calculations for both the formation of the IFP using the vascular parameters found in the DCE-MRI data and the mass transport of drugs were used in COMSOL and resolved using the same initial conditions and parameters as experimental setup. (Adapted from an Open Access Article and used under [Creative Common CC BY license](https://creativecommons.org/licenses/by/4.0/) with proper citation). [23]

4. Usage of CFD to judge the arrangement of artery with tumor vasculature

In a pancreatic tumor, there is a growth of blood vessels near the tumor, which allows for the transport of particles to the tumor vasculature. In case of our simulation, the geometry is made simpler by focusing on a main artery with pressure influenced flow, and another branching out to the zone of the tumor. The tumor vasculature can be simply thought of as stacked cylinders. This is illustrated in figure 5. Blood flows from P1 to P2 and possibly enters the tumor region at interstitial pressure P3, through a hole. The relation between the three stands as $P1 < P2 < P3$.




Figure 4 : Geometry of artery with tumor vasculature. [24]

5. CFD model of blood flow in solid tumors – Verification with test results :

When a multiphase CFD model combining basic relationships to measure hemodynamics in disturbed flow areas was used, simulation of the two-phase dilute flow simulation of solid human cells in a stable state fits well with test results in vortex formation, reuptake points and velocity profiles. In addition, the linear speed of the cross section in the middle of the vortex area showed the best correlation of the estimated speed.




Figure 5 : Simulation showing all the points under consideration. [24]

6. CFD Analysis of Topical Bronchial Tree Models for Lung Cancer Patients :

Lung surgery is the only possible treatment for lung cancer. The inevitable removal of active lung tissue and body weight requires a careful and systematic pre-surgery procedure for patients. In particular, the efficacy of post-lung residue surgery needs to be predicted. Clinically, these tests are algorithms based on pulmonary function tests (PFTs). However, these methods ignore the functioning of part of the local airway and provide inland exploration. CFD is designed to provide patient-specific, quantitative, and local information about energy flow and ventilation of the bronchial region. CFD is primarily used to detect flow fluctuations in 12 patients affected by lung cancer and to evaluate the effects of tumor mass on flow parameters and the distribution of lobar flow. Patient-specific airway models were reconstructed with Computed Tomography (CT) images, and the tumor mass was separated by hand. Lung

measurements and tumors are collected. The peripherality index is defined to describe the distance a tumor from the parenchyma. CFD simulations are performed on Fluent [®], and the results are analyzed in terms of flow and lobar volume flow rate (VFR).




Figure 6 : Distribution of pressure in the CFD bronchial tree model. (Licenses and permissions reserved, Used with Copyright permission : [View permission](#)). [25]

7. CFD modelling in the design and testing of an endovascular chemofilter device

Intra-Arterial Chemotherapy (IAC) is the preferred treatment for primary liver cancer, without its side effects. During the process, a blend of chemotherapeutic drugs, e. g. Doxorubicin, is injected into the veins that supply the tumor. A portion of Doxorubicin is absorbed by the tumor, but the remaining drug passes into the circulatory system, causing irreversible heart failure. The efficacy and safety of IAC can be enhanced by over-the-counter chemical analysis of catheter-based chemofilter, as suggested by a team of neurologists. CFD modeling is used to improve the hemodynamic and drug binding of the Chemofilter device.




Figure 7 : (a) Flow lines and shear pressure distribution of a 2x2 microcell matrix (b) Pressure and (c) Speed calculators are calculated to flow through two 2x2 matrix layers with 100 μm microcellites. (Adapted and used under [Creative Common CC BY license](#) with proper citation). [26]

8. Use of CFD model to understand Pharmacokinetics in solid tumors :

Of the three main types of cancer treatment — surgery, radiation, and chemotherapy — chemotherapy is one of the most powerful stages of new success. However, the biggest challenge of chemotherapy is to move the drug to the desired location at a high enough level to work. Computational fluid dynamics (CFD) simulations were used to test the behavior of several vectors of drug delivery in tumor capillaries — mainly moving bacteria, non-motile bacteria, and nanoparticles. Red blood cells, bacteria, and nanoparticles are placed in a stream using an immersed border method. The human capillary model was developed using a new method for the management of impaired red blood cells (RBCs). The model was verified with experimental data available from the literature. The stochastic model of bacterial motility was described based on tests of running and falling. Capillary models and bacteria are combined to mimic the intracapillary transport of bacteria. Non-motile bacteria, along with nanoparticles 0.2 μm , 0.3 μm , and 0.405 μm are also presented in the simulation in capillary flow in order to compare them with moving bacteria. Moving bacteria usually swim into a plasma layer near the capillary wall, whereas non-moving bacteria are often trapped in a bolus flow between RBCs. Nanoparticles are strongly influenced by Brownian movements and fluid fluctuations of small scale, so they are not inclined to a single flow area. Motile strains were found to have a long shelf life of 1 mm in length and a high radial velocity. This suggests that migrating bacteria can enter interstitium at a higher rate than nonmotile bacteria or nanoparticles with a diameter between 200 nm and 405 nm.




Figure 8 : Scheme of drug delivery to solid tumors. (Licenses and permissions reserved, Used with Copyright permission : [View permission](#)). [27]

9. Multiphysics CFD modeling for alpha-immunoconjugate delivery

Radiotherapy is a very widely used treatment for cancer. The research trend and success in radiotherapy focuses on high-energy radiation (Linear Energy Transfer -LET) radiation as alpha particles. Thus, the alpha-based method is an important means of the development of radiotherapy, and Targeted Alpha Therapy (TAT) is the most important program for the use of alpha particles over the past decade. This paper highlights research work done on TAT computer modeling at Canadian Nuclear Laboratories (CNL). This attempt demonstrates the feasibility of combining CFD mimicry and microdosimetric modeling to evaluate the effectiveness of TAT in a realistic and accurate way.




Figure 9 : Dose in the target per alpha particle (at 5.8 MeV). (Licenses and permissions reserved, Used with Copyright permission : [View permission](#)). [28]

10. CFD use to assess adhesion of cancer cells to white blood cells

Penn State researchers are learning how cancer cells attach to white blood cells, circulatory protectors, and how blood flow affects this adhesion. The study by Meghan Hoskins (a PhD student at Penn State) is sponsored by the National Cancer Institute and the PSU Applied Research Laboratory. Her goal is to accurately mimic previous experimental conditions of this condition in order to validate her model for further study of the metastasis process. Tumor cell adhesion can be affected by changes in the flow rate within the micro capillaries of living tumor cells, according to the Canadian Institute for Medical Research (CIMR) by Jennifer Hoskins. Hoskins attempts to demonstrate dynamic and biochemistry activity in the workplace during in vitro cell adhesion. She uses computational fluid dynamics (CFD) to create a 3D geometric grid for the test room. Researchers at the University of Bristol study how cells interact, collide, and coalesce, and measure the intensity of this activity. So far, Hoskins plans to model two sets of experiments. Cancer cell adhesion to white blood cells is affected by shear level and shear stress, or the energy produced by flow. In the second model, the room is closed, but researchers can watch as the cells interact with each other internally. It aims to understand why and how the migration of cancer cells is affected by the fluid dynamics of the system. If they can detect the presence of cancer-causing proteins in the bloodstream making white blood cells more efficient, perhaps medical researchers could devise a way to block the activity of those proteins. Hoskins hopes that her model will see continuous improvement, especially with regard to simulating adherence within the geometric structure of real blood vessels.




Figure 10 : Model for blood flow. (Licenses and permissions reserved, Used with Copyright permission : [View permission](#)). [29]

11. CFD used to create 3D model of bone metastasis and to check perfusion

Breast cancer usually spreads to the bone marrow. Bone metastatic cancer is incurable and causes widespread bone loss, leading to serious illness and death. Mechanical markers in the skeletal muscle are important environmental parameters that alter tumor formation, osteolysis, and tumor cell-bone signaling, but which mechanisms are most beneficial and the corresponding cell mechanisms are unknown. Interstitial fluid flow based on its well-known role in bone resorption and primary breast cancer was concentrated. A full-scale computer model, based on the microCT of the 3D model of bone metastasis used for filling to predict the internal machine space during in vitro testing was created. The perfusion used causes uniform dispersion, different liquid speeds and shear pressure on the wall throughout the inside of the scaffold. The liquid velocity distribution and shear pressure on the wall did not change between the model sub-domains of varying widths and spaces. Additionally, the magnitude of these stimuli lies within the range of anabolic mechanical signals in the skeletal structure, which ensures that our 3D model reflects previous in vivo studies using machine-based anabolic loading in the context of bone metastasis. Our results show that the local population of cells throughout the scaffold will experience the same mechanical environment.




Figure 11 : The flow rate in the scaffolds was very high in the high-contact regions, and the hydrostatic pressure on the wall varied by line. (A) On the inlet side, the incoming speed lines reveal the lateral flow to the scaffold area characterized by a greater connection. (B) On the outlet side of the scaffold, the exit lines showed the flow of the dispersed liquid evenly, although a small area of high velocity was noted, corresponding to the upper lateral flow region at the surface. (C) Frequency histograms have shown that the fluid flow velocity at the entry point of the scaffold is higher and more concentrated than the outside of the outlet. Note that histograms drop speeds in the upper extremities, 0 due to non-slip boundary conditions, although these values are included in all frequency calculations. (D) A contour map of the inner basement showed the pressure distributed by the line. (Licenses and permissions reserved, Used with Copyright permission : [View permission](#)). [30]

12. CFD modelling for a growing bronchial tumor

The methods of diagnosis and treatment of a lung tumor vary from place to place. Such methods include surgery, radiation, chemotherapy, etc. Knowing the location and size of the tumor will be very helpful in planning the treatment. In addition, accurate drug delivery to the infected area will enhance the treatment effect and reduce unwanted side effects in some regions. Currently, such information (tumor location and size) can only be obtained using CT, PET, or MRI. The development of the newly developed CFD-Fractal system is used to measure the presence and progression of lung diseases. This is done using the CFD-Fractal method in a lung model with a growing bronchial squamous tumor. Ventilated aerosol patterns will be detected using a body-based model, and will also be expressed using mathematical and fractal analysis. Many aspects of aerosol pattern will be examined in relation to tumor growth, including particle

transformation, spatial distribution potential, mono-fractal size, and multifractal spectrum. These factors can be used to identify the location of the lung disease, to measure the severity of the disease, and to later design a treatment plan with the intended drug delivery and appropriate dosage. The proposed approach will pave the way for the development of a therapeutic (therapeutic-diagnostic) system for the diagnosis and treatment of lung diseases.




Figure 12 : Pressure distribution in ten-generation bronchial tube model. (Licenses and permissions reserved, Used with Copyright permission : [View permission](#)). [31]

III. DISCUSSIONS

In our study, we have employed Computational Fluid Dynamics (CFD) based simulation models to gather experimental information driven results on certain methodologies related to cancer treatment, the usage of CFD has hugely reduced the hectic costs of carrying out physical experiments. We numerically investigated the flow distribution of anticancer agents into the branches of the external carotid artery in IAC models. The CFD simulation results of IAC showed that a curved catheter tip may be placed both below and toward the target artery to increase the distribution rate of anticancer agent into the tumor-feeding artery. Although super-selective IAC can reliably supply anticancer agents to the target artery, high wall shear stress at the target artery can occur, depending on vessel geometry of the patient, and this may cause serious complications during treatment. In case of further research, examination of the fluid structure interaction is vital to account for the elasticity of the vessel walls and catheter movement in many patient models. The noninvasive flow in the patients hepatic artery can be statistically evaluated using CFD, as part of a new dosimetry method for radioembolization of liver cancer. The number of microsphere delivered to different liver segments was calculated at different injection sites based on clinical data. The results showed a varied distribution of 90Y microspheres in liver segments, indicating the importance of CFD patient-specific mimicry for personal dosimetry. Compared with current dosimetry methods, personal dosimetry has the potential to increase the volume brought while reducing the number of microspheres embedded in other healthy tissues. The three-dimensional CFD model for mass drug delivery to tumor nodules during Intraperitoneal (IP) chemotherapy to incorporate actual tumor geometry and flexible vascular properties has also been emphasized on. Dynamic Contrast-Enhanced Magnetic Resonance Imaging (DCE-MRI) imaging has made it possible to differentiate between tumor tissues, healthy surrounding tissues and necrotic areas based on differential vascular structure. We found that the effect of interstitial pressure profiles within tumors was highly dependent on abnormal geometry and location. The depth of the cisplatin penetration of a particular tumor ranged from 0.32 mm to 0.50 mm. It can be stated that the positive relationship between tumor size and the Interstitial Fluid Pressure (IFP) is no longer maintained where there are areas with different vascular conditions, at a same time, a positive relationship has been noted between the percentage of active tumor and the main IFP. In a pancreatic tumor,

there is a growth of blood vessels near the tumor, which allows for the transport of particles to the tumor vasculature. In our study, the geometry is made simpler by focusing on a main artery with pressure influenced flow, and another branching out to the zone of the tumor. The tumor vasculature can be simply thought of as stacked cylinders. Blood flows from P1 to P2 and possibly enters the tumor region at interstitial pressure P3, through a hole. The relation between the three stands as $P1 < P2 < P3$. When a multiphase CFD model combining basic relationships to measure hemodynamics in disturbed flow areas was used, simulation of the two-phase dilute flow simulation of solid human cells in a stable state fits well with test results in vortex formation, reuptake points and velocity profiles. In addition, the linear speed of the cross section in the middle of the vortex area showed the best correlation of the estimated speed. Lung surgery is the only possible treatment for lung cancer. The inevitable removal of active lung tissue and body weight requires a careful and systematic pre-surgery procedure for patients. In particular, the efficacy of post-lung residue surgery needs to be predicted. Clinically, these tests are algorithms based on Pulmonary Function Tests (PFTs). However, these methods ignore the functioning of part of the local airway and provide inland exploration. CFD is designed to provide patient-specific, quantitative, and local information about energy flow and ventilation of the bronchial region.

Of the three main types of cancer treatment surgery, radiation, and chemotherapy chemotherapy is one of the most powerful stages of new success. However, the biggest challenge of chemotherapy is to move the drug to the desired location at a high enough level to work. CFD simulations were used to test the behavior of several vectors of drug delivery in tumor capillaries mainly moving bacteria, non-motile bacteria, and nanoparticles. Red blood cells, bacteria, and nanoparticles are placed in a stream using an immersed border method. The human capillary model was developed using a new method for the management of impaired Red Blood Cells (RBCs). The model was verified with experimental data available from the literature. The stochastic model of bacterial motility was described based on tests of running and falling. Capillary models and bacteria are combined to mimic the intracapillary transport of bacteria. Non-motile bacteria, along with nanoparticles 0. 2 mm, 0. 3 mm, and 0. 405 mm are also presented in the simulation in capillary flow in order to compare them with moving bacteria. Moving bacteria usually swim into a plasma layer near the capillary wall, whereas non-moving bacteria are often trapped in a bolus flow between RBCs. Nanoparticles are strongly influenced by Brownian movements and fluid fluctuations of small scale, so they are not inclined to a single flow area. Motile strains were found to have a long shelf life of 1 mm in length and a high radial velocity. This suggests that migrating bacteria can enter interstitium at a higher rate than nonmotile bacteria or nanoparticles with a diameter between 200 nm and 405 nm. Radiotherapy is a very widely used treatment for cancer. The research trend and success in radiotherapy focuses on high-energy radiation Linear Energy Transfer (LET) radiation as alpha particles. Thus, the alpha-based method is an important means of the development of radiotherapy, and Targeted Alpha Therapy (TAT) is the most important program for the use of alpha particles over the past decade. This paper highlights research work done on TAT computer modeling at Canadian Nuclear Laboratories (CNL). This attempt demonstrates the feasibility of combining CFD mimicry and microdosimetric modeling to evaluate the effectiveness of TAT in a realistic and accurate way. Penn State researchers are learning how cancer cells attach to white blood cells, circulatory protectors, and how blood flow affects this adhesion. The study by Meghan Hoskins a PhD student at Penn State is sponsored by the National Cancer Institute and the PSU Applied Research Laboratory. Her goal is to accurately mimic previous experimental conditions of this condition in order to validate her model for further study of the metastasis process. Tumor cell adhesion can be affected by changes in the flow rate within the micro capillaries of living tumor cells, according to the Canadian Institute for Medical Research (CIMR) by Jennifer Hoskins. Hoskins attempts to demonstrate dynamic and biochemistry activity in the workplace during in vitro cell adhesion. She uses computational fluid dynamics CFD to create a 3D geometric grid for the test room. Researchers at the University of Bristol study how cells interact, collide, and coalesce, and measure the intensity of this activity. So far, Hoskins plans to model two sets of experiments. Cancer cell adhesion to white blood cells is affected by shear level and shear stress, or the energy produced by flow. In the second model, the room is closed, but researchers can watch as the cells interact with each other internally. It aims to understand why and how the migration of cancer cells is affected by the fluid dynamics of the system. If they can detect the presence of cancer-causing proteins in the bloodstream making white blood cells more efficient, perhaps medical researchers could devise a way to block the activity of those proteins. Hoskins hopes that her model will see continuous improvement, especially with regard to simulating adherence within the geometric structure of real blood vessels. Breast cancer usually spreads to the bone marrow. Bone metastatic cancer is incurable and causes widespread bone loss, leading to serious illness and death. Mechanical markers in the skeletal muscle are important environmental parameters that alter tumor formation, osteolysis, and tumor cell-bone signaling, but which mechanisms are most beneficial and the corresponding cell mechanisms are unknown. Interstitial fluid flow based on its well-known role in bone resorption and primary breast cancer was concentrated. A full-scale computer model, based on the microCT of the 3D model of bone metastasis used to predict the internal machinery during in vitro testing was created.

The perfusion used causes uniform dispersion, different liquid speeds and shear pressure on the wall throughout the inside of the scaffold. The liquid velocity distribution and shear pressure on the wall did not change between the model sub-domains of varying widths and spaces. Additionally, the magnitude of these stimuli lies within the range of anabolic mechanical signals in the skeletal structure, which ensures that our 3D model reflects previous in vivo studies using machine-based anabolic loading in the context of bone metastasis.

IV. CONCLUSION

We have made use of Computational Fluid Dynamics (CFD) simulation models to understand the effectiveness of several cancer treatment methodologies while avoiding the huge costs, skills and labour required to conduct the corresponding physical experiments at the same time. CFD also provides the platform of carrying out certain experimentations which are very difficult to be performed under real-life circumstances. CFD is also much less time consuming, it provides quicker and more error-free results. The CFD model outcome of intra-arterial chemotherapy revealed that a curved catheter tip may be placed both below and toward the target artery to increase the distribution rate of anticancer agent into the tumor-feeding artery. Investigation of the fluid model interaction shall serve as a major point of attention to interpret the elasticity of the vessel walls and catheter movement in case studies of patients for future research. The noninvasive flow in the patients hepatic artery can be judged employing statistics, using CFD, as part of a dosimetry method for radioembolization of liver cancer. It can be stated that the positive relationship between tumor size and the Interstitial Fluid Pressure (IFP) is no longer maintained under situations featuring areas with different vascular conditions, at a same time, a positive relationship has been observed between the amount of active tumor and the main IFP. In a pancreatic tumor, there is angiogenesis near the tumor, allowing for the transport of particles to the tumor vasculature. We have kept it simple in our simulation by considering a main artery with pressure driven flow, and another branching out to the zone of the tumor. Additionally, the linear speed of the cross section in the middle of the vortex area showed the best correlation of the estimated speed. CFD is designed to provide patient-specific, quantitative, and local information about energy flow and ventilation of the bronchial region. CFD simulations are used to analyze the behaviour of several vectors of drug delivery in tumor capillaries mainly moving bacteria, non-motile bacteria, and nanoparticles. Capillary models and bacteria are combined to mimic the intracapillary transport of bacteria. We have been able to conveniently infer the results of combining CFD simulation and microdosimetric modeling that has served to evaluate the effectiveness of Targeted Alpha Therapy (TAT) in a realistic and accurate way. In addition to this, We may also conclude that tumor cell adhesion can be affected by changes in the flow rate within the micro capillaries of living tumor cells. Our CFD model provides precise information about perfusion and it may be efficiently used to develop treatments on bone metastasis. However, there is an avenue for further research and diversified study on the employment of CFD for understanding cancer treatment procedures.

Conflict of Interest

The author declares no conflict of interest.

REFERENCES

1. Abramson, D. H., Francis, J. H., & Gobin, Y. P. (2019). What's new in intra-arterial chemotherapy for retinoblastoma? *International Ophthalmology Clinics*, 59(2). <https://doi.org/10.1097/IIO.0000000000000266>
2. Chen, Q., Zhang, B., Dong, Y., Mo, X., Zhang, L., Huang, W., Jiang, H., Xia, J., & Zhang, S. (2018). Comparison between intravenous chemotherapy and intra-arterial chemotherapy for retinoblastoma: A meta-analysis. *BMC Cancer*, 18(1). <https://doi.org/10.1186/s12885-018-4406-6>
3. Erratum: Intra-arterial chemotherapy in retinoblastoma - A paradigm change (*Indian Journal of Ophthalmology* (2019) 67 (740-754) DOI: 10.4103/ijo.IJO_866_19). (2019). In *Indian Journal of Ophthalmology* (Vol. 67, Issue 8). <https://doi.org/10.4103/0301-4738.263157>
4. Homma, A., Onimaru, R., Matsuura, K., Robbins, K. T., & Fujii, M. (2016). Intra-arterial chemoradiotherapy for head and neck cancer. In *Japanese Journal of Clinical Oncology* (Vol. 46, Issue 1). <https://doi.org/10.1093/jjco/hyv151>
5. Pekacka, A. (2020). The Role of Intraarterial Chemotherapy in the Management of Retinoblastoma. In *Journal of Ophthalmology* (Vol. 2020). <https://doi.org/10.1155/2020/3638410>



6. Rishi, P. , Agarwal, A. , Chatterjee, P. , Sharma, T. , Sharma, M. , Saravanan, M. , & Ravikumar, R. (2020). Intra-Arterial Chemotherapy for Retinoblastoma: Four-Year Results from Tertiary Center in India. *Ocular Oncology and Pathology*, 6(1). <https://doi.org/10.1159/000500010>
7. Shields, C. L. , Dockery, P. W. , Yaghy, A. , Duffner, E. R. , Levin, H. J. , Taylor, O. S. , Sajjadi, Z. , Lally, S. E. , Shields, J. A. , Rosenwasser, R. , Tjournakaris, S. , & Jabbour, P. (2021). Intra-arterial chemotherapy for retinoblastoma in 341 consecutive eyes (1,292 infusions): comparative analysis of outcomes based on patient age, race, and sex. *Journal of AAPOS*, 25(3). <https://doi.org/10.1016/j.jaapos.2020.12.006>
8. Shields, C. L. , Manjandavida, F. P. , Lally, S. E. , Pieretti, G. , Arepalli, S. A. , Caywood, E. H. , Jabbour, P. , & Shields, J. A. (2014). Intra-arterial chemotherapy for retinoblastoma in 70 eyes: Outcomes based on the international classification of retinoblastoma. *Ophthalmology*, 121(7). <https://doi.org/10.1016/j.ophtha.2014.01.026>
9. Sweid, A. , Hammoud, B. , Weinberg, J. H. , Texakalidis, P. , Xu, V. , Shivashankar, K. , Baldassari, M. P. , Das, S. , Ramesh, S. , Tjournakaris, S. , Shields, C. L. , Ancona-Lezama, D. , Lim, L. A. S. , Dalvin, L. A. , & Jabbour, P. (2020). Intra-arterial chemotherapy for retinoblastoma in infants ≤ 10 kg: 74 treated eyes with 222 IAC sessions. *American Journal of Neuroradiology*, 41(7). <https://doi.org/10.3174/AJNR.A6590>
10. Zanaty, M. , Barros, G. , Chalouhi, N. , Starke, R. M. , Manasseh, P. , Tjournakaris, S. I. , Shields, C. L. , Hasan, D. , Bulsara, K. , Rosenwasser, R. H. , & Jabbour, P. (2014). Update on intra-arterial chemotherapy for retinoblastoma. In *Scientific World Journal* (Vol. 2014). <https://doi.org/10.1155/2014/869604>
11. Anwanwan, D. , Singh, S. K. , Singh, S. , Saikam, V. , & Singh, R. (2020). Challenges in liver cancer and possible treatment approaches. In *Biochimica et Biophysica Acta - Reviews on Cancer* (Vol. 1873, Issue 1). <https://doi.org/10.1016/j.bbcan.2019.188314>
12. Bester, L. , Meteling, B. , Boshell, D. , Chua, T. C. , & Morris, D. L. (2014). Transarterial chemoembolisation and radioembolisation for the treatment of primary liver cancer and secondary liver cancer: A review of the literature. *Journal of Medical Imaging and Radiation Oncology*, 58(3). <https://doi.org/10.1111/1754-9485.12163>
13. Feng, M. , Pan, Y. , Kong, R. , & Shu, S. (2020). Therapy of Primary Liver Cancer. In *The Innovation* (Vol. 1, Issue 2). <https://doi.org/10.1016/j.xinn.2020.100032>
14. Fujita, M. , Yamaguchi, R. , Hasegawa, T. , Shimada, S. , Arihiro, K. , Hayashi, S. , Maejima, K. , Nakano, K. , Fujimoto, A. , Ono, A. , Aikata, H. , Ueno, M. , Hayami, S. , Tanaka, H. , Miyano, S. , Yamaue, H. , Chayama, K. , Kakimi, K. , Tanaka, S. , ... Nakagawa, H. (2020). Classification of primary liver cancer with immunosuppression mechanisms and correlation with genomic alterations. *EBioMedicine*, 53. <https://doi.org/10.1016/j.ebiom.2020.102659>
15. Ma, L. , Hernandez, M. O. , Zhao, Y. , Mehta, M. , Tran, B. , Kelly, M. , Rae, Z. , Hernandez, J. M. , Davis, J. L. , Martin, S. P. , Kleiner, D. E. , Hewitt, S. M. , Ylaya, K. , Wood, B. J. , Greten, T. F. , & Wang, X. W. (2019). Tumor Cell Biodiversity Drives Microenvironmental Reprogramming in Liver Cancer. *Cancer Cell*, 36(4). <https://doi.org/10.1016/j.ccell.2019.08.007>
16. Qiu, Z. , Li, H. , Zhang, Z. , Zhu, Z. , He, S. , Wang, X. , Wang, P. , Qin, J. , Zhuang, L. , Wang, W. , Xie, F. , Gu, Y. , Zou, K. , Li, C. , Li, C. , Wang, C. , Cen, J. , Chen, X. , Shu, Y. , ... Hui, L. (2019). A Pharmacogenomic Landscape in Human Liver Cancers. *Cancer Cell*, 36(2). <https://doi.org/10.1016/j.ccell.2019.07.001>
17. Rashed, W. M. , Kandeil, M. A. M. , Mahmoud, M. O. , & Ezzat, S. (2020). Hepatocellular Carcinoma (HCC) in Egypt: A comprehensive overview. In *Journal of the Egyptian National Cancer Institute* (Vol. 32, Issue 1). <https://doi.org/10.1186/s43046-020-0016-x>
18. Sia, D. , Villanueva, A. , Friedman, S. L. , & Llovet, J. M. (2017). Liver Cancer Cell of Origin, Molecular Class, and Effects on Patient Prognosis. In *Gastroenterology* (Vol. 152, Issue 4). <https://doi.org/10.1053/j.gastro.2016.11.048>
19. Subramaniam, S. , Kelley, R. K. , & Venook, A. P. (2013). A review of hepatocellular carcinoma (HCC) staging systems. In *Chinese Clinical Oncology* (Vol. 2, Issue 4). <https://doi.org/10.3978/j.issn.2304-3865.2013.07.05>
20. Wang, Z. G. , He, Z. Y. , Chen, Y. Y. , Gao, H. , & Du, X. L. (2021). Incidence and survival outcomes of secondary liver cancer: A surveillance epidemiology and end results database analysis. *Translational Cancer Research*, 10(3). <https://doi.org/10.21037/tcr-20-3319>
21. H. Kitajima, T. Iwai, Y. Yajima, K. Mitsudo, Computational Fluid Dynamics Study of Superselective Intra-arterial Chemotherapy for Oral Cancer: Flow Simulation of Anticancer Agent in the Linguofacial Trunk, Applied Sciences 10 (2020) 7496.
22. Sangro, B. , Iñarrairaegui, M. , & Bilbao, J. I. (2012). Radioembolization for hepatocellular carcinoma. In *Journal of Hepatology* (Vol. 56, Issue 2). <https://doi.org/10.1016/j.jhep.2011.07.012>



23. Steuperaert, M. , Debbaut, C. , Carlier, C. , De Wever, O. , Descamps, B. , Vanhove, C. , Ceelen, W. , & Segers, P. (2019). A 3D CFD model of the interstitial fluid pressure and drug distribution in heterogeneous tumor nodules during intraperitoneal chemotherapy. *Drug delivery*, 26(1), 404–415. <https://doi.org/10.1080/10717544.2019.1588423>
24. Chakraborty, N. , & Basu, S. (2020). Computational modeling of perfusion in dense solid tumors. *arXiv preprint arXiv:2012.04095*.
25. T. Gemci, V. Ponyavin, Y. Chen, H. Chen, R. Collins, Computational model of airflow in upper 17 generations of human respiratory tract, *Journal of Biomechanics*, Volume 41, Issue 9, 2008, Pages 2047-2054, ISSN 0021-9290, <https://doi.org/10.1016/j.jbiomech.2007.12.019>.
26. Maani, Nazanin (2019): CFD MODELING IN DESIGN AND EVALUATION OF AN ENDOVASCULAR CHEMOFILTER DEVICE. Purdue University Graduate School. Thesis. <https://doi.org/10.25394/PGS.11303480.v1>
27. Wenbo Zhan, Moath Alamer, Xiao Yun Xu, Computational modelling of drug delivery to solid tumour: Understanding the interplay between chemotherapeutics and biological system for optimised delivery systems, *Advanced Drug Delivery Reviews*, Volume 132, 2018, Pages 81-103, ISSN 0169-409X, <https://doi.org/10.1016/j.addr.2018.07.013>.
28. Tao Xu, Yu Liu, Tong Liu, Gang Li, Nusret Aydemir, Multiphysics Modelling of Background Dose by Systemic Targeted Alpha Therapy, *Journal of Medical Imaging and Radiation Sciences*, Volume 49, Issue 3, 2018, Pages 270-276, ISSN 1939-8654, <https://doi.org/10.1016/j.jmir.2018.06.002>.
29. Gautier Follain, Naël Osmani, Ana Sofia Azevedo, Guillaume Allio, Luc Mercier, Matthia A. Karreman, Gergely Solecki, Maria Jesús García Leòn, Olivier Lefebvre, Nina Fekonja, Claudia Hille, Vincent Chabannes, Guillaume Dollé, Thibaut Metivet, François Der Hovsepian, Christophe Prudhomme, Angélique Pichot, Nicodème Paul, Raphaël Carapito, Siamak Bahram, Bernhard Ruthensteiner, André Kemmling, Susanne Siemonsen, Tanja Schneider, Jens Fiehler, Markus Glatzel, Frank Winkler, Yannick Schwab, Klaus Pantel, Sébastien Harlepp, Jacky G. Goetz, Hemodynamic Forces Tune the Arrest, Adhesion, and Extravasation of Circulating Tumor Cells, *Developmental Cell*, Volume 45, Issue 1, 2018, Pages 33-52. e12, ISSN 1534-5807, <https://doi.org/10.1016/j.devcel.2018.02.015>.
30. Boyuan Liu, Suyue Han, Brandon P. Hedrick, Yahya Modarres-Sadeghi, Maureen E. Lynch, Perfusion applied to a 3D model of bone metastasis results in uniformly dispersed mechanical stimuli, Volume 115, Issue 4, April 2018, Pages 1076-1085, <https://doi.org/10.1002/bit.26524>.
31. Bela Soni, Shahrouz Aliabadi, Large-scale CFD simulations of airflow and particle deposition in lung airway, *Computers & Fluids*, Volume 88, 2013, Pages 804-812, ISSN 0045-7930, <https://doi.org/10.1016/j.compfluid.2013.06.015>.
32. Ali, D., Ozalp, M., Blanquer, S. B. G., & Onel, S. (2020). Permeability and fluid flow-induced wall shear stress in bone scaffolds with TPMS and lattice architectures: A CFD analysis. *European Journal of Mechanics, B/Fluids*, 79. <https://doi.org/10.1016/j.euromechflu.2019.09.015>
33. Aliboni, L., Pennati, F., Sarti, M., Iorio, V., Carrinola, R., Palleschi, A., & Aliverti, A. (2021). Computational Fluid Dynamics (CFD) Analysis of Subject-specific Bronchial Tree Models in Lung Cancer Patients. *Proceedings of the Annual International Conference of the IEEE Engineering in Medicine and Biology Society, EMBS*. <https://doi.org/10.1109/EMBC46164.2021.9629765>
34. Aliboni, Lorenzo, Tullio, M., Pennati, F., Lomauro, A., Carrinola, R., Carrafiello, G., Nosotti, M., Palleschi, A., & Aliverti, A. (2022). Functional analysis of the airways after pulmonary lobectomy through computational fluid dynamics. *Scientific Reports*, 12(1). <https://doi.org/10.1038/s41598-022-06852-x>
35. Aramburu, J., Antón, R., Rivas, A., Ramos, J. C., Sangro, B., & Bilbao, J. I. (2018). Liver Radioembolization: An Analysis of Parameters that Influence the Catheter-Based Particle-Delivery via CFD. *Current Medicinal Chemistry*, 27(10). <https://doi.org/10.2174/0929867325666180622145647>
36. Beckett, E. M., Cyrs, W. D., Abelman, A., Monnot, A. D., Gaffney, S. H., & Finley, B. L. (2019). Derivation of an occupational exposure limit for diacetyl using dose-response data from a chronic animal inhalation exposure study. In *Journal of Applied Toxicology* (Vol. 39, Issue 5). <https://doi.org/10.1002/jat.3757>
37. Chari, S., Sridhar, K., Walenga, R., & Kleinstreuer, C. (2021). Computational analysis of a 3D mucociliary clearance model predicting nasal drug uptake. *Journal of Aerosol Science*, 155. <https://doi.org/10.1016/j.jaerosci.2021.105757>
38. Deng, F., Liu, L., Li, Z., & Liu, J. (2021). 3D printed Ti6Al4V bone scaffolds with different pore structure effects on bone ingrowth. *Journal of Biological Engineering*, 15(1). <https://doi.org/10.1186/s13036-021-00255-8>



39. Dieguez-Elizondo, P. M., O'Donohoe, P. G., Gil-Lopez, T., Castejon-Navas, J., & Gálvez-Huerta, M. A. (2019). Calculation methods of Radon-222 radiological activity for NORM plant with ventilation. *Journal of Petroleum Science and Engineering*, 183. <https://doi.org/10.1016/j.petrol.2019.106360>
40. Kantová, N., Jandačka, J., Sládek, S., Holubčík, M., & Čaja, A. (2019). The flowing of particulate matter through baffles depending on their number in the flue gas tract. *Transportation Research Procedia*, 40. <https://doi.org/10.1016/j.trpro.2019.07.102>
41. Liu, Z., Sun, D., Zhu, Q., & Liu, X. (2021). The screening of immune-related biomarkers for prognosis of lung adenocarcinoma. *Bioengineered*, 12(1). <https://doi.org/10.1080/21655979.2021.1911211>
42. Longest, P. W., Tian, G., Walenga, R. L., & Hindle, M. (2012). Comparing MDI and DPI aerosol deposition using in vitro experiments and a new stochastic individual path (SIP) model of the conducting airways. *Pharmaceutical Research*, 29(6). <https://doi.org/10.1007/s11095-012-0691-y>
43. Pires, T., Dunlop, J. W. C., Fernandes, P. R., & Castro, A. P. G. (2022). Challenges in computational fluid dynamics applications for bone tissue engineering. In *Proceedings of the Royal Society A: Mathematical, Physical and Engineering Sciences* (Vol. 478, Issue 2257). <https://doi.org/10.1098/rspa.2021.0607>
44. Roncali, E., Taebi, A., Foster, C., & Vu, C. T. (2020). Personalized Dosimetry for Liver Cancer Y-90 Radioembolization Using Computational Fluid Dynamics and Monte Carlo Simulation. *Annals of Biomedical Engineering*, 48(5). <https://doi.org/10.1007/s10439-020-02469-1>
45. Sedakov, I., Dubinina, V., Bondar, O., Lukianchuk, O., & Zavoloka, O. (2016). COMPARATIVE CHARACTERISTICS OF THE METHODS OF CHEMODRUGS ADMINISTRATION IN THE TREATMENT OF PATIENTS WITH LOCALLY-ADVANCED BREAST CANCER. *EUREKA: Health Sciences*, 1. <https://doi.org/10.21303/2504-5679.2016.00038>
46. Solodkiy, V. A., Sotnikov, V. M., Chkhikvadze, V. D., Nudnov, N. V., & Trotsenko, S. D. (2022). Impact of Postoperative Radiotherapy on Overall Survival in Patients with pN1 Non-Small Cell Lung Cancer. *Journal of Radiology and Nuclear Medicine*, 102(6). <https://doi.org/10.20862/0042-4676-2021-102-6-338-348>
47. Tsuzuki, Y., Roncali, E., Cherry, S. R., & Aldredge, R. C. (2017). The effect of hepatic-artery geometry on the particle release map. *International Conference on Fluid Flow, Heat and Mass Transfer*. <https://doi.org/10.11159/ffhmt17.122>
48. Villa-Roel, N., Ryu, K., Gu, L., Fernandez Esmerats, J., Kang, D. W., Kumar, S., & Jo, H. (2022). Hypoxia inducible factor 1 α inhibitor PX-478 reduces atherosclerosis in mice. *Atherosclerosis*, 344. <https://doi.org/10.1016/j.atherosclerosis.2022.01.002>
49. Xi, J., Si, X. A., Kim, J., Zhang, Y., Jacob, R. E., Kabilan, S., & Corley, R. A. (2016). Anatomical Details of the Rabbit Nasal Passages and Their Implications in Breathing, Air Conditioning, and Olfaction. *Anatomical Record*, 299(7). <https://doi.org/10.1002/ar.23367>
50. Yang, A. S., Chuang, F. C., Chen, C. K., Lee, M. H., Chen, S. W., Su, T. L., & Yang, Y. C. (2015). A high-performance micromixer using three-dimensional Tesla structures for bio-applications. *Chemical Engineering Journal*, 263. <https://doi.org/10.1016/j.cej.2014.11.034>
51. Yoo, S. J., & Ito, K. (2018a). Assessment of transient inhalation exposure using in silico human model integrated with PBPK-CFD hybrid analysis. *Sustainable Cities and Society*, 40. <https://doi.org/10.1016/j.scs.2018.04.023>
52. Yoo, S. J., & Ito, K. (2018b). Numerical prediction of tissue dosimetry in respiratory tract using computer simulated person integrated with physiologically based pharmacokinetic-computational fluid dynamics hybrid analysis. *Indoor and Built Environment*, 27(7). <https://doi.org/10.1177/1420326X17694475>



INNO SPACE
SJIF Scientific Journal Impact Factor
Impact Factor: 8.118



ISSN INTERNATIONAL
STANDARD
SERIAL
NUMBER
INDIA



INTERNATIONAL JOURNAL OF INNOVATIVE RESEARCH

IN SCIENCE | ENGINEERING | TECHNOLOGY

 **9940 572 462**  **6381 907 438**  **ijirset@gmail.com**



www.ijirset.com

Scan to save the contact details