



e-ISSN: 2319-8753 | p-ISSN: 2347-6710

IJIRSET

International Journal of Innovative Research in
SCIENCE | ENGINEERING | TECHNOLOGY



INTERNATIONAL JOURNAL OF INNOVATIVE RESEARCH

IN SCIENCE | ENGINEERING | TECHNOLOGY

Volume 11, Issue 9, September 2022

ISSN INTERNATIONAL
STANDARD
SERIAL
NUMBER
INDIA

Impact Factor: 8.118



9940 572 462



6381 907 438



ijirset@gmail.com



www.ijirset.com

Blood Flow Restriction Training (BFRT): Definition, Historical Context and Development

Karan Pal Vishnoi, Dr. Sushil Kumar

Ph.D Research Scholar, Department of Physical Education, Sunrise University, Alwar (Rajasthan), India

Assistant Professor, Department of Physical Education, Sunrise University, Alwar (Rajasthan), India

ABSTRACT: Blood Flow Restriction Training (BFRT) is a progressive training modality that combines low-intensity resistance exercises with controlled partial occlusion of blood flow to the working muscles, leading to significant muscle hypertrophy and strength adaptations similar to those achieved through traditional high-intensity resistance training. This technique, first introduced in the 1960s in Japan under the name KAATSU training, has grown in popularity across various fields, including athletic performance, rehabilitation, and clinical applications, due to its ability to induce muscle growth while minimizing the mechanical load on joints and connective tissues.

The purpose of this paper is to explore the definition of BFRT, its physiological mechanisms, and its historical development, as well as the current scientific consensus surrounding its use in both healthy populations and those with limitations. BFRT allows individuals to achieve hypertrophic and strength-related outcomes with loads as low as 20-30% of their one-repetition maximum (1RM), making it especially beneficial for populations such as the elderly, post-surgical patients, or athletes recovering from injuries who cannot tolerate the heavy loads of traditional resistance training.

The physiological mechanisms of BFRT revolve around the restriction of venous return while maintaining arterial inflow, creating a hypoxic and metabolically stressed environment in the muscle. This triggers several key processes, including increased metabolic byproduct accumulation, muscle fiber recruitment, and the enhanced release of anabolic hormones, such as growth hormone and insulin-like growth factor-1 (IGF-1). These adaptations stimulate muscle protein synthesis and hypertrophy, mimicking the effects of high-intensity exercise. Research also shows that BFRT can increase the recruitment of fast-twitch muscle fibers, which are usually engaged during heavy lifting, further contributing to muscle growth and strength gains.

Over the decades, technological advancements in BFRT equipment, such as the development of pneumatic cuffs with controlled pressure monitoring, have improved the safety and effectiveness of the technique, allowing for widespread adoption in clinical and fitness settings. BFRT has been successfully applied in various domains, including athletic performance enhancement, rehabilitation from injury, and the management of muscle atrophy in aging populations. However, despite its numerous benefits, BFRT is not without risks. Improper use or excessive pressure can lead to complications such as blood clots, nerve damage, or ischemia, underscoring the importance of proper training, supervision, and adherence to safety protocols.

This paper also discusses the future prospects of BFRT, including its potential to revolutionize resistance training by offering an alternative to traditional high-load exercises. Future research directions include optimizing BFRT protocols, investigating its long-term effects on muscle health, and expanding its application in special populations, such as individuals with cardiovascular conditions or chronic diseases. Furthermore, advancements in digital tools, such as real-time muscle oxygenation monitoring, promise to enhance the precision and safety of BFRT, paving the way for its integration into personalized training and rehabilitation programs.

KEYWORDS: Blood Flow Restriction Training (BFRT), Muscle Hypertrophy, Occlusion Training, Low-Intensity Resistance Exercise, Rehabilitation, Strength Training

I. IMPORTANCE OF BFRT IN MODERN TRAINING

The rise of BFRT is largely attributed to its efficacy in reducing the mechanical stress placed on the body during traditional strength training. This is particularly advantageous for older adults, those recovering from injury, and athletes looking to improve muscle mass and strength without risking overuse injuries. In contrast to traditional high-

intensity resistance training, BFRT can be used in contexts where heavy weights are impractical or unsafe, thereby broadening the accessibility of strength training to a wider demographic.

Purpose of the Study: The purpose of this paper is to provide a thorough exploration of BFRT, including its definition, historical background, mechanisms, and practical applications. It also aims to examine the benefits and risks associated with the technique and explore its potential for future development in the fields of fitness and rehabilitation.

Scope and Limitations: This paper covers BFRT's role in enhancing muscle growth and rehabilitation, focusing on its historical development and current scientific research. While it provides an in-depth analysis of BFRT's benefits and risks, it does not cover specific training protocols for individual needs or delve into the psychological aspects of training.

Overview of Research Methodology: The research methodology involves an extensive review of peer-reviewed studies, clinical trials, and historical documents related to BFRT. The data is analyzed to provide a comprehensive understanding of the method's efficacy and safety in various settings, including sports performance and clinical rehabilitation.

II. INTRODUCTION

Blood Flow Restriction Training (BFRT) is a game-changer in the world of physical fitness and rehabilitation, allowing individuals to achieve muscle growth and strength gains through low-intensity exercise, making it accessible to populations with different fitness levels and physical limitations. Its increasing popularity is evident in both athletic and clinical settings due to its ability to promote hypertrophy without subjecting the body to the heavy loads typically required for muscle growth.

Definition of Blood Flow Restriction Training (BFRT) : BFRT, also known as occlusion training, is a specialized technique that involves the partial restriction of blood flow to a specific muscle group during low-intensity exercise. This restricted blood flow stimulates muscular adaptations normally seen in high-intensity resistance training, despite the use of lighter weights or resistance.

BFRT involves the application of cuffs or bands to the upper or lower limbs to restrict venous blood flow while allowing arterial inflow during exercise. This restriction creates an oxygen-deprived environment in the muscle, increasing metabolic stress and promoting muscle growth even at low exercise intensities. The goal is to create a hypoxic environment within the muscles, simulating the effects of heavy lifting while minimizing joint and tendon stress.

The fundamental principle of BFRT is occlusion, which reduces the ability of the muscle to clear metabolic byproducts such as lactic acid. This accumulation of metabolites, combined with the mechanical tension generated by low-intensity exercise, signals the body to initiate muscle protein synthesis and other anabolic processes. The most common technique involves applying pressure cuffs or elastic bands to the limbs and performing resistance exercises at around 20-30% of the individual's one-repetition maximum (1RM), a fraction of the load typically required for muscle hypertrophy.

Traditional strength training requires high-intensity loads, typically 70-80% of an individual's 1RM, to induce significant muscle hypertrophy. BFRT, on the other hand, achieves similar hypertrophic and strength gains with much lighter loads by amplifying the metabolic stress on the muscles. This makes BFRT an ideal option for individuals with joint issues or those recovering from injuries, as the lighter loads reduce strain on the skeletal and muscular systems.

The scientific consensus is that BFRT is an effective method for increasing muscle size and strength, particularly in populations that are unable to engage in traditional high-load resistance training. Multiple studies have confirmed the method's efficacy, particularly in clinical settings for rehabilitation, but there are ongoing debates regarding its long-term safety and the best practices for implementation in both athletic and medical contexts.

III. HISTORICAL CONTEXT OF BFRT

The concept of blood flow restriction training originated in the 1960s, but it wasn't until the 2000s that it gained widespread recognition as a legitimate form of exercise. Its development can be traced to Japan, where Dr. Yoshiaki Sato first experimented with what he called "KAATSU training," a precursor to modern BFRT.

Dr. Yoshiaki Sato, a Japanese scientist, is credited with developing the first version of BFRT in the early 1960s after observing the effects of restricted blood flow on muscle growth. Through self-experimentation, Dr. Sato began to refine his methods, using bicycle inner tubes to restrict blood flow to his limbs while performing basic exercises. His early research demonstrated that muscle hypertrophy could be achieved with light weights when blood flow to the working muscles was restricted, which laid the foundation for modern BFRT practices.

Dr. Sato's work was further developed by several key researchers, including Dr. Jim Stray-Gundersen and Dr. Jeremy Loenneke, who contributed significantly to understanding the physiological mechanisms behind BFRT. These pioneers helped bring BFRT into the scientific spotlight and facilitated its acceptance in clinical and athletic training communities.

Over the decades, BFRT has evolved from a rudimentary practice using basic equipment to a highly sophisticated training method. As its popularity grew, researchers began to explore its potential applications in both fitness and rehabilitation. Advances in technology have allowed for more precise control of blood flow restriction, leading to safer and more effective applications in various contexts.

Numerous case studies have demonstrated the effectiveness of BFRT in enhancing muscle growth and recovery. For example, studies have shown that BFRT can significantly reduce the time required for rehabilitation following ACL surgery, making it a valuable tool in physical therapy. Additionally, professional athletes have used BFRT to maintain muscle mass and strength during recovery periods when high-intensity training is not feasible.

IV. MECHANISMS OF BFRT

The primary mechanism by which BFRT works is through the creation of a hypoxic (low oxygen) environment within the muscle, leading to metabolic stress. This triggers several physiological responses, including increased muscle protein synthesis, enhanced muscle growth, and the production of growth hormones.

When blood flow to the muscles is restricted, the muscle becomes deprived of oxygen (hypoxia), leading to an accumulation of metabolites such as lactic acid. This metabolic stress triggers a cascade of cellular signals that promote muscle protein synthesis, leading to hypertrophy. The combination of mechanical tension from the exercise and metabolic stress is believed to be the primary driver of muscle growth during BFRT.

One of the major advantages of BFRT is its ability to produce muscle growth with low-intensity exercise. Traditional strength training relies on heavy loads to stimulate muscle fibers, but BFRT achieves similar results by creating an environment that mimics the effects of heavy lifting, even though the loads used are significantly lighter.

The reduced blood flow causes an accumulation of metabolites, including lactate, which is believed to stimulate muscle growth. This metabolic stress increases the production of anabolic hormones, such as growth hormone and insulin-like growth factor-1 (IGF-1), which are critical for muscle repair and growth. Additionally, BFRT enhances the recruitment of fast-twitch muscle fibers, which are typically activated during high-intensity exercise.

Research shows that BFRT significantly increases the production of growth hormone, which plays a critical role in muscle repair and growth. After a BFRT session, growth hormone levels can increase by as much as 290%, promoting both muscle hypertrophy and faster recovery. This hormonal response is one of the key mechanisms that makes BFRT so effective, especially when combined with low-intensity exercises. The endocrine response to BFRT helps facilitate muscle growth without placing excessive mechanical strain on the joints and tendons, which is particularly beneficial for those recovering from injuries or with joint limitations.

BFRT is versatile and applicable in a wide range of contexts, from enhancing athletic performance to aiding in rehabilitation and improving the quality of life for aging populations. Its ability to stimulate muscle growth and strength using lower loads makes it an attractive option for individuals across various fields and physical conditions.

For athletes, BFRT can be a powerful tool to increase muscle size and strength without the risk of overuse injuries associated with heavy lifting. During periods of recovery or off-season, when athletes want to maintain muscle mass and strength while reducing the risk of injury, BFRT can serve as a lower-risk alternative to traditional resistance training. Athletes from various disciplines, including bodybuilding, soccer, and track and field, have incorporated BFRT into their training regimens to achieve strength gains with reduced joint stress.

One of the most significant applications of BFRT is in rehabilitation. Patients recovering from surgery, such as anterior cruciate ligament (ACL) repair, or from injuries that limit their ability to lift heavy loads, can benefit from BFRT. Physical therapists have found that BFRT allows patients to regain muscle strength and mass without putting excessive strain on healing tissues. Additionally, studies have shown that incorporating BFRT into rehabilitation programs can reduce the time it takes to return to full function compared to traditional methods.

BFRT has been used to help patients who are suffering from muscle atrophy due to conditions like sarcopenia (age-related muscle loss) or prolonged immobility. For older adults or individuals with chronic diseases who may not be able to perform high-intensity resistance exercises, BFRT provides an effective alternative to prevent muscle loss and maintain functional independence. Clinicians use BFRT to improve strength and mobility while minimizing the risk of injury.

As people age, muscle mass naturally decreases, leading to a decline in functional abilities and an increased risk of falls and other injuries. BFRT offers a solution for older adults to maintain muscle mass and strength with lower intensity exercises, making it safer and more practical for this demographic. Studies have shown that older adults who engage in BFRT experience improvements in muscle mass, strength, and overall mobility, contributing to better quality of life and independence.

V. BENEFITS AND RISKS OF BFRT

Like any training modality, BFRT comes with its set of advantages and potential risks. Understanding these factors is crucial for determining whether BFRT is appropriate for certain individuals and for ensuring that it is applied safely.

BFRT offers numerous advantages for both athletes and non-athletes. One of the most significant benefits is that it allows for substantial strength and hypertrophy gains with low-intensity exercises, reducing the risk of injury associated with heavy lifting. This is particularly beneficial for individuals recovering from injury, those with joint or mobility issues, and older adults. Additionally, BFRT can help to improve muscular endurance, enhance cardiovascular function, and increase the production of anabolic hormones, which play a key role in muscle repair and growth.

While BFRT is generally considered safe when applied correctly, there are potential risks, particularly if the technique is not executed properly. Excessive occlusion pressure or improper application of the bands can lead to complications such as nerve damage, blood clots, or muscle ischemia. Additionally, individuals with certain medical conditions, such as cardiovascular disease, should consult with a healthcare professional before engaging in BFRT. Overuse of BFRT without adequate recovery can also lead to excessive muscle damage or delayed-onset muscle soreness (DOMS).

To minimize risks, it is essential to follow established guidelines for the safe application of BFRT. This includes using appropriate occlusion pressures based on the individual's limb size, monitoring the duration of BFRT sessions, and ensuring that exercises are performed under the supervision of trained professionals. The cuffs or bands used for BFRT should be specifically designed for this purpose, as using makeshift bands or applying too much pressure can increase the risk of adverse effects.

To mitigate risks, it is important to assess the individual's health status before implementing BFRT. Individuals with cardiovascular, vascular, or neurological conditions should be evaluated by a medical professional before starting BFRT. Additionally, using pneumatic cuffs with built-in pressure sensors can help ensure that the correct amount of occlusion is applied, reducing the risk of complications. Proper education and supervision are also crucial in ensuring that BFRT is performed safely.

VI. DEVELOPMENT OF BFRT TECHNOLOGY

The technological advancements in BFRT equipment have played a crucial role in its safety, efficacy, and widespread adoption. As the demand for BFRT has grown, so too has the development of specialized tools and technologies that allow for precise control over blood flow restriction during training sessions.

Early versions of BFRT involved the use of rudimentary bands or makeshift equipment to restrict blood flow. Today, however, specialized equipment, such as pneumatic cuffs and automated systems, are available that allow for more controlled and safer blood flow restriction. These devices often include pressure gauges or sensors to monitor occlusion levels and ensure that blood flow restriction is applied appropriately. This advancement has made BFRT safer and more accessible, particularly in clinical and rehabilitation settings where precision is key.

Modern BFRT devices allow for adjustable and consistent pressure, reducing the risk of over-occlusion or under-occlusion. These devices can be programmed to apply the ideal pressure based on the individual's limb size, ensuring that blood flow is restricted enough to create the desired muscle adaptations without compromising safety. The integration of these devices into gyms, rehabilitation centers, and even home fitness equipment has broadened the use of BFRT in various settings.

As BFRT has grown in popularity, the use of technology to monitor and control training sessions has become more widespread. Digital tools, such as mobile apps and wearable sensors, can now be used to track an individual's progress during BFRT sessions. These tools provide real-time feedback on muscle activation, oxygenation levels, and overall performance, allowing for more personalized and optimized training programs.

The use of digital tools in BFRT has improved the ability to fine-tune training protocols and ensure safety. For example, some devices are equipped with sensors that monitor oxygen levels in the muscle during exercise, allowing trainers and clinicians to adjust the pressure of the cuffs based on real-time data. This ensures that the individual is receiving the right amount of occlusion for their specific goals and physical condition, making BFRT more efficient and safer.

Blood Flow Restriction Training (BFRT) has emerged as a revolutionary technique in the fields of fitness, rehabilitation, and sports performance. Its unique approach of combining low-intensity resistance exercise with partial occlusion of blood flow offers significant benefits in terms of muscle hypertrophy, strength gains, and recovery, without the need for the heavy loads typically required in traditional resistance training. This makes it an appealing option for a wide range of populations, including athletes, older adults, and individuals recovering from injury or surgery.

Historically, BFRT has developed significantly from its early beginnings in Japan with KAATSU training to its widespread use across the globe. The introduction of more advanced, controlled equipment has improved the safety and accessibility of BFRT, allowing it to become a staple in both clinical and performance-based settings. This has led to its application not only for enhancing athletic performance but also as an effective tool for preventing muscle atrophy in populations such as the elderly or individuals with chronic diseases.

BFRT represents a significant development in the field of strength training and rehabilitation, offering a safe and effective alternative to high-intensity resistance training. As BFRT continues to evolve, its applications in both clinical and athletic contexts are likely to expand, providing new opportunities for individuals across a broad range of physical conditions to improve muscle strength and function while minimizing the risks associated with traditional resistance training.

Blood Flow Restriction Training (BFRT) is an innovative training method that combines low-intensity resistance exercise with the partial occlusion of blood flow, leading to muscle hypertrophy and strength gains similar to traditional high-intensity training. This method has become increasingly popular in both fitness and rehabilitation settings due to its ability to enhance muscle growth with reduced mechanical stress on joints. This paper explores the definition of BFRT, its historical development, mechanisms of action, applications, and potential risks and benefits. Furthermore, it discusses the advancement of BFRT technology and provides recommendations for its safe implementation. The future potential of BFRT in both clinical and athletic contexts is also addressed, providing insight into its growing importance in the fitness and rehabilitation fields.

VII. CONCLUSION

Through the review of current research, it is clear that BFRT effectively stimulates the physiological processes that lead to muscle growth, largely by increasing metabolic stress and the recruitment of fast-twitch muscle fibers. Additionally, the endocrine response to BFRT, which includes heightened levels of growth hormone and insulin-like growth factor-1 (IGF-1), further enhances muscle protein synthesis and recovery. These responses, typically associated with high-intensity training, can now be achieved with much lower physical stress on joints and connective tissues, which is particularly beneficial for individuals with limited mobility or those at risk of injury.

However, despite its many advantages, BFRT is not without risks. Improper application, such as using excessive pressure or not following established guidelines, can lead to serious complications, including nerve damage, blood clots, or muscle ischemia. Therefore, it is essential for BFRT to be performed under proper supervision, particularly in clinical settings. The use of modern equipment, such as pneumatic cuffs with built-in pressure control, can help mitigate some of these risks and ensure that the technique is applied safely and effectively.

Looking ahead, the future of BFRT is promising. There is considerable potential for further refinement of BFRT protocols, including individualized approaches that optimize pressure, duration, and frequency of training for different populations. The integration of digital tools, such as real-time monitoring of blood flow and muscle oxygenation, could further enhance the precision of BFRT, making it even safer and more effective. Additionally, as more research emerges, BFRT may find new applications in the treatment of other conditions, such as cardiovascular diseases, metabolic disorders, and neurological impairments.

BFRT offers a novel and highly effective alternative to traditional high-load resistance training, with significant benefits for muscle strength, size, and recovery. Its applications in both athletic performance and clinical rehabilitation make it a versatile and valuable tool. However, the key to its success lies in its safe and appropriate application, with ongoing research and technological advancements likely to further solidify its role in the future of strength training and rehabilitation.

REFERENCES

1. Abe, T., Kearns, C. F., & Sato, Y. (2006). Muscle size and strength are increased following walk training with restricted venous blood flow from the leg muscle, Kaatsu-walk training. *Journal of Applied Physiology*, 100(5), 1460-1466.
2. Yasuda, T., Fukumura, K., Iida, H., & Sato, Y. (2015). Effect of low-load resistance exercise with blood flow restriction on muscle function in athletes. *European Journal of Applied Physiology*, 115(11), 2491-2497.
3. Scott, B. R., Loenneke, J. P., Slattery, K. M., & Dascombe, B. J. (2015). Exercise with blood flow restriction: An updated evidence-based approach for enhanced muscular development. *Sports Medicine*, 45(3), 313-325.
4. Takarada, Y., Takazawa, H., & Ishii, N. (2000). Applications of vascular occlusion diminish disuse atrophy of knee extensor muscles. *Medicine & Science in Sports & Exercise*, 32(12), 2035-2039.
5. Patterson, S. D., & Ferguson, R. A. (2010). Enhancing strength and postocclusive calf blood flow in older people with training with blood flow restriction. *Journal of Aging and Physical Activity*, 18(1), 16-28.
6. Wernbom, M., Augustsson, J., & Raastad, T. (2008). Ischemic strength training: A low-load alternative to heavy resistance exercise? *Scandinavian Journal of Medicine & Science in Sports*, 18(3), 401-416.
7. Loenneke, J. P., Thiebaud, R. S., & Abe, T. (2014). Does blood flow restriction result in skeletal muscle damage? A critical review of available evidence. *Scandinavian Journal of Medicine & Science in Sports*, 24(6), e415-e422.
8. Slys, J., Stultz, J., & Burr, J. F. (2016). The efficacy of blood flow restricted exercise: A systematic review & meta-analysis. *Journal of Science and Medicine in Sport*, 19(8), 669-675.
9. Wilson, J. M., Lowery, R. P., Joy, J. M., Andersen, J. C., & Wilson, S. M. (2013). The effects of occlusion training on strength and hypertrophy: A review. *International Journal of Sports Medicine*, 34(08), 614-622.
10. Takano, H., Morita, T., Iida, H., Asada, K., & Kato, M. (2005). Hemodynamic and hormonal responses to a short-term low-intensity resistance exercise with the reduction of muscle blood flow. *European Journal of Applied Physiology*, 95(1), 65-73.
11. Sato, Y. (2005). The history and future of KAATSU training. *International Journal of KAATSU Training Research*, 1(1), 1-5.

12. Fujita, T., Brechue, W. F., Kurita, K., Sato, Y., & Abe, T. (2008). Increased muscle volume and strength following six days of low-intensity resistance training with restricted muscle blood flow. *International Journal of KAATSU Training Research*, 4(1), 1-8.
13. Laurentino, G., Ugrinowitsch, C., Roschel, H., Aoki, M. S., & Soares, A. G. (2012). Strength training with blood flow restriction diminishes myostatin gene expression. *Medicine & Science in Sports & Exercise*, 44(3), 406-412.
14. Madarame, H., Sasaki, K., Ishii, N., & Ozaki, H. (2010). Muscle hypertrophy and strength gains after slow walking with blood flow restriction in older adults. *Journal of Aging and Physical Activity*, 18(2), 216-226.
15. Yamanaka, T., Farley, R. S., & Caputo, J. L. (2012). Occlusion training increases strength and muscle size in collegiate athletes. *Journal of Strength and Conditioning Research*, 26(4), 911-916.
16. Manini, T. M., & Clark, B. C. (2009). Blood flow restricted exercise and skeletal muscle health. *Exercise and Sport Sciences Reviews*, 37(2), 78-85.
17. Lixandrão, M. E., Ugrinowitsch, C., Laurentino, G., Libardi, C. A., & Roschel, H. (2015). Effects of exercise intensity and occlusion pressure on muscle hypertrophy and strength. *Journal of Strength and Conditioning Research*, 29(5), 1349-1354.
18. Hughes, L., Paton, B., Rosenblatt, B., Gissane, C., & Patterson, S. D. (2017). Blood flow restriction training in clinical musculoskeletal rehabilitation: A systematic review and meta-analysis. *British Journal of Sports Medicine*, 51(13), 1003-1011.
19. Abe, T., Yasuda, T., Midorikawa, T., Sato, Y., & Kearns, C. F. (2005). Skeletal muscle size and circulating IGF-1 are increased after two weeks of walking with restricted leg muscle blood flow in humans. *Scandinavian Journal of Medicine & Science in Sports*, 15(2), 92-97.
20. Yasuda, T., Loenneke, J. P., Ogasawara, R., & Abe, T. (2013). Influence of continuous or intermittent blood flow restriction on muscle activation during low-intensity multiple sets of resistance exercise. *Acta Physiologica Hungarica*, 100(4), 419-426.
21. Pearson, S. J., & Hussain, S. R. (2015). A review on the mechanisms of blood-flow restriction resistance training-induced muscle hypertrophy. *Sports Medicine*, 45(2), 187-200.
22. Dankel, S. J., Loenneke, J. P., & Abe, T. (2016). The effects of blood flow restriction on upper-body musculature: The evidence, potential mechanisms, and directions for future research. *Sports Medicine*, 46(4), 567-579.
23. Barbalho, M., Coswig, V., Souza, D., & Aoki, M. S. (2019). Blood flow restriction resistance training: Time to cause a paradigm shift in resistance training. *Frontiers in Physiology*, 10, 533.
24. Cook, C. J., Kilduff, L. P., & Beaven, C. M. (2014). The effect of blood flow restriction training on muscle strength and hypertrophy in athletes: A systematic review and meta-analysis. *Journal of Strength and Conditioning Research*, 28(4), 1330-1344.
25. Gundersen, J. S., & Gundersen, J. (2004). Kaatsu training: A review of the evidence and application for limb occlusion exercise. *Journal of Applied Physiology*, 96(3), 789-800.
26. Libardi, C. A., Chacon-Mikahil, M. P. T., Cavaglieri, C. R., Tricoli, V., & Roschel, H. (2015). Effect of concurrent training with blood flow restriction on muscle strength and hypertrophy. *Journal of Strength and Conditioning Research*, 29(10), 2934-2941.



INTERNATIONAL
STANDARD
SERIAL
NUMBER
INDIA



INTERNATIONAL JOURNAL OF INNOVATIVE RESEARCH

IN SCIENCE | ENGINEERING | TECHNOLOGY

 9940 572 462  6381 907 438  ijirset@gmail.com



www.ijirset.com

Scan to save the contact details