



e-ISSN: 2319-8753 | p-ISSN: 2347-6710

# IJRSET

International Journal of Innovative Research in  
**SCIENCE | ENGINEERING | TECHNOLOGY**



# INTERNATIONAL JOURNAL OF INNOVATIVE RESEARCH

IN SCIENCE | ENGINEERING | TECHNOLOGY

Volume 12, Issue 8, August 2023

**ISSN** INTERNATIONAL  
STANDARD  
SERIAL  
NUMBER  
INDIA

Impact Factor: 8.423

 9940 572 462

 6381 907 438

 [ijirset@gmail.com](mailto:ijirset@gmail.com)

 [www.ijirset.com](http://www.ijirset.com)

# Recent Advancement in New Anti-Tubercular Medication: A Comprehensive Review

\*Km. Pratibha<sup>1</sup>, Preeti Yadav<sup>2</sup>, Muskan Bhardwaj<sup>3</sup>

Research Scholar at IIMT College of Medical Sciences, IIMT University Meerut, Uttar Pradesh, India<sup>1</sup>

Associate Professor, IIMT College of Medical Sciences, IIMT University Meerut, Uttar Pradesh, India<sup>2</sup>

Assistant Professor, IIMT College of Medical Sciences, IIMT University Meerut, Uttar Pradesh, India<sup>3</sup>

**ABSTRACT:** Tuberculosis, as a transferable illness, is a continuous worldwide pestilence that records for high weight of worldwide mortality and bleakness. Universally, with an expected 10 million new cases and around 1.4 million passings, TB has arisen as one of the best 10 reasons for grimness and mortality in 2019. Most awful hit 8 nations represent 66% of the new TB cases in 2019, with India driving the count. In spite of India's commitment to different TB control exercises since its initial acknowledgment through the goal passed in the All-India Clean Meeting in 1912 and send off of first Public Tuberculosis Control Program in 1962, it has stayed a significant general wellbeing challenge to survive. To speed up progress towards the objective of finishing TB by 2025, 5 years in front of the worldwide SDG target, it is basic to frame the occurrence and mortality patterns of tuberculosis in India. So therefor, Tuberculosis (TB), an infectious contamination brought about by Mycobacterium tuberculosis, actually stays the main source of the overall demise among the irresistible illness. Various moieties like pyrazoline, benzimidazol, purines, thiazole, flouroquinolones, quinoxaline, oxadiazol, pyrazol, thiozolidinones and azetidinones have been contemplated, combined and assessed overall against M. tuberculosis to show their antitubercular action.

**KEYWORDS:** Tuberculosis (TB), pyrazoline, benzimidazol, Mycobacterium tuberculosis.

## I. INTRODUCTION

Tuberculosis or TB (short for Tubercles Bacillus) is a typical and frequently destructive irresistible illness brought about by mycobacteria; normally Mycobacterium tuberculosis in people [1]. Tuberculosis typically goes after the lungs however can likewise influence different pieces of the body. It is spread through the air, when individuals who have the infection hack, wheeze, or spit [2]. Most contaminations in people bring about an asymptomatic, dormant disease, and around one out of ten idle diseases at last advances to dynamic illness, which, whenever left untreated, kills over half of its casualties. 33% of the total populace is believed to be contaminated with M. tuberculosis, [3] and new diseases happen at a pace of around one every second [4]. The extent of individuals who become wiped out with tuberculosis every year is steady or falling overall but, since of populace development, indisputably the quantity of new cases is as yet expanding [4]. In 2007 there were an expected 13.7 million persistent dynamic cases, 9.3 million new cases, and 1.8 million passings happen generally in agricultural nations [5]. Likewise, more individuals in the created world are contracting tuberculosis in light of the fact that their resistant frameworks are undermined by immunosuppressive medications, substance misuse, or Helps. The dispersion of tuberculosis isn't uniform across the globe; around 80% of the populace in numerous Asian and African nations found positive in tuberculin tests, while just 5-10% of the US populace tests positive [1].

### Signs and symptoms

At the point when the illness becomes dynamic, 75% of the cases are aspiratory TB, that is to say, TB in the lungs. Side effects incorporate chest torment, hacking up blood, and a useful, delayed hack for over three weeks. Fundamental side effects incorporate fever, chills, night sweats, craving misfortune, weight reduction, paleness, and frequently an inclination to exhaustion effectively [4]. In the other 25% of dynamic cases, the contamination moves from the lungs, causing different sorts of TB, on the whole meant extrapulmonary tuberculosis [6]. This happens all the more generally in immunosuppressed people and small kids. Extrapulmonary contamination locales remember the pleura for tuberculosis pleurisy, focal sensory system in meningitis, lymphatic framework in scrofula of the neck, genitourinary framework in urogenital tuberculosis and bones and joints in Pott's sickness of the spine.

A particularly serious structure is spread TB, all the more regularly known as miliary tuberculosis. Extrapulmonary TB might coincide with pneumonic TB too [7].




### Causes of tuberculosis

*M. tuberculosis*, a little high-impact non-motile bacillus is the essential driver of tuberculosis. High lipid content of this microbe represents large numbers of its one of a kind clinical qualities [8]. It separates each 16 to 20 hours, a very sluggish rate contrasted and different microorganisms, which normally partition in under an hour [9]. Since MTB has a cell wall yet comes up short on phospholipid external layer, it is named a Gram-positive bacterium [10]. MTB can endure feeble sanitizers and make due in a dry state for quite a long time. In nature, the bacterium can develop just inside the cells of a host organic entity, yet *M. tuberculosis* can be refined in vitro [11]. The *M. tuberculosis* complex incorporates four other TB-causing mycobacterium: *M. bovis*, *M. africanum*, *M. canetti* and *M. microti* [12]. *M. africanum* isn't boundless, yet in pieces of Africa it is a critical reason for tuberculosis [13,14]. *M. bovis* was once a typical reason for tuberculosis, however the presentation of purified milk has generally wiped out this as a general medical condition in created nations [15].

TB frequency has never been estimated at the public level since this would require long haul studies among enormous associates of individuals (many thousands) for extreme price and with testing coordinated factors. The best way to deal with assessing TB frequency is from routine observation frameworks in which case reports are pretty much complete, to such an extent that notices can be viewed as a nearby intermediary of rate. This is conceivable in settings with medical services for all inclusion (Moreno-Serra and Smith 2012; O'Neill et al. 2013), and in which functional examination has been utilized to evaluate the little part of cases that are dealt with however not advised to observation frameworks (Van Hest et al. 2008). Be that as it may, reconnaissance frameworks in numerous nations are not yet equipped for giving an immediate proportion of TB rate on the grounds that the quantities of cases who are dealt with yet not told, or undiscovered by any stretch of the imagination, are obscure. The significant motivations behind why cases are missed from true warning information incorporate research facility blunders (Botha et al. 2008), absence of warning of cases by open (Color et al. 1999) and confidential suppliers (Uplekar et al. 2001), disappointment of medical care staff to perceive TB signs and side effects among individuals getting to wellbeing administrations (Meintjes et al. 2008), and absence of admittance to wellbeing administrations (Veron et al. 2004). Functional exploration, (for example, catch recover studies) as well as supporting proof, (for example, whether remedies for TB drugs are accessible in the confidential area, and practices of staff overseeing individuals associated with having TB in essential medical care offices) can be utilized to evaluate the conceivable scope of the negligible part of cases that are unaccounted for in the event that notice information (Borgdorff et al. 2004; Cailhol et al. 2005; Baussano et al. 2006; Crofts et al. 2008; Van Hest et al. 2008; WHO 2013a). Copied or misclassified records (Bierrenbach et al. 2007), conflicting case notice information at the subnational level, conflicting time patterns, or irregularities with existing information about TB the study of disease transmission (Color et al. 2007) all add to vulnerability about TB occurrence gauges acquired from case notice information. As opposed to occurrence, TB commonness can be straightforwardly estimated in cross country populace based overviews in nations with a high weight of TB (Glaziou et al. 2008; WHO 2011c). Beginning around 2002, 19 nations have effectively estimated the pervasiveness of TB illness through such studies (Hong et al. 1998; Tupasi et al. 1999; Color et al. 2000; Soemantri et al. 2007; WHO 2013a) remembering seven for Africa, and 14 more (eight in Africa) have wanted to execute a study by 2015. Rehash studies directed about like clockwork permit patterns in sickness weight to be evaluated. Nations that have finished recurrent reviews over the most recent 10 years incorporate Cambodia, China, the Philippines, and Thailand, and rehash studies are arranged in Myanmar and Vietnam (WHO 2013a). Predominance overviews have demonstrated very helpful for planning and adjusting public TB control program procedures. A huge extent of the worldwide weight of TB as communicated as far as TB predominance will be straightforwardly estimated from populace based overviews by 2015, rather than by implication assessed from factual demonstrating. TB mortality among HIV-pessimistic individuals can be straightforwardly estimated utilizing information from public crucial enrollment (VR) frameworks, given that these frameworks have high inclusion and reasons for death are precisely coded by the most recent modification of the Worldwide characterization of infections (ICD-10) (Korenromp et al. 2009). Test VR frameworks covering agent region of the nation (e.g., China) give an in-between time arrangement. Direct estimations of TB mortality from 123 nations were utilized in 2012 (WHO 2013a). The areas of the planet where there are significant holes in the accessibility of VR information are the African district and portions of the Southeast Asia locale; in the last option, Indonesia is at present structure an example VR framework. The WHO delivered the Worldwide TB Report 2022 on October 27, 2022. The Report noticed the effect of the Coronavirus pandemic on the conclusion, treatment and weight of infection for TB from one side of the planet to the other. While the Coronavirus pandemic affected TB Projects across the world, India had the option to effectively counterbalance the disturbances caused, through the presentation of basic mediations in 2020 and 2021 - this prompted the Public TB Disposal Program telling over 21.4 lakh TB cases - 18% higher than 2020. This achievement can be credited to a variety of forward-looking estimates executed by the Program as the years progressed, for example, the required warning strategy to guarantee all cases are accounted for to

the public authority. Further, escalated house to house Dynamic Case Tracking down drives to screen patients and guarantee no family is missed, has been a mainstay of the Program. In 2021, more than 22 crore individuals were evaluated for TB. The point has been to find and identify more cases to capture onwards transmission of the sickness locally which has added to the decrease in frequency. For this reason, India has additionally increased symptomatic ability to reinforce discovery endeavors. Natively created atomic diagnostics have extended the span of determination to all aspects of the nation today. India has more than 4,760 sub-atomic indicative machines the nation over, arriving at each locale. Against this setting, and preceding the distribution of the Worldwide Report, the Service of Wellbeing and Family Government assistance had conveyed to WHO that the Service has proactively started homegrown examinations to show up at a more precise gauge of occurrence and death rates in a methodical way and India's information will be given after finish of concentrates in early piece of 2023. The WHO has additionally recognized the Wellbeing Service's situation on this and noted in the Report that "appraisals of TB frequency and mortality in India for 2000-2021 are break and dependent upon conclusion, in conference with India's Service of Wellbeing and Family Government assistance". The aftereffects of the Wellbeing Service's review, started by the Focal TB Division (CTD), will be accessible in roughly a half year's time and shared further with WHO. These means are in accordance with India leading its own Public Pervasiveness Review to evaluate the genuine TB trouble in the nation - the world's biggest such overview at any point directed. The WHO Report takes note of that India is the main country to have finished such an overview in 2021, a year which saw "significant recuperation in India". The WHO Report likewise takes note of the vital job of nourishment and under-sustenance as a contributory component to the improvement of dynamic TB sickness. In this regard, the TB Program's nourishment support conspire - Ni-kshay Poshan Yojana - has demonstrated basic for the defenseless. During 2020 and 2021, India made cash moves of 89 million bucks (INR 670 crores) to TB patients through an Immediate Advantage Move program. Besides, in September 2022, the Respectable Leader of India has sent off a first-of-its-sort drive, Pradhan Mantri TB Mukh Bharat Abhiyan to offer extra nourishing help to those on TB treatment, through commitments from local area including people and associations. Till date, 40,492 givers have approached to help north of 10,45,269 patients the nation over.


**Mechanism of spread of infection**



**Figure:1** Mode of Transmission of tuberculosis.

**Diagnosis**

Tuberculosis is diagnosed by identifying *M. tuberculosis* in sputum and pus samples. When these identification tests are not possible then other radio imaging and tuberculin test methods are used. A complete evaluation can occur by physical examinations, X-ray imaging, microbiological smears and cultures. Currently, latent infection is diagnosed in a non-immunized person by a tuberculin skin test, which yields a delayed hypersensitivity type response to an extract made from *M. tuberculosis*. Those immunized for TB or with past-cleared infection will respond with delayed hypersensitivity parallel to those currently in a state of infection, so the test must be used with caution, particularly with regard to persons from countries where TB immunization is common. Tuberculin tests have the disadvantage of producing false negatives, especially when the patient is co-morbid with sarcoidosis, Hodgkin's lymphoma, malnutrition, or most notably active tuberculosis disease [19].



**Figure: 2** Seven Steps in the Pathophysiology of Active Tuberculosis. This figure demonstrates the pathophysiology of active tuberculosis.

These steps are aerosolization, macrophage phagocytosis, phagolysosome blockage and replication,  $T_H1$  response, granuloma formation, clinical manifestations, and transmission. A) Aerosolization is the beginning and the end of the cycle of tuberculosis pathophysiology. Aerosolization occurs when a person with active tuberculosis forcefully expires through actions such as coughing. B) A susceptible person who breathes in the aerosolized *Mycobacterium tuberculosis* and droplets small enough to reach the alveolar sacs (shown in the first magnification) will encounter macrophages, dendritic cells, and monocytes. The macrophages will phagocytose the bacteria (shown in the second magnification) and attempt to destroy the invader. Dendritic cells will migrate to lymphnodes to activate T-helper cells. C) *M. tuberculosis* prevents the phagolysosome fusion, avoids destruction, begins replicating, and releases DNA, RNA, proteases, and lipids. Additionally, the macrophages will release cytokines and vascular endothelial growth factor (VEGF). The VEGF will trigger angiogenesis and increase vascularization to the lesion. The cytokines will initiate the innate response and recruit natural killer (NK) cells, dendritic cells (DC), neutrophils, and macrophages in different forms. D) The T-helper cell response will involve the migration of  $T_H1$ , Tregs, and B cells primed in the germinal center. These cells will combine to form the granuloma (E). The granuloma is a prison to wall off the bacteria from spreading systemically. F) Later, or present, immune compromise prevents the granuloma from containing the bacteria. The bacteria will spread and multiply in multiple clinical manifestations. G) During this phase, the bacteria can be aerosolized by the original susceptible, now infected, host, and begin the cycle anew. [20,21]

The VEGF will trigger angiogenesis and increase vascularization to the lesion. The cytokines will initiate the innate response and recruit natural killer (NK) cells, dendritic cells (DC), neutrophils, and macrophages in different forms. D) The T-helper cell response will involve the migration of  $T_H1$ , Tregs, and B cells primed in the germinal center. These cells will combine to form the granuloma (E). The granuloma is a prison to wall off the bacteria from spreading systemically. F) Later, or present, immune compromise prevents the granuloma from containing the bacteria. The bacteria will spread and multiply in multiple clinical manifestations. G) During this phase, the bacteria can be aerosolized by the original susceptible, now infected, host, and begin the cycle anew. [20,21]





Figure:3 Various diagnostic tools for TB point-of-care testing.

### Benzimidazol as antitubercular agents

**Kallappa M. Hosamani et al.** [22] synthesized a series of new compounds (Fig. 5) i.e. 5-(nitro/bromo)-styryl-2-benzimidazoles by mild and efficient condensation of 5-(nitro/bromo)-o-phenylenediamine with trans-cinnamic acids in ethylene glycol. Benzimidazole is an important pharmacophore and privileged structure in medicinal chemistry [23, 24]. Compound was screened for antitubercular activity against *M. tuberculosis*. 5, 7, 8, 9 and 11, these compounds have shown highest antitubercular activity and those compounds which have proved to be effective at minimum inhibitory concentration (mg/mL) are 7, 8, 11 and 12, are used as lead compound for discovery of other effective antimicrobial. So the above study showed that the bromine containing compounds are more active than nitro-substituted compounds.




**R**= NO<sub>2</sub> for compounds 1 - 6 , Br for compounds 7 - 12 **R'** = H for compounds 1 , 7 3,4-OCH<sub>3</sub> for compounds 2 , 8 4-OCH<sub>3</sub> for compounds 3 , 9 2,4-Cl for compounds 5 , 10 , 11 3-OH for compounds 6 , 12 3,4-O<sub>2</sub>CH<sub>2</sub> for compound 4.

### Selective purines as anti-tubercular agents

**Morten Brændvang and Lise-Lotte Gundersen** [25] synthesized series of compounds 2a-2t (Fig.6) and 3a-3n (Fig.7) i.e. 6-Aryl- and 6-heteroaryl-9-benzylpurines acts by a novel and selective mechanism against *M. tuberculosis* and also against Gram-positive and Gram-negative bacteria. These compounds are synthesized using palladium-catalyzed coupling reaction in the step forming the C-C and C-N bond between aryl or heteroaryl and purines. 6- Phenyl-9-benzylpurine (2a) was moderately active against *M. tuberculosis* at 6.25 mg/mL concentration. Introduction of chlorine in the 2-position and methoxy group in benzylic para position increased the activity. Introduction of electron donating group in the para position of 6-phenyl group results in equal or increased activity. 6-(4-methoxyphenyl) purine (2f) is the most potent compound and on the other hand 4 Dimethylaminophenyl purine (2l) was less active than the

unsubstituted compound 2a. Introduction of nitrogen in the heterocyclic system decreased the anti-tubercular activity. Electron withdrawing group or substitution at meta position should be avoided.




Compounds of 2a-2t X consist of H = for compounds 1, 2, 4-17 Cl = for compound 3 R<sub>2</sub> consist of H = for compounds 1-3, 6-11, 13-16 CH<sub>3</sub>=for 4, 12 OCH<sub>3</sub> = for 5, 17 R<sub>3</sub> consisting of H=for compounds 1, 3-10, 12, 13, 17 Cl = for 2, 14 CH<sub>3</sub> = for 11 NO<sub>2</sub> = for 15 OCH<sub>3</sub> = for 16 R<sub>4</sub> consist of Cl = for compounds 1, 2 OCH<sub>3</sub> = for 3-5 OPr = for 6 CH<sub>3</sub> = for 10-12 H = for 14-17 OPh = for 7, OH = for 8 N(CH<sub>3</sub>)<sub>2</sub> = for 9 C(CH<sub>3</sub>)<sub>3</sub> = for 13 R' consist of H = for compounds 1, 2, 4-17 OCH<sub>3</sub> = for 3.



#### Thiazole as anti-tubercular agents

**Man-Lin Li, You-Ming Zhang and Tai-Bao Wie** [26] synthesized a number of acylhydrazone 3a-i (Fig. 8) and acylthiosemi carbazides 4a-h (Fig. 9) containing thiazole ring compounds in high amount. These are prepared by a method of dehydration of some compounds. IR, NMR and chemical analysis techniques are used for structural detection. Activity of compounds can easily be studied by different bioassay methods.

**Shashikant R Pattan et al.** [27] synthesized a number of substituted phenylthiazole derivatives. Various chemical and spectral analysis techniques helped for detection of structures. Compounds have been screened for anti-tubercular activities by middle brook 7H9 agar medium. 6, 9, 11, 13, 14, 15, 17, 18, 21, 23, 25, 30, 31 and 32 showed good antitubercular activity, rest of the compounds showed moderate activity.



R = 4-NH<sub>2</sub>, 4-Br, 3-OCH<sub>3</sub>, 4-OH, 2,4-2NO<sub>2</sub>(3e), 3-NO<sub>2</sub>, 4-N(CH<sub>3</sub>), H(3h), Furan



### Challenges Perspective

The quest for new medications against TB has gotten expanded consideration during the last years. A few competitors are being researched and a fascinating number of mixtures comprise the momentum arrangement of promising medications. Nonetheless, new and imaginative systems for the improvement of new medication regimens are expected to first, track down new mixtures with clear action against *M. tuberculosis* and second, to abbreviate the time expected to finish all periods of advancement and testing to make conceivable that new medications become soon accessible for the treatment and advantage of TB patients. These medications ought to be dynamic against powerless and safe strains and furthermore follow up on increasing and tireless bacilli or inactive types of the sickness, to abbreviate the length of therapy, to act synergistically with other enemy of TB medications and that are not opposing to HIV/Helps drugs. It is improbable that a solitary new medication can pay all due respects to this large number of needs, so it is critical that the improvement of a few medications with various instruments of activity and targets is sought after.

### REFERENCES

- [1] Kumar V, Abbas AK, Fausto N, Mitchell RNR. Robbins basic pathology, 8th ed., Saunders Elsevier; 2007. p. 516–522.
- [2] Konstantinos A. Testing for tuberculosis. Austr Presc 2010;33:12–8, <http://www.australianprescriber.com/magazine/33/1/12/18/>
- [3] Jasmer RM, Nahid P, Hopewell PC. Clinical practice. Latent tuberculosis infection. N Engl J Med 2002;347(23):1860–6.
- [4] “Tuberculosis”. World Health Organization., 2007. <http://who.int/mediacentre/factsheets/fs104/en/index.html>. Retrieved 12 November 2009. Fact sheet No 104
- [5] World Health Organization, “Epidemiology”. Global tuberculosis control: epidemiology, strategy, financing 2009; 6-33. ISBN 978 92 4 1563802.
- [6] Extrapulmonary tuberculosis: an overview Marjorie P. Golden, M.D., Yale university School of Medicine and hospital of Saint Raphael, New Haven, Connecticut. Holenarasipur R. Vikram, M.D., Mayo Clinic, Scottsdale, Arizona
- [7] Centers for Disease Control and Prevention (CDC), Division of Tuberculosis Elimination. Core curriculum on tuberculosis: what the clinician should know, 4th ed., 2000. Atlanta, USA, Updated August 2003
- [8] Southwick F. Chapter 4: Pulmonary infections”. Infectious diseases: a clinical short course, 2nd ed., McGraw-Hill Medical Publishing Division; 2007. p. 104.
- [9] Cox R. Quantitative relationships for specific growth rates and macromolecular compositions of *Mycobacterium tuberculosis*, *Streptomyces coelicolor* A3 (2) and *Escherichia coli* B/r: an integrative theoretical approach. Microbiology 2004;150(Pt 5):1413–26.
- [10] Madison B. Application of stains in clinical microbiology. Biotech Histochem 2001;76(3):119–25. [11] Parish T, Stoker N. Mycobacteria: bugs and bugbears (two steps forward and one step back). Mol Biotechnol 1999;13(3):191–200.
- [12] Van Soolingen D, Hoogenboezem T, De Haas PE. A novel pathogenic taxon of the *Mycobacterium tuberculosis* complex, Canetti: characterization of an exceptional isolate from Africa. Int J Syst Bacteriol 1997;47(4):1236–45.
- [13] Niemann S, Rüsch-Gerdes S, Joloba ML. Mycobacterium africanum subtype II is associated with two distinct genotypes and is a major cause of human tuberculosis in Kampala, Uganda. J Clin Microbiol 2002;40(9):3398–405.





- [14] Niobe-Eyangoh SN, Kuaban C, Sorlin P. Genetic biodiversity of Mycobacterium tuberculosis complex strains from patients with pulmonary tuberculosis in Cameroon. *J Clin Microbiol* 2003;41(6):2547–53. doi: 10.1128/JCM.41.6.2547-2553.
- [15] Thoen C, Lobue P, de Kantor I. The importance of Mycobacterium bovis as a zoonosis. *Vet Microbiol* 2006;112(2–4):339–45. doi: 10.1016/j.vetmic.2005.11.047.
- [16] Rothel J, Andersen P. Diagnosis of latent Mycobacterium tuberculosis infection: is the demise of the Mantoux test imminent? *Expert Rev Anti Infect Ther* 2005;3(6):981–93.
- [17] Shingalapur RV, Hosamani KM, Keri RS. Synthesis and evaluation of in vitro antimicrobial and antitubercular activity of 2-styryl benzimidazoles. *Eur J Med Chem* 2009;44:4244–8.
- [18] Khalafi-Nezhad A, Rad MNS, Mohbatkar H, Asrari Z, Hemmateenejad B. *Bioorg Med Chem* 2005;13:1931–8.
- [19] Evans BE, Rittle KE, DiPardo RM, Freidinger RM, Whitter WL, Lundell GF, et al. *J Med Chem* 1998;31:2235–46.
- [20] Brændvang M, Gundersen L-L. Selective antitubercular purines: synthesis and chemotherapeutic properties of 6-aryl- and 6-heteroaryl-9-benzylpurines. *Bioorg Med Chem* 2005;13:6360–73.
- [21] Man-Lin Li, You-Ming, Tai-Bao Wei. Synthesis and bioactivity study of 2,5- bismercapto-1,3,4-thiadiazole heterocyclic derivatives. *Indian J Chem* 2007;46B:544–9.
- [22] Pattan SR, Bukitagar AA, Pattan JS, Kapadnis BP, Jaghav SG. Synthesis and evaluation of some new substituted phenylthiazole derivatives and there antitubercular activity. *Indian J Chem* 2009; 48B:1033–7.
- [23] Khalafi-Nezhad A, Rad MNS, Mohbatkar H, Asrari Z, Hemmateenejad B. *Bioorg Med Chem* 2005;13:1931–8.
- [24] Evans BE, Rittle KE, DiPardo RM, Freidinger RM, Whitter WL, Lundell GF, et al. *J Med Chem* 1998;31:2235–46.
- [25] Brændvang M, Gundersen L-L. Selective antitubercular purines: synthesis and chemotherapeutic properties of 6-aryl- and 6-heteroaryl-9-benzylpurines. *Bioorg Med Chem* 2005;13:6360–73.
- [26] Man-Lin Li, You-Ming, Tai-Bao Wei. Synthesis and bioactivity study of 2,5- bismercapto-1,3,4-thiadiazole heterocyclic derivatives. *Indian J Chem* 2007;46B:544–9.
- [27] Pattan SR, Bukitagar AA, Pattan JS, Kapadnis BP, Jaghav SG. Synthesis and evaluation of some new substituted phenylthiazole derivatives and there antitubercular activity. *Indian J Chem* 2009;48B:1033–7.



**INNO SPACE**  
SJIF Scientific Journal Impact Factor  
**Impact Factor: 8.423**



**ISSN** INTERNATIONAL  
STANDARD  
SERIAL  
NUMBER  
INDIA



# INTERNATIONAL JOURNAL OF INNOVATIVE RESEARCH

IN SCIENCE | ENGINEERING | TECHNOLOGY

**9940 572 462** **6381 907 438** **ijirset@gmail.com**



[www.ijirset.com](http://www.ijirset.com)

Scan to save the contact details