



e-ISSN: 2319-8753 | p-ISSN: 2347-6710

# IJRSET

International Journal of Innovative Research in  
**SCIENCE | ENGINEERING | TECHNOLOGY**

# INTERNATIONAL JOURNAL OF INNOVATIVE RESEARCH

IN SCIENCE | ENGINEERING | TECHNOLOGY

Volume 12, Issue 13, April 2023

**ISSN** INTERNATIONAL  
STANDARD  
SERIAL  
NUMBER  
INDIA

9940 572 462

6381 907 438

[ijirset@gmail.com](mailto:ijirset@gmail.com)

[www.ijirset.com](http://www.ijirset.com)

# Therapeutics Role and Active Constituents of *Ficus Carica* : A Review

P Selvakumar<sup>1</sup>, Anantha Valli V<sup>2</sup>, Ann Praveena S<sup>3</sup>, Boomika S<sup>4</sup>, Swetha T<sup>5</sup>

Associate Professor, Dhaanish Ahmed Institute of Technology, Coimbatore, Tamilnadu, India<sup>1</sup>

U.G Student, Department of Food Technology, Dhaanish Ahmed Institute of Technology, Coimbatore,

Tamilnadu, India<sup>2,3,4,5</sup>

**ABSTRACT:** *Ficus carica* (Moraceae) is a deciduous tree, which grows in tropical and subtropical region of India, commonly known as fig tree. Dried figs are nutritionally rich fruits. *Ficus carica* Linn. (Moraceae) is commonly known as Angir is a middle sized laticiferous deciduous tree, widely distributed in all tropical and sub-tropical countries. The fruit extracts possessed activity in anaemia, latex as anthelmintic (due to ficin) and anticarcinogenic. Traditionally, the plant is being used as purgative, aphrodisiac, anti-inflammatory, expectorant, diuretic, anti-anxiety (mild sedative). Pharmacological studies carried out on the fresh plant materials, crude extracts, and isolated components of *Ficus carica* provide an experimental support for its numerous traditional uses. However, the potent bioactive secondary metabolite for anticancer, haemostatic effect, antifungal activity, scavenging effect, and irritant potential is described by earlier researchers of this field. The present review is therefore, an effort to give a detailed survey of the literature on its pharmacognosy, phytochemistry, and pharmacological properties. Figs are one of the highest source of calcium, copper, magnesium. The seeds are real fruit in figs. In traditional medicine the roots are used in the treatment of leucoderma and ringworms. Fruits have antipyretic aphrodisiac property. The present review in therefore, an effort to give a detailed survey of literature on its pharmacognostic, traditional and pharmacological uses..

**KEYWORDS:** *Ficus Carica* Linn., Pharmacognosy, Phytochemistry, Pharmacological properties.

## I. INTRODUCTION

*Ficus* (Moraceae) comprises one of the largest genera of angiosperms with more than 800 species of trees, shrubs, hemiepiphytes, climbers, and creepers in the tropics and subtropics worldwide. This genus is an important genetic resource due to its high economic and nutritional values and also an important part of the biodiversity in the rainforest ecosystem [1-4]. It is also a good source of food for fruit-eating animals in tropical areas. The genus is divided into six subgenera based on preliminary morphology. The monoecious subgenus *Urostigma* is the largest with about 280 species all inclusive, and most of them display distinctive hemiepiphytic habits [5-7]. *Ficus* includes 23 species of hemiepiphytes and lithophytes which produce aerial and creeping root systems. The fig is an important harvest worldwide for its dry and fresh consumption. Its common edible part is the fruit which is fleshy, hollow, and receptacle. Figs are an excellent source of phenolic compounds, such as proanthocyanidins, whereas red wine and tea, which are two good sources of phenolic compounds, contain phenols lower than those in fig [8-10]. Its fruit, root, and leaves are used in traditional medicine to treat various ailments such as gastrointestinal (colic, indigestion, loss of appetite, and diarrhea), respiratory (sore throats, coughs, and bronchial problems), and cardiovascular disorders and as anti-inflammatory and antispasmodic remedy [11-13].

### Scientific Classification

Kingdom : Plantae  
Clade : Tracheophytes  
Clade : Angiosperms  
Clade : Eudicots  
Clade : Rosids  
Order : Rosales  
Family : Moraceae  
Genus : *Ficus*  
Subgenus : *F. subg. Ficus*  
Species : *F. carica*

## II. RESEARCH REVIEW

### PHYTOCHEMICAL PROPERTIES

Phytochemicals are the chemicals produced by plants. Literature survey indicated the presence of coumarins, flavonoids, sterols, triterpenoids, anthocyanins etc, in various parts of the plant. Dried seeds contain fixed oil containing the fatty acids viz oleic acid, linoleic acid, linolenic acid, palmitic acid, stearic acid, arachidic acid. caoutchouc, resin, albumin, cerin, sugar and malic acid, rennin, proteolytic enzymes, diastase, esterase, lipase, catalase, and peroxidase. Fruits contain cyanidin-3-O-glucoside, cyanidin-3-Orhamnoglucoside, saturated fat, cholesterol, sodium, insoluble sugars, protein, vitamin A, vitamin C, calcium, iron[14-16].

### PHARMACOLOGICAL REPORTS

#### Antipyretic Activity

The significant antipyretic effect of an ethanol extract of *Ficus carica* was demonstrated in a study, where this extract was effective at dose of 100, 200, and 300 mg/kg in reducing normal body temperature. Furthermore, the effect extended up to 5 h after drug administration when compared with that of paracetamol (150 mg/kg.), a standard antipyretic agent[17-18].

#### Anti-inflammatory Activity

The probable anti-inflammatory effect of petroleum ether (PEE), chloroform (CE) and ethanol (EE) extracts obtained from the leaves of *Ficus carica* Linn.. Antiinflammatory activity was studied by carrageenan-induced rat paw edema and cotton pellet granuloma methods. The petroleum ether (PEE), chloroform (CE) and ethanol (EE) extracts significantly reduced carrageenan-induced paw edema and cotton pellet granuloma in rats. These extracts showed a greater anti-inflammatory effect comparative to standard drug Indomethacin[19-20].

#### Antispasmodic and antiplatelet Activity

The aqueous ethanol extract (AEE) of *Ficus carica* fruit was studied for antispasmodic effect on rabbit jejunum preparations and for antiplatelet effect using ex vivo model of human platelets. This study exhibits the remarkable spasmolytic property in the ripe dried fruit of *Ficus carica* along with antiplatelet activity that provides sound pharmacological basis for its medicinal use in the gut motility and inflammatory disorders[21-22].

#### Antihelmintic Activity

As per WHO, only a few drugs are frequently used in the treatment of helminthes in human beings. Antihelmintics from the natural sources may play a key role in the treatment of parasite infections. This type of activity is also reported in different members of *Ficus*, i.e., *Ficus benghalensis* Linn and *Ficus racemosa* Linn[23-24].

#### Hepatoprotective Activity

The methanol extract of the leaves of *Ficus carica* was evaluated for hepatoprotective activity in CCl<sub>4</sub>-induced liver damage in a rat model. Hepatoprotective activity was also reported in the leaf extracts of *Ficus racemosa* Linn and *Ficus hispida* Linn. possess significant hepatoprotective activity against carbon tetrachloride- and paracetamol-induced hepatotoxicity in rats, respectively[25-27].

#### Hypoglycemic Activity

The hypoglycemic effect of an aqueous extract of leaves has been demonstrated in streptozotocin-induced diabetic rats, where weight loss was prevented in these animals. Additionally, treatment resulted in an increase in the survival index that correlated with increased plasma insulin levels. A similar activity has also been reported for the fruit of *Ficus carica*[28-29].

#### Hypocholesterolemic Activity

The leaves of fig have hypocholesterolemic activity. Chloroform extract is prepared from the aqueous decoction of fig leaves. It causes decline in the levels of total cholesterol and decrease in the total cholesterol/HDL cholesterol ratio, together with a reduction of the hyperglycemia. In addition to this, the cell content of cholesterol in HepG2 cells appreciates the reduction of blood cholesterol level in streptozocin-induced diabetic rats[30].

#### Anticancer activity

Anticancer activity reported a mixture of 6-O-acyl- $\beta$ -D-glucosyl- $\beta$ -sitosterols (where acyl moiety being primarily palmitoyl and linoleyl with minor amount of stearyl and oleyl) found in *Ficus carica* Linn. resin as potent cytotoxic agents. Both the natural and the synthetic compounds showed in vitro inhibitory effects on proliferation of various cancer cell lines[31].

#### Antimicrobial Activity

The antimicrobial activity of methanol (MeOH) extract of figs against oral bacteria. The MeOH extract (MICs, 0.156 to 5 mg/ml; MBCs, 0.313 to 5 mg/ml) showed a strong antibacterial activity against oral bacteria. The combination effects of MeOH extract with ampicillin or gentamicin were synergistic against oral bacteria[32-33].

#### Cytotoxicity Activity

The approximate  $IC_{50}$  values of ethanol, ethyl acetate, and dichloromethane extracts of the leaves and fruits are 10, 13, 12  $\mu$ g/ml and 12, 12, 11.5  $\mu$ g/ml, respectively, and for latex 17  $\mu$ g/ml. Thus, fig containing active principles may be responsible for cytotoxicity[34-35].

#### Antimutagenic Activity

Antimutagenic action of plant extracts of *Armoracia rusticana*, *Ficus carica*, *Zea mays* and their mixture on environmental xenobiotics has been investigated[36-37]. The plant extracts and their mixture decreased the level of mutations induced by N-metil-N'-nitro-N-nitrozoguanidin (MNNG) in *Vicia faba* cells, chlorophyll mutations in *Arabidopsis thaliana*, and NaF-induced mutability in rat marrow cells. The plant extracts and their mixture demonstrate the ability to decrease the genotoxicity of environmental mutagens[38-40].

#### Anti-angiogenic activity

The anti-angiogenic and anti-proliferative potentials of *Ficus carica* latex extract using human umbilical vein endothelial cells (HUVECs). The results clearly indicated that latex extracts of *Ficus carica* contain strong anti-angiogenic and anti-proliferative activities. Therefore, latex extract could be an ideal candidate as a potential agent for the prevention of angiogenesis in cancer and other chronic disorders[41-45].

#### Antioxidant activity

Phytochemical analysis reported significant antioxidant activity in dried fruits of *Ficus carica*. Dried figs are in vitro antioxidants after human consumption. These findings suggest that dried fruits should be a greater part of the diet as they are dense in phenol antioxidants and nutrients most probably fiber[46-48].

#### Free radical scavenging Activity

The method to study the ultrasonic assisted extraction of total flavonoids from the fruit of *Ficus carica* and their scavenging activities against hydroxyl and superoxide anion free radicals. The total flavonoid extract from the leaves had marked scavenging effects on both hydroxyl and superoxide anion free radicals in a concentration-dependent fashion[49-50].

### III. CONCLUSION

The majority of the pharmacological studies that have been carried out on *Ficus carica* were conducted using crude extracts. Thus, it is difficult to reproduce the outcomes of these studies and to locate the precise bioactive metabolite. Hence, there is a need for phytochemical standardization and identification of potent bioactive candidates. Phytochemical research carried out on *Ficus carica* has led to the isolation of few classes of plant metabolites. However, the vast traditional use and proven pharmacological activities of *Ficus carica* indicate that an immense scope still exists for its phytochemical exploration. An extensive literature survey revealed that *Ficus carica* is a sacred and important medicinal plant used for the ethnomedicinal treatment of anemia, bronchitis, constipation, diabetes, fever (jaundice), hemorrhoids, inflammation, liver disorders, infectious diseases, and many more throughout the world. Pharmacological studies carried out on the fresh plant materials, crude extracts, and isolated components of *Ficus carica* provide an experimental support for its numerous traditional uses. They failed to recognize the referred biological activities. Therefore, there is a vast scope for establishing a relation between given phytoconstituents and biological activities. The outcome of the future research in the above-mentioned areas will provide a convincing support for the future clinical uses of *Ficus carica* in modern medicine.

### REFERENCES

- [1] Husain SZ, Malik RN, Javaid M, Bibi S. (2008). Ethnobotanical properties and uses of medicinal plants of Morgah Biodiversity Park, Rawalpindi. *Pak J Bot* 40:1897–911

- [2] Ibrar M, Hussain F, Sultan A. (2007). Ethnobotanical studies on plant resources of Ranyal hills, district Shangla, Pakistan. *Pak J Bot* 39:329–37.
- [3] Idolo M, Motti R, Mazzoleni S. (2010). Ethnobotanical and phytomedicinal knowledge in a long-history protected area, the Abruzzo, Lazio and Molise National Park (Italian Apennines). *J Ethnopharmacol* 127:379–95.
- [4] Selvakumar P and Loganathan, V. Phytochemical screening and antioxidant activity of red flowered *Mirabilis jalapa* leaf in different solvents, *International Journal of Pharma and Bio Sciences*, 2012, 3(4), pp. 440–446.
- [5] Roshan Kumar, Purabi Saha, Priya Lokare, Kunal Datta, P. Selvakumar, & Anurag Chourasia. (2022). A Systemic Review of *Ocimum sanctum* (Tulsi): Morphological Characteristics, Phytoconstituents and Therapeutic Applications. *International Journal for Research in Applied Sciences and Biotechnology*, 9(2), 221–226. <https://doi.org/10.31033/ijrasb.9.2.15>.
- [6] Ishtiaq M, Hanif W, Khan MA, et al. (2007). An ethnomedicinal survey of important medicinal folklore food phytonyms of flora of Samahni Valley (Azad Kashmir) Pakistan. *Pak J Biol Sci* 10:2241–56.
- [7] İsmet U, Sinan O, Ersin K, Sezgin Ç. (2010). Ethnobotanical aspects of Kapıdağ Peninsula (Turkey). *Biol Diversity Conserv* 3:15–22.
- [8] Mukesh Kr. Singh, Ajay Kumar, Roshan Kumar, Dr. P. Satheesh Kumar, P.Selvakumar and Anurag Chourasia, Effects of Repeated Deep Frying on Refractive Index and Peroxide Value of Selected Vegetable Oils, *International Journal for Research in Applied Sciences and Biotechnology*, Volume-9, Issue-3 (May 2022).
- [9] Selvakumar P. Biomedical activities of floradol. *J Prev Diagn Treat Strat Med* 2022;1:158-62. DOI: 10.4103/jpdtm.jpdtm\_46\_22.
- [10] Jang S, Kelley KW, Johnson RW. (2008). Luteolin reduces IL-6 production in microglia by inhibiting JNK phosphorylation and activation of AP-1. *Proc Natl Acad Sci USA* 105:7534–9.
- [11] Jaradat NA. (2005). Medical plants utilized in Palestinian folk medicine for treatment of diabetes mellitus and cardiac diseases. *J Al-Aqsa Univ* 9:1–27.
- [12] P. Selvakumar and V Loganathan, In Vitro Phytochemical, Antimicrobial and Antioxidant Activity Studies on *Alocasia Sanderiana* W.Bull. *Indo American Journal of Pharmaceutical Sciences*, 2016, 3 (3), 252-264.
- [13] Solomon A, Golubowicz S, Yablowicz Z, et al. (2010). Protection of fibroblasts (NIH-3T3) against oxidative damage by cyanidin-3-rhamnoglucoside isolated from fig fruits (*Ficus carica* L.). *J Agric Food Chem* 58:6660–5.
- [14] P Selvakumar and V Loganathan, In Vitro pharmacology studies on *Alocasia Sanderiana* W.Bull. *Journal of Pharmacognosy and Phytochemistry* 2016; 5(2): 114-120.
- [15] Jeong M, Kim H, Cha J. (2009). Antimicrobial activity of methanol extract from *Ficus carica* leaves against oral bacteria. *J Bacteriol Virol* 39:97–102.
- [16] Jeong WS, Lachance PA. (2001). Phytosterol and fatty acids in fig (*Ficus carica* var. Mission) fruit and tree components. *J Food Sci* 66:278–81.
- [17] Loganathan V and Selvakumar P, In-Vitro assessment of antimicrobial and antioxidant activities of different extracts of *Mallotus rhamnifolius*. *Journal of Pharmacognosy and Phytochemistry* 2018. ; 7(6): 1995-1999.
- [18] Sruthi B, Sunny G, Hajra N, Sakthivel S. (2012). Diuretic activity of ethanolic extracts of *Ficus carica* L. fruits. *Int J Res Pharmacol Pharmacother* 1:25–8.
- [19] Loganathan V and Selvakumar P, A Study of the Physico-Chemical and Phytochemical Parameters of Leaves of *Mallotus rhamnifolius*. *International Journal of Pharmacognosy and Phytochemical Research* 2017; 9(6); 858-863. DOI number: 10.25258/phyto.v9i6.8191.
- [20] Joseph B, Raj SJ. (2011). Pharmacognostic and phytochemical properties of *Ficus carica* Linn – An overview. *Int J PharmTech Res* 3:8–12.
- [21] P.Selvakumar, In vitro antioxidant, antibacterial activities and physicochemical evaluation of ethanolic extracts of three medicinal plant leaves, *International Journal of Medicine Research*. Volume 1; Issue 1; March 2016; 01-04.
- [22] Stepek G, Buttle DJ, Duce IR, et al. (2005). Assessment of the anthelmintic effect of natural plant cysteine proteinases against the gastrointestinal nematode, *Heligmosomoides polygyrus*, in vitro. *Parasitology* 130:203–11.
- [23] P.Selvakumar and V. Loganathan, Preliminary phytochemical investigation of extract of leaves and stem of *Euphorbia hirta*, *International journal of current science* 2012, 48-51. ISSN 2250-1770.
- [24] Patil VV, Bhangale SC, Patil VR. (2010a). Evaluation of antipyretic potential of *Ficus carica* leaves. *Int J Pharm Sci Rev Res* 2:48–50.
- [25] P. Selvakumar and V. Loganathan, Preliminary phytochemical investigation of extract of leaves and stem of *Euphorbia hirta* in *International journal current science* 2012, 48-51. ISSN 2250-1770.
- [26] Singh D, Singh B, Goel RK. (2011). Traditional uses, phytochemistry and pharmacology of *Ficus religiosa*: A review. *J Ethnopharmacol* 134:565–83.
- [27] Solomon S, Golubowicz Z, Yablowicz S, et al. (2006). Antioxidant activities and anthocyanin content of fresh fruits of common fig (*Ficus carica* L.). *J Agric Food Chem* 54:7717–23.

## Organized by

Dhaanish Ahmed Institute of Technology, KG chavadi, Coimbatore, India

- [28] Khan KY, Khan MA, Ahmad M, et al. (2011). Ethno-medicinal species of genus *Ficus* L. used to treat diabetes in Pakistan. *J Appl Pharm Sci* 1:209–11.
- [29] Khan SW, Khatoun S. (2007). Ethnobotanical studies on useful trees and shrubs of Haramosh and Bugrote valleys, in Gilgit Northern areas of Pakistan. *Pak J Bot* 39:699–710.
- [30] P. Selvakumar and V. Loganathan, “In vitro antibacterial and antifungal activities of morinda tinctoria leaf in different solvents” in *Pharmacia*, vol - 2, issue - 1, June 2013 (53). ISSN 0976-9692.
- [31] Kalaskar MG, Shah DR, Raja NM, et al. (2010). Pharmacognostic and phytochemical investigation of *Ficus carica* Linn. *Ethnobot Leaflets* 14:599–609.
- [32] Rubnov S, Kashman Y, Rabinowitz R, et al. (2001). Suppressors of cancer cell proliferation from fig (*Ficus carica*) resin, isolation and structure elucidation. *J Nat Prod* 64:993–6.
- [33] Sgariberi VC, Gupte SM, Karmer DE, Whitaker JR. (1964). *Ficus* enzymes, I. Separation of the proteolytic enzymes of *Ficus carica* and *Ficus glabrata* lattices. *J Biol Chem* 239:2170–7.
- [34] P. Selvakumar and V. Loganathan, Physicochemical Analysis of *Alocasia Sanderiana* W. *Bull. International Journal of Pharmaceutical Science and Research*. V-1; Issue 3; March 2016; 01-03.
- [35] Khanom F, Kayahara H, Tadasa K. (2000). Superoxide-scavenging and prolyl endopeptidase inhibitory activities of Bangladeshi indigenous medicinal plants. *Biosci Biotech Biochem* 64:837–40.
- [36] P. SelvaKumar “Comparison of ABTS, FRAP assays for estimating antioxidant activity from *Alternanthera sessilis* extracts” in *International journal current science* 2012, 7-10. ISSN 2250-1770.
- [37] Khodarahmi GA, Ghasemi N, Hassanzadeh F, Safaie M. (2011). Cytotoxic effects of different extracts and latex of *Ficus carica* L. on HeLa cell line. *Iran J Pharm Res* 10:273–7.
- [38] Patil AP, Patil VP, Patil VR, Chaudhari RY. (2010b). Antihelmintic and preliminary phytochemical screening of leaves of *Ficus carica* Linn against intestinal helminthiasis. *Int J Res Ayurveda Pharm* 1:601–5.
- [39] Patil DR, Patil PS, Patil DA. (2011). Indigenous home remedies as applied in Shirpur Taluka of Dhule district (Maharashtra) India. *Curr Bot* 2:34–5.
- [40] Patil VV, Patil VR. (2011b). Evaluation of anti-inflammatory activity of *Ficus carica* Linn leaves. *Indian J Nat Prod Resour* 2:151–5.
- [41] Selvakumar P. Phytochemical and pharmacological profile review of *Bryophyllum pinnatum*. *Biomed Biotechnol Res J* 2022;6:295-301.
- [42] Krishna MG, Pallavi E, Ravi KB, et al. (2007). Hepatoprotective activity of *Ficus carica* Linn. leaf extract against carbon tetrachloride-induced hepatotoxicity in rats. *DARU* 15:162–7.
- [43] Shiraishi SC, Kawakami K, Widodo SE, et al. (1996). Organic acid profile in the juice of fig fruits. *J Fac Agric, Kyushu Univ* 41:29–33.
- [44] Oliveira AP, Silva LR, Andrade PB, et al. (2010). Further insight into the latex metabolite profile of *Ficus carica*. *J Agric Food Chem* 58:10855–63.
- [45] P.Selvakumar, Pharmacognostical and phytopharmacological review on coatbuttons. *J PrevDiagn Treat Strategies Med* 2022;1:202-8.
- [46] Akkala Abhilasha, Anthati Sreenivasulu, T Manimozhi, P Satheesh Kumar, P Selvakumar, Prabhdeep Singh, The Model of Smart Sensing Device For Sensitive Nanoclusters Modification in Sensing Properties, 2022 2nd International Conference on Advance Computing and Innovative Technologies in Engineering (ICACITE), IEEE, 1043-1047
- [47] Mavlonov GT, Ubaidullaeva KA, Rakhmanov MI, et al. (2008). Chitin-binding antifungal protein from *Ficus carica* latex. *Chem Nat Compd* 44:216–19.
- [48] P Selvakumar, V Loganathan, Phytochemical Screening and In vitro Antioxidant Activity of Male and Female Flowers Ethanolic Extracts of *Cocos nucifera*, *international journal of life science*, 2016, 5, 3, pp-127-131.
- [49] Selvakumar P, Fiber Reinforced Polymer Composites: Classification, Chemical Treatment, Mechanical and Tribological Properties, *Research Journal of Chemistry and Environment*, Vol. 27 (1) January (2023).
- [50] Oliveira AP, Valentão P, Pereira JA, et al. (2009). *Ficus carica* L., metabolic and biological screening. *Food Chem Toxicol* 47:2841–6.



# INTERNATIONAL JOURNAL OF INNOVATIVE RESEARCH

IN SCIENCE | ENGINEERING | TECHNOLOGY

 9940 572 462  6381 907 438  [ijirset@gmail.com](mailto:ijirset@gmail.com)



[www.ijirset.com](http://www.ijirset.com)

Scan to save the contact details