



e-ISSN: 2319-8753 | p-ISSN: 2347-6710

IJRSET

International Journal of Innovative Research in
SCIENCE | ENGINEERING | TECHNOLOGY

INTERNATIONAL JOURNAL OF INNOVATIVE RESEARCH

IN SCIENCE | ENGINEERING | TECHNOLOGY

Volume 12, Issue 13, April 2023

ISSN INTERNATIONAL
STANDARD
SERIAL
NUMBER
INDIA

9940 572 462

6381 907 438

ijirset@gmail.com

www.ijirset.com

Medicinal Significance and its Uses of *Musa Paradisiaca* : A Review

P Selvakumar¹, Roshni J², Priyadharshini D³, Praveena P⁴, Hilmi Faliha Y⁵

Associate Professor, Dhaanish Ahmed Institute of Technology, Coimbatore, Tamilnadu, India¹

U.G Student, Department of Food Technology, Dhaanish Ahmed Institute of Technology, Coimbatore, Tamilnadu, India^{2,3,4,5}

ABSTRACT: *Musa paradisiaca* is a plant normally named as banana which is a member of the Musaceae group. The various elements of the plant are widely used to cure various illness in humans like diabetes, diarrhoea, dysentery, hypertension, hysteria, epilepsy, leprosy, haemorrhages, renal calculi, and ulcers. The documented pharmacological advantage of this plant are antilithiatic, antioxidant, antibacterial, antidiabetic, antiulcer, antidiarrheal, hypocholesterolaemic, hepatoprotective, anti-snake venom, wound healing, hair growth promoting, antifungal, and antimenorrhagic activity. This review converse the features on predictable uses and pharmacological advantages of *Musa paradisiaca*. Phytochemical studies have reported several secondary metabolites from the plant. *Musa paradisiaca* has numerous phytochemicals and has a substantial pharmacological activity that can help to boost different health issues.

KEYWORDS: *Musa paradisiaca*, Ethno medicine, Photochemical, Bioactivity.

I. INTRODUCTION

Banana plants represent some of the largest herbaceous plants existing in the present, with some reaching up to 9 metres (30 ft) in height. The large herb is composed of a modified underground stem (rhizome), a false trunk, a network of roots, and a large flower spike[1-5]. The false trunk is an aggregation of the basal portion of leaf sheathes; it is not until the plant is ready to flower that a true stem grows up through the sheath and droops back down towards the ground. At the end of this stem grows a peduncle with many female flowers protected by large purple-red bracts[6-10]. The extension of the stem (this part called the rachis) continues growth downward where a terminal male flower grows. The leaves originate from a pseudostem and unroll to show a leaf blade with two lamina halves. *Musa paradisiaca* reproduces by both sexual (seed) and asexual (suckers) processes, utilizing asexual means when producing sterile (non-seedy) fruits[11-13]. Further qualities to distinguish *Musa paradisiaca* include spirally arranged leaves, fruits as berries, latex-producing cells present, 5 connate and 1 member of the inner whorl distinct, and petiole with one row of air channels[14-17].

Taxonomy

Kingdom	: Plantae
Subkingdom	: Tracheobionta
Super Division	: Spermatophyta
Division	: Magnoliophyta
Class	: Liliopsida
Subclass	: Zingiberidae
Order	: Zingiberales
Family	: Musaceae
Genus	: Musa
Species	: paradisiaca

Cultivated Bananas

A number of distinct groups of plants bearing edible fruit have been developed from species of *Musa paradisiaca*. In English, fruits which are sweet and used for dessert are usually called "bananas", whereas starchier varieties used for cooking are called "plantains", but these terms do not have any botanical significance[18-20]. By far the largest and now the most widely distributed group of cultivated bananas is derived from section *Musa paradisiaca*, particularly *M. acuminata* and *M. balbisiana*, either alone or in various hybrid combinations. The next but much smaller group is derived

Organized by

Dhaanish Ahmed Institute of Technology, KG chavadi, Coimbatore, India

from members of section *Musa paradisiaca* (previously classified as *AustraliMusa paradisiaca*) and is restricted in importance to Polynesia[21-22].

Morphology

The Banana plant, *Musa paradisiaca* often erroneously referred to as a "tree", is a large herb, with succulent, very juicy stem, which is a cylinder of leaf-petiole sheaths, reaching a height of 20 to 25 ft (6-7.5 m) and arising from a fleshy rhizome or corm. Leaves are tender, smooth, oblong or elliptic numbering 4 or 5 to 15, arranged spirally and they unfurl, as the plant grows, at the rate of one per week. The inflorescence, a transformed growing point, is a terminal spike shooting out from the heart in the tip of the stem. At first, it is a large, long-oval, tapering, purple-clad bud. As it opens, it is seen that the slim, nectar-rich, tubular, toothed, white flowers are clustered in whorled double rows along the floral stalk, each cluster covered by a thick, waxy, hood like bract, purple outside, deep-red within. Female flowers occupy the lower 5 to 15 rows[23-24].

Traditional uses

Many traditional uses of banana have been well documented, for example, the leaf and stem are used to treat diarrhoea; the stem is good for asthenia and wounds, and the leaf for the treatment of inflammation, headache and rheumatism. Previous studies reported that *M. Paradisiaca* had antimicrobial and healing activities. Nevertheless, only a few studies have reported on the efficacy of this plant against nematodes. Various parts of *M. Paradisiaca* have been used for various medicinal purposes. It has traditionally been used for antidepressant, antibacterial, antihypertensive, antiulcerogenic, urolithiasis laxatives, antihelmin, analgesic, antifungal, constipation, wound healing, fevers, burns, diarrhoea, inflammation, pains and antivenomic for snake bites[25-26].

II. RESEARCH REVIEW**Phytochemicals and mineral contents**

Catecholamines such as norepinephrine, serotonin, dopamine. Tryptophan, indole compounds. pectin have been found in the pulp. Several flavonoids and related compounds (Leucocyanidin, quercetin and its 3-O-galactoside, 3-O-glucoside, and 3-O-rhamnosyl glucoside) were isolated from the unripe pulp of plantain. Serotonin, nor-epinephrine, tryptophan, indole compounds, tannin, starch, iron, crystallisable and non-crystallisable sugars, vitamin C, B-vitamins, albuminoids, fats, mineral salts have been found in the fruit pulp of *M. paradisiaca* and *M. sapientum*. Carbohydrates have been isolated from *M. sapientum*[27-28].

Pharmacological Actions

The various effects of *Musa paradisiaca* Linn. are documented in traditional as well as scientific literature. The main pharmacological effects of this plant are anti-diabetic activity, antiulcer activity, antioxidant property, wound healing, hair growth promoter, diuretic, analgesic, augmenting action on skeletal muscle contraction, antihypertensive activity, anti-allergic activity and mutagenic effects and haemostatic activity in which a few are reported[29-31].

Antifungal activity

Effect of plantain *Musa paradisiaca* peel and stalk extracts were investigated for determining the antifungal action, using percentage inhibition test. The results further justify the claim that *Musa paradisiaca* stalk and peel extract demonstrated antifungal action in which methanol was seen to be a better solvent for extracting active ingredients from medicinal plants considering the high susceptibility of test organisms to methanol extract than ethanol extract[32-33].

Antidiabetic activity

Aqueous extracts and methanolic fractions of *M. paradisiaca* flowers and bract produced antidiabetic effect in male Wistar rats experimentally induced diabetes using intravenous streptozotocin (STZ). Validate the antidiabetic activity of methanolic flower extract attributable to the polyphenolic compound and dietary fibres present, thus backing calls for the exploration of inflorescence as a potential functional food and/or nutraceutical with excellent nutritional and organoleptic properties[34-35].

Antiulcer activity

Histological studies showed that banana treatment sections showed a greater aggregation and intensity of pink spots when compared to controls. This study suggests that banana powder treatment not only strengthens mucosal resistance against ulcerogens but also promotes healing by inducing cellular proliferation. The extract (50mg/kg twice daily for 5 days) showed significant antiulcer effect and antioxidant activity in gastric mucosa homogenates where it reversed the increase in ulcer index, lipid peroxidation and superoxide dismutase values induced by stress[36-37].

Antioxidant property

The antioxidant behavior of the extracts was evaluated by using the thiocyanate method, β -carotene bleaching method and 1, 1-diphenyl-2-picrylhydrazyl (DPPH) free radical elimination. Antioxidant activity of water extracts was comparable to those of synthetic antioxidants such as butylated hydroxyanisole and butylated hydroxytoluene and it shows a significant antioxidant property. The antioxidant effects of crude extracts from green banana and yellow peel were investigated and the results indicated that the extract of green peel recorded more significant activities than that of yellow peel at other solvents extracts[38-39].

Wound healing activity

The rats were given graded dose of (50-200 Kg/day) of aqueous and methanolic extract of *Musa paradisiaca sapientum* var. *paradisiaca* orally for a period of 10-21 days depending upon the type of study. Both extracts when studied for incision and dead space wounds parameters increased wound breaking strength and levels of hydroxyl proline, hexuronic acid, hexosamine, superoxide dismutase, reduced glutathione in the granulation tissue and decreased percentage of wound area, scar area when compared with the control group both the extracts showed good safety profile[40-41].

Hair growth promoting activity

For the evaluation of the hair growth promoting activity of *Musa paradisiaca paradisiaca* unripe fruit extract, the study was aimed to investigate the hair growth promoting activity of *Musa paradisiaca paradisiaca* unripe fruit extract. The extract of *Musa paradisiaca paradisiaca* unripe fruit when tested for the hair growth activity was assayed by studying hair length and microscopic study of follicles in vehicle control, 2% minoxidil treated and extracts treated animals. The findings suggest that extract of *Musa paradisiaca paradisiaca* unripe fruit has potential as a hair growth promoter[42].

Analgesic activity

The analgesic activity of aqueous extract of the plant was evaluated using the hot plate method and writhing test in mice. The hot plate method is useful in detecting centrally acting analgesics whereas acetic acid induced writhing method is useful to detect peripheral analgesic effects. As the phytochemical screening has shown the presence of carbohydrates, sterols, proteins, flavonoids, alkaloids in aqueous extract of *Musa paradisiaca paradisiaca* leaves, its potent activity may be attributed to the presence of these phytoconstituents[43].

Augmenting action on skeletal muscle contraction

Augmentation action in skeletal muscles was studied by taking an extract obtained from juice expressed from the stem of the plantain banana tree (*Musa paradisiaca sapientum* L., var. *paradisiaca*) induces twitch augmentation in skeletal muscle. The mechanism of this action was investigated in the mouse hemi-diaphragm preparation. Directly evoked twitches and potassium induced (K⁺) contractures were both increased by the extract[44].

Antidiarrheal activity

The banana cellulose (a soluble polymer) will make smoother and cease constipation. However, consumption of banana could be an advantage in individuals hurting from symptom. The investigators observed that the banana flake cluster had fewer symptoms clinically; with fifty seven of the topics were diarrhoea free on their last study day in differentiation to pure gold of the medically treated subjects. This study shows the disease curing activity of plantain tree[45].

Diuretic activity

Ash of the peel of *M. sapientum* showed an increase in urine volume and K⁺ as well as other electrolyte excretion than normal saline in a study in rats. Successive ethanolic extract also give this diuretic effect. Phytochemicals such as saponin, flavonoids and terpenoids are known to be responsible for this effect[46].

Antihypertensive activity

The antihypertensive effect of *M. paradisiaca* in albino rats was reported. Banana diet has a mean arterial blood pressure lowering as well as onset preventing effect in rats with elevated blood pressure induced by desoxycorticosterone acetate (DOCA) administration. The antihypertensive effect of ripe banana pulp in desoxycorticosterone acetate-induced hypertensive rats which may be due to the high tryptophan and carbohydrate content of banana that increases serotonin levels and gives serotonin-mediated natriuretic effect[47].

Antimenorrhagic activity

Consuming one cooked banana flower with one cup of curd or yogurt is one of the most efficient ways of treating excessive bleeding during menstruation. The cooked banana flower and curd combination increases the level of

Organized by

Dhaanish Ahmed Institute of Technology, KG chavadi, Coimbatore, India

progesterone in the body and thereby reduces bleeding associated with menorrhagia. The flowers are also taken as an infusion in normal doses for painful menstruation[48-49].

Hepatoprotective activity

Parts of *M. balbisiana* have also been reported to have hepatoprotective activity. The present study demonstrates that banana fruit and its peel have hepatoprotective, hypoglycemic effects against diabetic hepatotoxic rats. The result revealed that all cows responded well to the given treatment regimen showing therapeutic potential as hepatoprotective and strong antioxidant properties[50-51].

Anti-allergic activity

The water extract of pulp of ripe *M. sapientum* has been reported to have significant anti-allergic activity on antigen induced degranulation in RBL-2H3 cells with an IC₅₀ value of 13.5±2.4[52].

III. CONCLUSION

Medicinal plants have attracted significant world interest in past years. Banana tree could be a healthful plant with numerous pharmacological properties. The leading pharmacological outcome of this plant are antilithiatic, antioxidant, antibacterial, antidiabetic, antiulcer, antidiarrheal, hypocholesterolaemic, hepatoprotective, antsnake venom, wound healing, hair growth promoting, antifungal, and antimenorrhagic actions. Because of the therapeutic effect, there is a vast range of future research on *Musa paradisiaca*. Different parts of *M. paradisiaca* have been studied extensively and found to modulate physiological activities and antimicrobial effects on various microorganisms. However, as against most of the conventional antimicrobials used in pharmacological and cosmetics preparations, the mechanisms of actions of this potent antimicrobial and medically active plant on each of the applications have not been well studied. Similarly, while antimicrobial—generally—studies are extensive, limited antiviral and in vivo reports exist. Putatively, the bioactive metabolites that appear to mediate antimicrobial activity are stigmaterol and β-sitosterol (phytosterols) and the triterpenes trio of 31-norcyclolaudenone, 24-methylene-cycloartanol, and cycloeucalenone. The paucity of in vivo and antiviral studies are gaps that require to be earnestly. *Musa paradisiaca* has numerous photochemical and has a substantial pharmacological activity that can help to boost different health issues.

REFERENCES

- [1] Tewtrakul et al., Antiallergic and anti-microbial activities of some Thai crops. Songklanakarin J. Sci. Technol. 30 (4); 2008: 467-473.
- [2] Selvakumar P and Loganathan, V. Phytochemical screening and antioxidant activity of red flowered *Mirabilis jalapa* leaf in different solvents, International Journal of Pharma and Bio Sciences, 2012, 3(4), pp. 440–446.
- [3] Osim EE and Ibu JO. The Effect of Plantains (*Musa paradisiaca paradisiaca*) on DOCA-Induced Hypertension in Rats. Pharm. Biol. 29 (1); 1991: 9-13.
- [4] Roshan Kumar, Purabi Saha, Priya Lokare, Kunal Datta, P. Selvakumar, & Anurag Chourasia. (2022). A Systemic Review of *Ocimum sanctum* (Tulsi): Morphological Characteristics, Phytoconstituents and Therapeutic Applications. International Journal for Research in Applied Sciences and Biotechnology, 9(2), 221–226. <https://doi.org/10.31033/ijrasb.9.2.15>.
- [5] Agarwal et al., Evaluation of Wound healing activity of extracts of plantain Banana (*Musa paradisiaca sapientum* Var. *Paradisiaca*) in rats. Indian J Exp Biol., 47;2009:32–40.
- [6] Mukesh Kr. Singh, Ajay Kumar, Roshan Kumar, Dr. P. Satheesh Kumar, P. Selvakumar and Anurag Chourasia, Effects of Repeated Deep Frying on Refractive Index and Peroxide Value of Selected Vegetable Oils, International Journal for Research in Applied Sciences and Biotechnology, Volume-9, Issue-3 (May 2022).
- [7] Selvakumar P. Biomedical activities of florasol. J Prev Diagn Treat Strat Med 2022;1:158-62. DOI: 10.4103/jpdtm.jpdtm_46_22.
- [8] Wu D and Kress JW. *Musa paradisiaca* ceae. Flora of China. 24;2000:297–313
- [9] Robinson JC. Bananas and plantains. Cambridge, UK: CAB International, University Press; 1996.
- [10] Martin T.S, Ohtani K., Kasai R., Yamasaki K. A Hemiterpenoid Glucoside from *Musa paradisiaca paradisiaca*. Nat. Med. 2000; 54(4): 190-192.
- [11] P. Selvakumar and V Loganathan, In Vitro Phytochemical, Antimicrobial and Antioxidant Activity Studies on *Alocasia Sanderiana* W.Bull. Indo American Journal of Pharmaceutical Sciences, 2016, 3 (3), 252-264.
- [12] Lewis DA and Shaw GP. A natural flavonoid and synthetic analogues protect the gastric mucosa from aspirin-induced erosions. J. Nutr. Biochem. 12;2001: 95-100.

Organized by

Dhaanish Ahmed Institute of Technology, KG chavadi, Coimbatore, India

- [13] Mukhopadhyaya et al., Effect of Banana Powder (*Musa paradisiaca sapientum* var. *paradisiaca*) on Gastric Mucosal Shedding. *J. Ethnopharmacol.* 1987; 21:11-19.
- [14] Surbhi Gupta et al, Analgesic activity of aqueous extract of *Musa paradisiaca paradisiaca*. *Der Pharmacia Sinica.*, 2011, 2(4), 74.
- [15] Mokbel MS and Fumio H. Antibacterial and antioxidant activities of banana fruit peel. *American journal of Biochemistry and Biotechnology.*1(3); 2005, 125.
- [16] Andrade CUB., Perazzo FF., Maistro E.L. Mutagenicity of the *Musa paradisiaca paradisiaca* (*Musa paradisiaca* ceae) fruit peel extract in mouse peripheral blood cells in vivo. *Genet. Mol. Res.* 7(3);2008: 725-732.
- [17] P Selvakumar and V Loganathan, In Vitro pharmacology studies on *Alocasia Sanderiana*W.Bull. *Journal of Pharmacognosy and Phytochemistry* 2016; 5(2): 114-120.
- [18] Cidian ZD. Grand dictionary of Chinese traditional medicine. Shanghai: Shanghai Science and Technology Press. 2005:178.
- [19] Loganathan V and Selvakumar P, In-Vitro assessment of antimicrobial and antioxidant activities of different extracts of *Mallotusramnifolius*. *Journal of Pharmacognosy and Phytochemistry* 2018. ; 7(6): 1995-1999.
- [20] Jain et al., Study of antacid and diuretic activity of ash and extracts of *Musa paradisiaca sapientum* L. fruit peel. *Phcog. Mag.* 3(10); 2007: 116-119.
- [21] Loganathan V and Selvakumar P, A Study of the Physico-Chemical and Phytochemical Parameters of Leaves of *Mallotusramnifolius*. *International Journal of Pharmacognosy and Phytochemical Research* 2017; 9(6); 858-863. DOI number: 10.25258/phyto.v9i6.8191.
- [22] Ribeiro et al., Acute diuretic effects in conscious rats produced by some medicinal plants used in the state of Sao Paulo. *Brazil J Ethnopharmacol.*, 24;1988:19–29.
- [23] P.Selvakumar, In vitro antioxidant, antibacterial activities and physicochemical evaluation of ethanolic extracts of three medicinal plant leaves, *International Journal of Medicine Research.* Volume 1; Issue 1; March 2016; 01-04.
- [24] Pannangpetch et al., TheAntiulcerative Effect of Thai *Musa paradisiaca* Species in Rats. *Phytother. Res.* 15; 2001: 407–410.
- [25] P.Selvakumar and V. Loganathan, Preliminary phytochemical investigation of extract of leaves and stem of *Euphorbia hirta* in *International journal of current science* 2012, 48-51. ISSN 2250-1770.
- [26] Kailash P, Bharathi K and Srinivasan K. Evaluation of *Musa paradisiaca* (*Paradisiaca* Linn, Cul-tivar) “Put tubale” stem juice for antilithiatic activity in albino rats. *Indian J PhysiolPharmacol.*, 37;1993:337–41.
- [27] P. Selvakumar and V. Loganathan, “In vitro antibacterial and antifungal activities of *morindatinctoria* leaf in different solvents” in *Pharmacia*, vol - 2, issue - 1, June 2013 (53). ISSN 0976-9692.
- [28] Jachak SM and Saklani A. Challenges and opportunities in drug discovery from plants. *Curr. Sci.*, 92(9); 2007: 1251-1257.
- [29] P.Selvakumar and V. Loganathan, Physicochemical Analysis of *Alocasia Sanderiana*W.Bull. *International Journal of Pharmaceutical Science and Research.* V-1; Issue 3; March 2016; 01-03.
- [30] Newman DJ, Cragg GM and Snader KM. Natural products as sources of new drugs over the period 1981–2002. *J. Nat. Prod.*66 (7); 2003: 1022–1037.
- [31] Kennedy and Jean. “Pacific Bananas: Complex Origins, Multiple Dispersals?” *Asian Perspectives* 47(1); 2008: 75-94.
- [32] P. SelvaKumar “Comparison of ABTS, FRAP assays for estimating antioxidant activity from *Alternanthera sessilis* extracts” in *International journal current science* 2012, 7-10. ISSN 2250-1770.
- [33] Sharma et al., In vitro anthelmintic screening of indigenous medicinal plants against *Haemonchuscontortus* (Rudolphi, 1803) Cobbold, 1898 of sheep and goats. *Indian Journal of Animal Research* 5; 1971:33–38.
- [34] Selvakumar P. Phytochemical and pharmacological profile review of *Bryophyllumpinnatum*. *Biomed Biotechnol Res J* 2022;6:295-301.
- [35] Rahman MM and Kabir SMH. In: *Banglapedia*, 1st Ed. Asiatic Society of Bangladesh, Dhaka, Bangladesh1;2003: 403.
- [36] Marie-Magdeleine et al., In vitro effects of *Musa paradisiaca paradisiaca* extracts on four developmental stages of *Haemonchuscontortus*. *Research in Veterinary Science* 96;2014: 127–132.
- [37] Ghani A. Chemical Constituents and Uses. *Medicinal Plants of Bangladesh*; 2003: 315.
- [38] Khare CP. *Indian Medicinal Plants*, Springer Science+ Business Media, New York, USA 2007: 426.
- [39] Waalkes et al., Serotonin, Norepinephrine, and Related Compounds in Bananas. *Science*127 (3299); 1958: 648-650.
- [40] Selvakumar P, Fiber Reinforced Polymer Composites: Classification, Chemical Treatment, Mechanical and Tribological Properties, *Research Journal of Chemistry and Environment*, Vol. 27 (1) January (2023).
- [41] Shanmugavelu KG and Rangaswami G. Tryptophan and Indole Compounds in Banana Ovaries. *Nature*194; 1962: 775–776.



Organized by

Dhaanish Ahmed Institute of Technology, KG chavadi, Coimbatore, India

- [42] Lewis DL, Field WD and Shaw GP. A natural flavonoid present in unripe plantain banana pulp (*Musa paradisiaca sapientum* L. var. *paradisiaca*) protects the gastric mucosa from aspirin-induced erosions. *J.Ethnopharmacol.*65; 1999: 283–288.
- [43] Lewis DA and Shaw GP. A natural flavonoid and synthetic analogues protect the gastric mucosa from aspirin-induced erosions. *J. Nutr. Biochem.* 12; 2001: 95-100.
- [44] Loganathan V and Selvakumar P. In-Vitro assessment of antimicrobial and antioxidant activities of different extracts of *Mallotus rhamnifolius*. *J Pharmacogn Phytochem* 2018;7(6):1995-1999.
- [45] Loganathan V and Selvakumar P. In-Vitro assessment of antimicrobial and antioxidant activities of different extracts of *Mallotus rhamnifolius*. *J Pharmacogn Phytochem* 2018;7(6):1995-1999.
- [46] Houghton PJ, and Skari K. The effect of Indian plants used against snakebite on blood clotting. *J Pharm Pharmacol.* 44; 1992:1054–60.
- [47] P.Selvakumar and V. Loganathan, Preliminary phytochemical investigation of extract of leaves and stem of *Euphorbia hirta*, *International journal of current science* 2012, 48-51. ISSN 2250-1770.
- [48] Ketiku A.O. Chemical composition of unripe (green) and ripe plantain (*Musa paradisiaca paradisiaca*).*J. Sci. Food Agr.*24(6); 1973:703 – 707.
- [49] Emaga et al., Effects of the stage of maturation and varieties on the chemical composition of banana and plantain peels. *Food Chem.* 103; 2007: 590–600.
- [50] Duita PK, Das AK and Banerji N.A Tetracyclic Triterpenoid from *Musa paradisiaca paradisiaca*. *Phytochem.*22 (11); 1983:2563-2564.
- [51] T. Ramakrishnan, M. D. Mohan Gift, S. Chitradevi, R. Jegan, P. Subha Hency Jose, H.N. Nagaraja, Rajneesh Sharma, P. Selvakumar, Sintayehu Mekuria Hailegiorgis, "Study of Numerous Resins Used in Polymer Matrix Composite Materials", *Advances in Materials Science and Engineering*, vol. 2022, Article ID 1088926, 8 pages, 2022. <https://doi.org/10.1155/2022/1088926>
- [52] Ho LH, Noor Aziah AA and Rajeev Bhat. Mineral composition and pasting properties of banana pseudo-stem flour from *Musa paradisiaca acuminata* X *balbisiaca* CV. Awak grown locally in Perak, Malaysia. *International Food Research Journal* 19(4): 2012;1479-1485.



INTERNATIONAL JOURNAL OF INNOVATIVE RESEARCH

IN SCIENCE | ENGINEERING | TECHNOLOGY

 9940 572 462  6381 907 438  ijirset@gmail.com



www.ijirset.com

Scan to save the contact details