



# IJIRSET

International Journal of Innovative Research in  
**SCIENCE | ENGINEERING | TECHNOLOGY**

e-ISSN: 2319-8753 | p-ISSN: 2347-6710



# INTERNATIONAL JOURNAL OF INNOVATIVE RESEARCH

IN SCIENCE | ENGINEERING | TECHNOLOGY

Volume 12, Issue 6, June 2023

**ISSN** INTERNATIONAL  
STANDARD  
SERIAL  
NUMBER  
INDIA

**Impact Factor: 8.423**



9940 572 462



6381 907 438



ijirset@gmail.com



www.ijirset.com

# Common Coupled fixed point theorems for maps under (E.A) and occasionally w-compatible properties in Multiplicative and non-Multiplicative metric Spaces

**Dr Sk. Sadik, D. Anusha and K. Madhavi**

Department of Mathematics, Sir C R R College of Engineering, Eluru Andhra Pradesh, India

Department of Mathematics, Sir C R R College of Engineering, Eluru Andhra Pradesh, India

Department of Mathematics, Sir C R R College of Engineering, Eluru Andhra Pradesh, India

**ABSTRACT:** In this paper, we prove a unique common coupled fixed point theorems for four mappings using (E.A) property in multiplicative metric space and a unique common coupled fixed point theorem for maps satisfying occasionally w-compatible property in a non-multiplicative metric space.

**KEYWORDS:** Multiplicative metric space, E. A. property, occasionally w-compatible

## I. INTRODUCTION AND PRELIMINARIES

The notion of multiplicative metric was introduced by Bashirov et al. [3] in 2008. After that Ozavsar and Cevikel [8] investigated its topological properties and proved some fixed point theorems. For more works on fixed, common fixed point theorems in Multiplicative metric spaces, we refer [5, 2, 6, 9].

The Coupled fixed point is introduced by Bhaskar and Lakshmikantham [4]. Later some of authors proved coupled fixed and coupled common fixed point theorems (See [1, 7, 10, 11, 12]).

The main aim of this paper is to prove a unique common coupled fixed point theorems for four mappings using (E.A) property in multiplicative metric space and a unique common coupled fixed point theorem for maps satisfying occasionally w-compatible property in a non-multiplicative metric space.

Now we recall some basic definitions and examples in multiplicative metric spaces.

**Definition 1.1 ([3]):** Multiplicative metric on a nonempty set X is a mapping  $d: X \times X \rightarrow \mathbb{R}^+$

Satisfying the following conditions:

- (d<sub>1</sub>)  $d(x, y) \geq 1$  for all  $x, y \in X$ ,
- (d<sub>2</sub>)  $d(x, y) = 1$  if and only if  $x = y$ ,
- (d<sub>3</sub>)  $d(x, y) = d(y, x)$ ,
- (d<sub>4</sub>)  $d(x, y) \leq d(x, z).d(z, y)$  for all  $x, y, z \in X$ .

The pair  $(X, d)$  is called a multiplicative metric space.

**Example 1.2** ([9]): Let  $R_+^n$  be the collection of all n-tuples of positive real numbers. And let

$d^*: R_+^n \times R_+^n \rightarrow R$  be defined as

$$d^*(x, y) = \left| \frac{x_1}{y_1} \right|^*, \left| \frac{x_2}{y_2} \right|^*, \dots, \left| \frac{x_n}{y_n} \right|^*$$

Where  $x = (x_1, x_2, \dots, x_n)$ ,  $y = (y_1, y_2, \dots, y_n)$ ,  $\in R_+^n$  and  $|.|^*: R_+ \rightarrow R_+$  is defined as follows

$$|a|^* = \begin{cases} a & \text{if } a \geq 1 \\ \frac{1}{a} & \text{if } a < 1 \end{cases}$$

Then clearly  $d^*(x, y)$  is a multiplicative metric.

**Example 1.3** ([9]): Let  $d: R \times R \rightarrow [1, \infty)$  be defined as  $d(x, y) = a^{|x-y|}$  where  $x, y \in R$  and  $a > 1$ .

Then  $d(x, y)$  is multiplicative metric.

**Definition 1.4** ([8]): Let  $(X, d)$  be a multiplicative metric space,  $x_0$  an arbitrary point in  $X$ , and  $\epsilon > 1$ . A multiplicative open ball  $B(x_0, \epsilon)$  of radius  $\epsilon$  centred at  $x_0$  is the set  $\{z \in X: d(z, x_0) < \epsilon\}$ .

A sequence  $\{x_n\}$  in a multiplicative metric space  $(X, d)$  is said to be multiplicative convergent to some point  $x \in X$ , if for any given  $\epsilon > 1$ , then there exists  $n_0 \in N$  such that  $x_n \in B(x, \epsilon)$  for all  $n \geq n_0$ . If  $\{x_n\}$  converges to  $x$ , we write  $x_n \rightarrow x$  as  $n \rightarrow \infty$ .

**Definition 1.5** ([8]): A sequence  $\{x_n\}$  in a multiplicative metric space  $(X, d)$  is said to be multiplicative convergent to  $x$  in  $X$  if and only if  $d(x_n, x) \rightarrow 1$  as  $n \rightarrow \infty$ .

**Note 1.5(a):** Let  $\{x_n\}$  be a sequence converging to  $x$  in a multiplicative metric space  $(X, d)$  and then  $\lim_{n \rightarrow \infty} d(x_n, y) = d(x, y)$ , for all  $y \in X$ .

**Definition 1.6** ([8]): Let  $(X, d_X)$  and  $(Y, d_Y)$  be two multiplicative metric space and  $x_0$  an arbitrary but fixed element of  $X$ . A mapping  $f: X \rightarrow Y$  is said to be multiplicative continuous at  $x_0$  if and only if  $x_n \rightarrow x_0$  in  $(X, d_X)$  implies that  $f(x_n) \rightarrow f(x_0)$  in  $(Y, d_Y)$ . That is, given arbitrary  $\epsilon > 1$ , then there exists  $\delta > 1$  which depends on  $x_0$  and  $\epsilon$  such that  $d_Y(fx, fx_0) < \epsilon$  for all those  $x$  in  $X$  for which  $d_Y(x, x_0) < \delta$ .

**Definition 1.7** ([8]): A sequence  $\{x_n\}$  in a multiplicative metric space  $(X, d)$  is said to be multiplicative Cauchy sequence if, for any  $\epsilon > 1$ , there exists  $n_0 \in N$  such that  $d(x_m, x_n) < \epsilon$  for all  $m, n \geq n_0$ .

**Definition 1.8** ([8]): A multiplicative metric space  $(X, d)$  is said to be complete if every multiplicative Cauchy sequence  $\{x_n\}$  in  $X$  is multiplicative convergent in  $X$ .

**Definition 1.9** ([2]): Let  $\Psi$  denote the class of functions  $\varphi$ . Suppose  $\varphi: [1, \infty) \rightarrow [1, \infty)$  which are continuous, nondecreasing and  $\varphi(t) = 1$  if and only if  $t = 1$ .

**Note 1.9(a):** If  $\varphi, \psi \in \Psi$  then  $\psi(t) \leq \frac{\psi(t)}{\varphi(t)}$  implies  $t = 1$ .

**Definition 1.10** ([4]): An element  $(x, y) \in X \times X$  is called a coupled fixed point of mapping  $F: X \times X \rightarrow X$  if  $x = F(x, y)$  and  $y = F(y, x)$ .

**Definition 1.11** ([7]): An element  $(x, y) \in X \times X$  is called  $(g_1)$  a coupled coincident point of mappings  $F: X \times X \rightarrow X$  and  $S: X \rightarrow X$  if  $Sx = F(x, y)$  and  $Sy = F(y, x)$ .  $(g_2)$  a common coupled fixed point of mappings  $F: X \times X \rightarrow X$  and  $S: X \rightarrow X$  if  $x = Sx = F(x, y)$  and  $y = Sy = F(y, x)$ .

Now we define the following definitions.

**Definition 1.13:** Let  $(X, d)$  be a multiplicative metric space and  $F: X \times X \rightarrow X$  and  $S: X \rightarrow X$  be mappings. Then the pair  $(F, S)$  is said to satisfy (E.A) property if there exist sequences  $x_n$  and  $y_n$  in  $X$  such that  $\lim_{n \rightarrow \infty} F(x_n, y_n) = \lim_{n \rightarrow \infty} Sx_n = t$  and  $\lim_{n \rightarrow \infty} F(y_n, x_n) = \lim_{n \rightarrow \infty} Sy_n = t'$  for so  $t, t' \in X$

**Definition 1.14 :** Let  $X$  be a non-empty set and  $F: X \times X \rightarrow X$  and  $S: X \rightarrow X$  be mappings. Then the pair  $(F, S)$  is said to be Occasionally w- compatible if there exist  $x, y \in X$  such that  $F(x, y) = Sx$ ,  $F(y, x) = Sy$ ,  $S(F(x, y)) = F(Sx, Sy)$  and  $S(F(y, x)) = F(Sy, Sx)$ .

## MAIN RESULTS

### **Theorem 2.1**

Let  $(X, d)$  be a multiplicative metric space and  $F, G: X \times X \rightarrow X$  and  $S, T: X \rightarrow X$  be Mappings satisfying the following

$$(2.1.1) \psi(d(F(x, y), G(u, v))) \leq \frac{\psi(M(x, y, u, v))}{\varphi(M(x, y, u, v))} \quad \text{for all } x, y, u, v \in X \text{ where } \varphi, \psi \in \Psi \text{ and}$$

$$M(x, y, u, v) = \max \left\{ \begin{array}{l} d(Sx, Tu), d(Sy, Tv), d(Sx, F(x, y)), d(Sy, F(y, x)), \\ d(Tu, G(u, v)), d(Tv, G(v, u)), d(Sx, G(u, v)), \\ d(Sy, G(v, u)), d(Tu, F(x, y)), d(Tv, F(y, x)), \\ \frac{d(Sx, F(x, y))d(Tu, G(u, v))}{d(Sx, Tu)d(Sy, Tv)}, \frac{d(Sy, F(y, x))d(Tv, G(v, u))}{d(Sx, Tu)d(Sy, Tv)}, \\ \frac{d(Tu, F(x, y))d(Sx, G(u, v))}{d(Sx, Tu)d(Sy, Tv)}, \frac{d(Tv, F(y, x))d(Sy, G(v, u))}{d(Sx, Tu)d(Sy, Tv)} \end{array} \right\}$$

(2.1.2) the pairs  $(F, S)$  and  $(G, T)$  are w- compatible.

- (2.1.3) (a)  $(F, S)$  satisfies (E.A) property,)  $F(X \times X) \subseteq T(X)$  and  $S(X)$  is a closed subspace of  $X$  (or)  
(b)  $(G, T)$  satisfies (E.A) property,  $G(X \times X) \subseteq S(X)$  and  $T(X)$  is a closed subspace of  $X$ .

Then  $F, G, S$  and  $T$  have a unique common coupled fixed point.

**Proof.** Suppose (2.1.3) (a) holds. Then there exist sequence  $\{x_n\}$  and  $\{y_n\}$  in  $X$  such that

$$\begin{aligned} \lim_{n \rightarrow \infty} F(x_n, y_n) &= \lim_{n \rightarrow \infty} Sx_n = t \text{ and} \\ \lim_{n \rightarrow \infty} F(y_n, x_n) &= \lim_{n \rightarrow \infty} Sy_n = t^1 \end{aligned} \tag{1}$$

for some  $t, t^1 \in X$ .

Since  $F(X \times X) \subseteq T(X)$ , there exist sequences  $\{z_n\}$  and  $\{w_n\}$  in  $X$  such that  $F(x_n, y_n) = Tz_n \rightarrow t$  and  $F(y_n, x_n) = Tw_n \rightarrow t^1$  as  $n \rightarrow \infty$ . (2)

Consider

$$\Psi \lim_{n \rightarrow \infty} d(t, G(z_n, w_n)) \leq \psi \lim_{n \rightarrow \infty} d(t, F(x_n, y_n))d(F(x_n, y_n), G(z_n, w_n))$$

from  $(d_4)$  and  $\psi$  is non-decreasing

$$= \psi \lim_{n \rightarrow \infty} d(F(x_n, y_n), G(z_n, w_n))$$

$$\leq \lim_{n \rightarrow \infty} \frac{\psi(M(x_n, y_n, z_n, w_n))}{\phi(M(x_n, y_n, z_n, w_n))}, \text{ where}$$

$$M(x_n, y_n, z_n, w_n) = \max \left\{ \begin{array}{l} d(Sx_n, Tz_n), d(Sy_n, Tw_n), d(Sx_n, F(x_n, y_n)), \\ d(Sy_n, F(y_n, x_n)), d(Tz_n, G(z_n, w_n))d(Tw_n, G(w_n, z_n)) \\ d(Sx_n, G(z_n, w_n)), d(Sy_n, G(w_n, z_n)), d(Tz_n, F(x_n, y_n)) \\ d(Tw_n, F(y_n, x_n)), \frac{d(Sx_n, F(x_n, y_n))d(Tz_n, G(z_n, w_n))}{d(Sx_n, Tz_n)d(Sy_n, Tw_n)} \\ \frac{d(Sy_n, F(y_n, x_n))d(Tw_n, G(w_n, z_n))}{d(Sx_n, Tz_n)d(Sy_n, Tw_n)}, \frac{d(Tz_n, F(x_n, y_n))d(Sx_n, G(z_n, w_n))}{d(Sx_n, Tz_n)d(Sy_n, Tw_n)} \\ \frac{d(Tw_n, F(y_n, x_n))d(Sy_n, G(w_n, z_n))}{d(Sx_n, Tz_n)d(Sy_n, Tw_n)} \end{array} \right\}$$

$$\rightarrow \max \left\{ \begin{array}{l} 1, 1, 1, 1, \lim_{n \rightarrow \infty} d(t, G(z_n, w_n)), \\ \lim_{n \rightarrow \infty} d(t^1, G(w_n, z_n)), \lim_{n \rightarrow \infty} d(t, G(z_n, w_n)), \\ \lim_{n \rightarrow \infty} d(t^1, G(w_n, z_n)), 1, 1, \\ \lim_{n \rightarrow \infty} d(t, G(z_n, w_n)), \lim_{n \rightarrow \infty} d(t^1, G(w_n, z_n)) \\ \lim_{n \rightarrow \infty} d(t, G(z_n, w_n)), \lim_{n \rightarrow \infty} d(t^1, G(w_n, z_n)) \end{array} \right\}$$

from (1), (2), Note 1.5(a)

$$( \lim d(t, G(z_n, w_n))) \leq \frac{\psi \max \lim d(t, G(z_n, w_n)), \lim d(t^1, G(w_n, z_n))}{\varphi \max \lim d(t, G(z_n, w_n)), \lim d(t^1, G(w_n, z_n))},$$

Similarly we can show that

$$\psi \lim_{n \rightarrow \infty} d(t^1, G(w_n, z_n)) \leq \frac{\psi \max' \lim_{n \rightarrow \infty} d(t, G(z_n, w_n)), \lim_{n \rightarrow \infty} d(t^1, G(w_n, z_n))}{\varphi \max \lim_{n \rightarrow \infty} d(t, G(z_n, w_n)), \lim_{n \rightarrow \infty} d(t^1, G(w_n, z_n))}$$

Now

$$\psi(\max \{ \lim_{n \rightarrow \infty} d(t, G(z_n, w_n)), \lim_{n \rightarrow \infty} d(t^1, G(w_n, z_n)) \}) = \max$$

$$\psi \left\{ \lim_{n \rightarrow \infty} d(t, G(z_n, w_n)), \psi \lim_{n \rightarrow \infty} d(t^1, G(w_n, z_n)) \right\}$$

$$\leq \frac{\psi \max \lim_{n \rightarrow \infty} d(t, G(z_n, w_n)), \lim_{n \rightarrow \infty} d(t^1, G(w_n, z_n))}{\varphi \max \lim_{n \rightarrow \infty} d(t, G(z_n, w_n)), \lim_{n \rightarrow \infty} d(t^1, G(w_n, z_n))},$$

From Note 1.9(a), we have

$$\max \{ \lim_{n \rightarrow \infty} d(t, G(z_n, w_n)), \lim_{n \rightarrow \infty} d(t^1, G(w_n, z_n)) \} = 1$$

Which implies that

$$\lim_{n \rightarrow \infty} d(t, G(z_n, w_n)) = 1 = \lim_{n \rightarrow \infty} d(t^1, G(w_n, z_n))$$

Thus

$$\lim_{n \rightarrow \infty} G(z_n, w_n) = t \text{ and } \lim_{n \rightarrow \infty} G(w_n, z_n) = t^1 \quad (3)$$

Since  $S(X)$  is a closed subspace of  $X$ , there exist  $u, v \in X$  such that

$$\lim_{n \rightarrow \infty} Sx_n = t = Su \text{ and } \lim_{n \rightarrow \infty} Sy_n = t^1 = Sv. \quad (4)$$

From Note 1.5(a), (3),(4) and using continuity of  $\psi$ , we have

$$\begin{aligned} \psi(d(F(u, v), Su)) &= \lim_{n \rightarrow \infty} \psi(d(F(u, v), G(z_n, w_n))) \\ &\leq \lim_{n \rightarrow \infty} \frac{\psi(M(u, v, z_n, w_n))}{\varphi(M(u, v, z_n, w_n))} \quad \text{From (2.1.1)} \end{aligned}$$

$$\begin{aligned} M(u, v, z_n, w_n) &= \max \left\{ \begin{array}{l} d(Su, T, z_n), d(Sv, Tw_n), d(Su, F(u, v)), \\ d(Sv, F(v, u)), d(Tz_n, G(z_n, w_n)), d(Tw_n, G(w_n, z_n)), \\ d(Su, G(z_n, w_n)), d(Sv, G(w_n, z_n)), d(Tz_n, F(u, v)), \\ d(Tw_n, F(v, u)), \frac{d(Su, F(u, v))d(Tz_n, G(z_n, w_n))}{d(Su, Tz_n)d(Sv, Tw_n)}, \\ \frac{d(Sv, F(v, u))d(Tw_n, G(w_n, z_n))}{d(Su, Tz_n)d(Sv, Tw_n)}, \frac{d(Su, G(z_n, w_n))d(Tz_n, F(u, v))}{d(Su, Tz_n)d(Sv, Tw_n)}, \\ \frac{d(Sv, G(w_n, z_n))d(Tw_n, F(v, u))}{d(Su, Tz_n)d(Sv, Tw_n)}, \end{array} \right\} \\ &\rightarrow \max \{d(Su, F(u, v)), d(Sv, F(v, u))\} \text{ as } n \rightarrow \infty \end{aligned}$$

Thus

$$\psi(d(F(u, v), Su)) \leq \frac{\psi(\max\{d(Su, F(u, v)), d(Sv, F(v, u))\})}{\varphi(\max\{d(Su, F(u, v)), d(Sv, F(v, u))\})}$$

Similarly we can show that

$$\psi(d(F(u, v), Sv)) \leq \frac{\psi(\max\{d(Su, F(u, v)), d(Sv, F(v, u))\})}{\varphi(\max\{d(Su, F(u, v)), d(Sv, F(v, u))\})}$$

Thus

$$\psi(\max\{d(F(u, v), Su), d(F(v, u), Sv)\}) = \max\{\psi(d(F(u, v), Su)), \psi(d(F(v, u), Sv))\}$$

$$\leq \frac{\psi(\max\{d(Su, F(u, v)), d(Sv, F(v, u))\})}{\varphi(\max\{d(Su, F(u, v)), d(Sv, F(v, u))\})}$$

From Note 1.9 (a), we have

$$F(u, v) = Su = t \text{ and } F(v, u) = Sv = t^1. \quad (5)$$

Since the pair  $(F, S)$  is  $w$ -compatible, we have

$$F(t, t^1) = St \text{ and } F(t^1, t) = St^1. \quad (6)$$

Since  $F(X \times X) \subseteq T(X)$  there exist  $\alpha, \beta \in X$  such that  $t = F(u, v) = T\alpha$

And  $t^1 = F(v, u) = T\beta$ .

Consider

$$\Psi(d(t, G(\alpha, \beta))) = \psi(d(F(u, v)), G(\alpha, \beta))$$

$$\leq \frac{\psi(M(u, v, \alpha, \beta))}{\varphi(M(u, v, \alpha, \beta))}$$

$$M(u, v, \alpha, \beta) = \max \left\{ \begin{array}{l} d(Su, T\alpha), d(Sv, T\beta), d(Sv, F(u, v)) \\ d(Sv, F(v, u)), d(T\alpha, G(\alpha, \beta)), d(\beta, G(\beta, \alpha)), \\ d(Su, G(\alpha, \beta)), d(Sv, G(\beta, \alpha)), d(T\alpha, F(u, v)), \\ d(T\beta, F(v, u)), \frac{d(Sx_n, F(u, v))d(T\alpha, G(\alpha, \beta))}{d(Su, T\alpha) + d(Sv, T\beta)}, \\ \frac{d(Sv, F(v, u))d(T\beta, G(\beta, \alpha))}{d(Su, T\alpha) + d(Sv, T\beta)}, \frac{d(T\alpha, F(u, v))d(Su, G(\alpha, \beta))}{d(Su, T\alpha) + d(Sv, T\beta)} \\ \frac{d(T\beta, F(v, u))d(Sv, G(\beta, \alpha))}{d(Su, T\alpha) + d(Sv, T\beta)} \end{array} \right\}$$

$\rightarrow \max \{d(t, G(\alpha, \beta)), d(t^1, G(\beta, \alpha))\}$

$$\psi(d(t, G(\alpha, \beta))) \leq \frac{\psi(\max\{d(t, G(\alpha, \beta)), d(t^1, G(\beta, \alpha))\})}{\varphi(\max\{d(t, G(\alpha, \beta)), d(t^1, G(\beta, \alpha))\})}$$

Similarly we can show that

$$\psi(d(t^1, G(\beta, \alpha))) \leq \frac{\psi(\max\{d(t, G(\alpha, \beta)), d(t^1, G(\beta, \alpha))\})}{\varphi(\max\{d(t, G(\alpha, \beta)), d(t^1, G(\beta, \alpha))\})}$$

Now

$$\psi(\max\{d(t, G(\alpha, \beta)), d(t^1, G(\beta, \alpha))\}) = \max\{\psi(d(t, G(\alpha, \beta))), \psi(d(t^1, G(\beta, \alpha)))\}$$

Hence

$$\leq \frac{\psi(\max\{d(t, G(\alpha, \beta)), d(t^1, G(\beta, \alpha))\})}{\varphi(\max\{d(t, G(\alpha, \beta)), d(t^1, G(\beta, \alpha))\})} \leq 1$$

But  $\varphi(\max\{d(t, G(\alpha, \beta)), d(t^1, G(\beta, \alpha))\}) \geq 1$

Hence  $\varphi(\max\{d(t, G(\alpha, \beta)), d(t^1, G(\beta, \alpha))\}) = 1$

Implies  $(\max\{d(t, G(\alpha, \beta)), d(t^1, G(\beta, \alpha))\}) = 1$

Implies

$$d(t, G(\alpha, \beta)) = 1 = d(t^1, G(\beta, \alpha))$$

$$G(\alpha, \beta) = T\alpha = t \text{ and } G(\beta, \alpha) = T\beta = t^1. \quad (7)$$

Since  $(G, T)$  is w-compatible, we have

$$G(t, t^1) = Tt \text{ and } G(t^1, t) = Tt^1 \quad (8)$$

Now

$$\begin{aligned} \Psi(d(St, t)) &\leq \psi(d(F(t, t^1), t)) \\ &= \psi \lim_{n \rightarrow \infty} d(F(t, t^1), G(z_n, w_n)) \\ &= \lim_{n \rightarrow \infty} \psi(d(F(t, t^1), G(z_n, w_n))) \\ &\leq \lim_{n \rightarrow \infty} \frac{\psi(M(t, t^1, z_n, w_n))}{\varphi(M(t, t^1, z_n, w_n))} \end{aligned}$$

$$M(t, t^1, z_n, w_n) = \max \left\{ \begin{array}{l} d(St, Tz_n), d(st^1, Tw_n), d(St, F(t, t^1)), \\ d(St^1, F(t^1, t)), d(T, G(z_n, w_n)), d(Tw_n, G(w_n, z_n)), \\ d(St, G(z_n, w_n)), d(S t^1, G(w_n, z_n)), d(Tz_n, F(t, t^1)), \\ d(Tw_n, F(t^1, t)), \frac{d(St, F(t, t^1))d(Tz_n, G(z_n, w_n))}{d(St, Tz_n)d(St^1, Tw_n)} \\ \frac{d(S t^1, F(t^1, t))d(Tw_n, G(w_n, z_n))}{d(St, Tz_n)d(S t^1, Tw_n)}, \frac{d(Tz_n, F(t, t^1))d(S t^1, G(w_n, z_n))}{d(St, Tz_n)d(S t^1, Tw_n)} \\ \frac{d(Tw_n, F(t^1, t))d(St, G(w_n, z_n))}{d(St, Tz_n)d(St^1, Tw_n)} \end{array} \right\}$$

Thus

$$\rightarrow \max \{d(St, t), d(St^1, t^1)\}, \text{ from (2), (3), (6)}$$

$$\psi(d(St, t)) \leq \frac{\psi(\max\{d(St, t), d(St^1, t^1)\})}{\varphi(\max\{d(St, t), d(St^1, t^1)\})}$$

Similarly we can show that

$$\psi(d(St^1, t^1)) \leq \frac{\psi(\max\{d(St, t), d(St^1, t^1)\})}{\varphi(\max\{d(St, t), d(St^1, t^1)\})}$$

Now

$$\begin{aligned} \Psi(\max\{d(St, t), d(St^1, t^1)\}) &= \max\{\psi(d(St, t)), \psi(d(St^1, t^1))\} \\ &\leq \frac{\psi(\max\{d(St, t), d(St^1, t^1)\})}{\varphi(\max\{d(St, t), d(St^1, t^1)\})} \end{aligned}$$

By Note 1.9 (a), we have  $\max\{d(St, t), d(St^1, t^1)\} = 1$ . Which in turn implies that  $St = t$  and  $St^1 = t^1$ .

Similarly we can show that  $Tt = t$  and  $Tt^1 = t^1$ .

Now from (6) and (8), it follows that  $(t, t^1)$  is a common coupled fixed point of  $F, G, S$  and  $T$ . Uniqueness of common coupled fixed point follows easily from (2.2.1). Similarly the theorem follows when 2.1.3(b) holds.

Now we give following example to illustrate Theorem 2.1.

### Example 2.2

Suppose that  $X = [0, 1]$ . Define  $d(x, y) = 2^{|x-y|}$  for all  $x, y \in X$ . Let  $F, G: X \times X \rightarrow X$  and  $S, T: X \times X \rightarrow X$  be defined as

$$F(x, y) = \frac{x^2 + y^2}{16}, G(x, y) = \frac{x + y}{16}, S_x = \frac{x^2}{2}, T_x = \frac{x}{2}, \psi(t) = t, \varphi(t) = t^{\frac{3}{4}}$$

Then  $(F, S)$  satisfies (E.A) property with  $\{x_n\} = \{\frac{1}{n}\}$  and  $\{y_n\} = \{\frac{1}{2n}\} \in X$ .

$$\begin{aligned} d(F(x, y), G(u, v)) &= 2^{\left|\frac{x^2+y^2}{16} - \frac{u+v}{16}\right|} \\ &\leq 2^{\frac{1}{16}[|x^2-u|+|y^2-v|]} \\ &\leq 2^{\frac{1}{16}2\max[|x^2-u|, |y^2-v|]} \\ &= \max\left\{2^{\frac{|x^2-u|}{8}}, 2^{\frac{|y^2-v|}{8}}\right\} \\ &= \max\left\{2^{\left|\frac{x^2-u}{2}\right|}, 2^{\left|\frac{y^2-v}{2}\right|}\right\}^{\frac{1}{4}} \end{aligned}$$

$$\begin{aligned}
 &= (\max \{d(Sx, Tu), d(Sy, Tv)\})^{\frac{1}{4}} \\
 &= \frac{\psi(M(x, y, u, v))}{\varphi(M(x, y, u, v))}
 \end{aligned}$$

Thus, the inequality (2.1.1) is satisfied. One can easily verify the remaining conditions of Theorem 2.1. Clearly (0, 0) is unique common coupled point of F, G, S and T.

**Remark:** There are some functions  $d: X \times X \rightarrow [1, \infty)$  such that  $d(x, y) = 1 \Leftrightarrow x = y$ ,  $d(x, y) \neq d(y, x)$  and  $d(x, y) \nleq d(x, z)d(z, y)$ .

For example, Let  $X = \{1, 2, 3\}$ ,  $d(1, 1) = d(2, 2) = d(3, 3) = 1$ ,  $d(1, 2) = 2$ ,  $d(2, 1) = 1.2$ ,  $d(1, 3) = d(3, 1) = 1.3$ ,  $d(2, 3) = d(3, 2) = 1.5$ .

Then  $d(1, 2) \neq d(2, 1)$  and  $d(1, 2) \nleq d(1, 3)d(3, 2)$ .

**Theorem 2.3** Let  $X$  be a non-empty set and  $d: X \times X \rightarrow [1, \infty)$  be a function such that  $d(x, y) = 1$  iff  $x = y$ . Let  $F, G: X \times X \rightarrow X$  and  $S, T: X \rightarrow X$  be satisfying the following

(2.3.1)

$$\psi(d(F(x, y), G(u, v))) \leq (\psi(M(x, y, u, v))) / (\varphi(M(x, y, u, v))) \quad \text{for all } x, y, u, v \in X,$$

Where  $\psi: [1, \infty) \rightarrow [1, \infty)$  be such that  $\psi(t) = 1$  iff  $t = 1$  and  $\psi(t) = 1$  iff  $t = 1$  and

$$M(x, y, u, v) = \max \left\{ \frac{d(Sx, Tu), d(Sy, Tv), d(Sx, F(x, y)), d(Sy, F(y, x)),}{d(Tu, G(u, v)), d(Tv, G(v, u)), d(F(x, y), Tu)}, \right. \\ \left. \frac{d(F(y, x), Tv), d(Sx, G(u, v)), d(Sy, G(v, u)),}{d(Sx, F(x, y))d(Tu, G(u, v)), d(Sy, F(y, x))d(Tv, G(v, u))}, \right. \\ \left. \frac{d(Sx, G(u, v))d(F(x, y), Tu)}{d(Sx, Tu)d(Sy, Tv)}, \frac{d(Sy, G(v, u))d(F(y, x), Tv)}{d(Sx, Tu)d(Sy, Tv)} \right\}$$

(2.2.2) the pairs  $(F, S)$  and  $(G, T)$  are occasionally w-compatible Then there exists a unique

Common coupled fixed point of F, G, S and T

**Proof.** Since the pairs  $(F, S)$  and  $(G, T)$  are occasionally w-compatible, there exist points  $(x, y), (\tilde{x}, \tilde{y}) \in X \times X$  such that

$$Sx = F(x, y), Sy = F(y, x) \text{ and}$$

$$S(F(x, y)) = F(Sx, Sy), S(F(y, x)) = F(Sy, Sx) \quad (9)$$

$$T\tilde{x} = G(\tilde{x}, y), Ty = G(y, \tilde{x}) \text{ and}$$

$$T(G(\tilde{x}, y)) = G(T\tilde{x}, Ty), T(G(y, \tilde{x})) = G(Ty, T\tilde{x}) \quad (10)$$

Now

$$S^2x = F(Sx, Sy), S^2y = F(Sy, Sx) \quad (11)$$

$$T^2\tilde{x} = G(T\tilde{x}, Ty), T^2y = G(Ty, T\tilde{x}) \quad (12)$$

Now, we shall show that  $Sx = T\tilde{x}$  and  $Sy = T$

From (2.3.1), consider  $y'$

$$\psi(d(F(x, y), G(\tilde{x}, y'))) \leq \frac{\psi(M(x, y, \tilde{x}, y'))}{\varphi(M(x, y, \tilde{x}, y'))}$$

where

$$\begin{aligned}
 M(x, y, \tilde{x}, \tilde{y}) &= \max \left\{ \frac{d(Sx, Tx), d(Sy, Ty), d(Sx, F(x, y)), d(Sy, F(y, x)),}{d(Tx, G(\tilde{x}, \tilde{y})), d(Ty, G(\tilde{y}, \tilde{x})), d(F(x, y), Tx), d(F(y, x), Ty), d(Sx, G(x, y)), d(Sy, G(y, x))}, \right. \\
 &\quad \left. \frac{d(Sx, F(x, y))d(Tx, G(\tilde{x}, \tilde{y}))}{d(Sx, Tx)d(Sy, Ty)}, \frac{d(Sy, F(y, x))d(Ty, G(\tilde{y}, \tilde{x}))}{d(Sx, Tx)d(Sy, Ty)} \right. \\
 &\quad \left. \frac{d(Sx, G(\tilde{x}, \tilde{y}))d(F(x, y), Tx)}{d(Sx, Tx)d(Sy, Ty)}, \frac{d(Sy, G(\tilde{y}, \tilde{x}))d(F(y, x), Ty)}{d(Sx, Tx)d(Sy, Ty)} \right. \\
 &= \max \quad d(Sx, T\tilde{x}), d(Sy, T\tilde{y})
 \end{aligned}$$

Thus

$$\psi(d(Sx, T\tilde{x})) \leq \frac{\psi(\max\{d(Sx, T\tilde{x}), d(Sy, T\tilde{y})\})}{\varphi(\max\{d(Sx, T\tilde{x}), d(Sy, T\tilde{y})\})} \quad 13$$

Similarly we can show that

$$\psi(d(Sy, T\tilde{y})) \leq \frac{\psi(\max\{d(Sx, T\tilde{x}), d(Sy, T\tilde{y})\})}{\varphi(\max\{d(Sx, T\tilde{x}), d(Sy, T\tilde{y})\})} \quad 14$$

Now we have

$$\begin{aligned}
 \psi(\max\{d(Sx, T\tilde{x}), d(Sy, T\tilde{y})\}) &= \max\{\psi(d(Sx, T\tilde{x})), \psi(d(Sy, T\tilde{y}))\} \\
 &\leq \frac{\psi(\max\{d(Sx, T\tilde{x}), d(Sy, T\tilde{y})\})}{\varphi(\max\{d(Sx, T\tilde{x}), d(Sy, T\tilde{y})\})}
 \end{aligned}$$

**From (13), (14)**

Hence  $\varphi(\max\{d(Sx, T\tilde{x}), d(Sy, T\tilde{y})\}) \leq 1$  so that

$$\max\{d(Sx, T\tilde{x}), d(Sy, T\tilde{y})\} = 1$$

Hence We have  $Sx = T\tilde{x}$  and  $Sy = T\tilde{y}$

Now write

$$u = Sx = T\tilde{x} \text{ and } v = Sy = T\tilde{y}. \quad (15)$$

Thus, from (11) and (12), we get

$$Su = S^2x = F(Sx, Sy) = F(u, v), Sv = S^2y = F(Sy, Sx) = F(v, u) \quad (16)$$

$$Tu = T^2\tilde{x} = G(S\tilde{x}, S\tilde{y}) = G(u, v), Tv = S^2\tilde{y} = G(S\tilde{y}, S\tilde{x}) = G(v, u). \quad (17)$$

Now we shall show that  $u = Su = Tu$  and  $v = Sv = Tv$ .

From (2.1.2), consider

$$\psi(d(F(u, v), G(\tilde{x}, \tilde{y}))) \leq \frac{\psi(M(u, v, \tilde{x}, \tilde{y}))}{\phi(M(u, v, \tilde{x}, \tilde{y}))}$$

where

$$M(u, v, x^*, y^*) = \max \left\{ \begin{array}{l} d(Su, Tx^*), d(Sv, Ty^*), d(Su, F(u, v)), d(Sv, F(v, u)), \\ d(Tx^*, G(x^*, y^*)), d(Ty^*, G(y^*, x^*)), d(F(u, v), Tx^*), \\ d(F(v, u), Ty^*), d(Su, G(x^*, y^*)), d(Sv, G(y^*, x^*)), \\ \frac{d(Su, F(u, v))d(Tx^*, G(x^*, y^*))}{d(Su, Tx^*)d(Sv, Ty^*)}, \frac{d(Sv, F(v, u))d(Ty^*, G(y^*, x^*))}{d(Su, Tx^*)d(Sv, Ty^*)} \\ \frac{d(Su, G(x^*, y^*))d(F(u, v), Tx^*)}{d(Su, Tx^*)d(Sv, Ty^*)}, \frac{d(Sv, G(y^*, x^*))d(F(v, u), Ty^*)}{d(Su, Tx^*)d(Sv, Ty^*)} \end{array} \right\}$$

$$= \max d(Su, u), d(Sv, v)$$

Thus

$$\psi(d(Su, u)) \leq \frac{\psi(\max\{d(Su, u), d(Sv, v)\})}{\varphi(\max\{d(Su, u), d(Sv, v)\})} \quad 18$$

Similarly we can show that

$$\psi(d(Sv, v)) \leq \frac{\psi(\max\{d(Su, u), d(Sv, v)\})}{\varphi(\max\{d(Su, u), d(Sv, v)\})} \quad 19$$

Now

$$\begin{aligned} \psi(\max\{d(Su, u), d(Sv, v)\}) &= \max \{\psi(d(Su, u)), \psi(d(Sv, v))\} \\ &\leq \frac{\psi(\max\{d(Su, u), d(Sv, v)\})}{\varphi(\max\{d(Su, u), d(Sv, v)\})} \end{aligned}$$

From (18), (19)

Which in turn yields that  $\varphi(\max d(Su, u), d(Sv, v)) \leq 1$

So that  $\varphi(\max d(Su, u), d(Sv, v)) = 1$  implies  $\max\{d(Su, u), d(Sv, v)\} = 1$  Hence

$$Su = u \text{ and } Sv = v \quad (20)$$

In the similar lines we can show that

$$Tu = u \text{ and } Tv = v \quad (21)$$

Now from (16), (17), (20) and (21) it follows that  $(u, v)$  is a common coupledfixed point of  $F, G, S$  and  $T$ .

Uniqueness of common coupled fixed point follows easily from (2.3.1).

## REFERENCES

- [1] Abbas, M., Alikhan, M., Radenović S. Common coupled fixed point theorems in cone metric spaces for w-compatible mappings. Applied Mathematics and Computation 2010; Volume 217, Issue 1, 195 - 202.
- [2] Abbas, M., La Sen, M., Nazir,T., Common fixed points of generalized rational type cocyclic mappings in multiplicative metric spaces, Descrete Dynamics in nature and society, Volume 2015, Article ID 532725, 10pages, <http://dx.doi.org/10.1155/2015/532725>.
- [3] Bashirov, A. E., Kurplnara, E. M., Ozyapici, A., Multiplicative calculus and its applications, J. Math. Anal. Appl., 337 (2008), 36 - 48. <http://dx.doi.org/10.1016/j.jmaa.2007.03.081>
- [4] Bhaskar, T.G. and Lakshmikantham V. Fixed point theorems in partially ordered metric spaces and applications. Nonlinear Analysis 2006, 65, 1379 - 1393.
- [5] He, X., Song, M. and Chen, D., Common fixed points for weak commutative mappings on a multiplicative metric space, Fixed Point Theory Appl., 48 (2014), 9 pages. <http://dx.doi.org/10.1186/1687-1812-2014-48>
- [6] Kang, S. M., Nagpal, P., Sudhir Kumar, G. and Sanjay Kumar, Fixed points for multiplicative

expansive mappings in multiplicative metric spaces, International Journal of Mathematical Analysis ,Vol. 9, 2015, no. 39, 1939 - 1946 ,doi.org/10.12988/ijma.2015.54130.

- [7] Lakshmikantham V. and Čirić Lj. Coupled fixed point theorems for non- linear contractions in partially ordered metric spaces. Nonlinear Analysis 2009, 70, 4341 - 4349.
- [8] Ozavsar, M. and Cevikel, A. C., Fixed point of multiplicative contraction mappings on multiplicative metric spaces, arXiv:1205.5131v1 [math.GM],2012.
- [9] Sarwar, M. and Badshah-e-Rome, Some unique fixed point theorems in multiplicative metric space, arXiv:1410.3384v2 [math.GM] 29 Dec 2014.
- [10] Rao, K.P.R., Kishore, G.N.V. and Van Luong N., A unique common cou- pled fixed point theorem for four maps under  $\psi$  -  $\varphi$  contractive condition in partial metric spaces. CUBO A Mathematical Journal 2012, Vol.14, No03, 115 - 127.
- [11] Rao, K.P.R., Kishore, G.N.V. and Raju, V.C.C., A coupled fixed point theorem for two pairs of w - compatible maps using altering distance function in partial metric space. Journal of Advanced Research in Pure Mathematics 2012, 4(4), 96 - 114.
- [12] Shatanawi, W., Samet, B., Abbas, M., Coupled fixed point theorems for mixed monotone mappings in ordered partial metric spaces. Mathematical and Computer Modeling 2012, 55, 680 - 687.



SJIF Scientific Journal Impact Factor

Impact Factor: 8.423



# INTERNATIONAL JOURNAL OF INNOVATIVE RESEARCH

IN SCIENCE | ENGINEERING | TECHNOLOGY

9940 572 462 6381 907 438 ijirset@gmail.com

[www.ijirset.com](http://www.ijirset.com)



Scan to save the contact details