



(A Monthly, Peer Reviewed, Refereed, Scholarly Indexed, Open Access Journal)



Impact Factor: 8.699

Volume 14, Issue 5, May 2025

⊕ www.ijirset.com 🖂 ijirset@gmail.com 🖄 +91-9940572462 🕓 +91 63819 07438





www.ijirset.com |A Monthly, Peer Reviewed & Refereed Journal| e-ISSN: 2319-8753| p-ISSN: 2347-6710|

Volume 14, Issue 5, May 2025

|DOI: 10.15680/IJIRSET.2025.1405211|

Production of Algae-Based Biofertilizer using Bio Extracted Nutrient Solution

Vishnushree T.S, Jeyanthi Jeyaraman

Student, Department of Civil (Environmental) Engineering, Government college of Technology, Coimbatore,

Tamil Nadu, India

Professor, Department of Civil (Environmental) Engineering, Government college of Technology, Coimbatore,

Tamil Nadu, India

ABSTRACT: This study explores a sustainable and ecofriendly methodfor producing algae based biofertilizer utilizing bio extracted nutrient solutions derived from organic waste materials. The nutrient medium was prepared using ethanol extracted potassium from musaparadisiaca(banana peels) and nitrogen from azadirachtaindica (neem leaves), both collected from local solid waste sources. Algal biomass was sourced from AlapuramLake, Dharmapuri, and cultured using the extracted nutrients. Growth was monitored over a 30-day period, with periodic measurements of Ph, turbidity, color and dry weight to assess algal development. The optimal growth was observed on the 15th day with p H 7.4, Turbidity 33.4 NTU, and temperature 29.5 C. the final dry algal biomass was processed into powdered biofertilizer, achieving 14.53 g from an initial 9.9 g dry weight. The results confirm the viability of utilizing plant waste derived nutrients for algae cultivation and biofertilizer production, offering a low cost and environmentally sustainable, alternative to chemical fertilizers. This project highlights the potential for integrating waste valorization in modern agriculture.

KEY WORDS: Biofertilizer, algae, Nutrient extraction, Neem leaves, banana peels, organic waste, sustainability.

I. INTRODUCTION

The increasing global demand for sustainable agricultural practices has highlighted the importance of bio fertilizers as an eco friendly alternative to chemical fertilizers. Among various bio fertilizer sources, algae have emerged as a promising source due to their ability to enhance soil fertility and plant growth while reducing environmental harm(Ammar et al.,2022). Algae are rich in essential nutrients such a nitrogen, phosphorous and potassium, which are vital for plant health. Due to its benefits many small scale and large scale sectors develop algae at huge scale for various purposes. At the peak of its demand and competence, the algae are grown with chemicals for its rapid and bulk growth(chatterjee. A, 2017). Algal blooms affect the lives of organisms that live in ponds, lakes, streams, rivers etc. The excessive use of chemical fertilizers has raised concerns over soil degradation, water pollution, and ecosystem imbalance. Biofertilizers, particularly those derived from algae, offer a sustainable solution by enhancing nutrient cycling and promoting healthier plant growth. Algae-based fertilizers not only improve soil structure but also increase microbial activity, which is crucial for long-term soil health. Furthermore, the integration of organic waste materials boosts the nutrient value of the biofertilizer. This holistic approach supports both agricultural productivity and environmental conservation.

Even though its harm, it provides essential nutrients and act as soilconditioners..Hence, it was selected as raw material for algal based biofertilizer. The waste such as shredded azadirachtaindica and the musaparadisiacalinn peel was taken for the nitrogen and potassium extract (godlewska.k et al., 2021). In the study, an attempt had been made to investigate the effect of bio extracted nutrient solution on the algal growth. The potential analysis had been made by analyzing the algal growth with applying varying dosage of bio extracted nutrient solution for the period of 30 days. The qualitative parameters had been observed.





www.ijirset.com |A Monthly, Peer Reviewed & Refereed Journal| e-ISSN: 2319-8753| p-ISSN: 2347-6710|

Volume 14, Issue 5, May 2025

|DOI: 10.15680/IJIRSET.2025.1405211|

II. MATERIALS AND METHODS

2.1 Materials

Equipment included a hot air oven, ph meter, turbidity meter, porcelain mortar and pestle, whatman filter paper, weighing balance and white plastic culture containers.

2.2 Bio Nutrient Extraction

Neem leaves and banana peels were dried, powdered, and soaked in ethanol at a ratio 1:10 ratio. Neem extract provided nitrogen rich nutrients while banana peel extract was high in potassium (kattubwe et al., <u>2020</u>) (sriraj. P et al., <u>2022</u>). The p H of the potassium extract was 8.43 and the nitrogen extract was 5.49, confirming successful extraction.



2.3 Algal Collection and Culturing

Algae were collected fromAlapuram Lake. Fresh samples were cultured in water supplemented with 2ml each of nitrogen and potassium extracts on the 10th and 20th day of the 30- day cycle. Environmental factors such as sunlight, aeration, and temperature were maintained naturally.



2.4 Biofertilizer Preparation

The grown algae were dried in hot air oven and powdered. The final dry weight of cultured algae was 14.53 g, compared to 9.9 g initially. Neem and banana powder were also used as carrier material to test future soil application. (Gajalakshmi.S et al., <u>2004</u>)





www.ijirset.com |A Monthly, Peer Reviewed & Refereed Journal| e-ISSN: 2319-8753| p-ISSN: 2347-6710|

Volume 14, Issue 5, May 2025 |DOI: 10.15680/IJIRSET.2025.1405211|





Cultured Algal Biomass Dried in oven



Grinded biomass grinding process

III. RESULTS AND DISCUSSION

3.1 Nutrient Extraction

246 ml of potassium and 117 ml of nitrogen solution were extracted from 25.24g of banana peel and 12.14 g of neem leaves, respectively.

3.1.1 potassium extraction

Serial Number	Parameters	Values	Units
1	Initial pH	7.33	-
2	Final pH	8.43	-
3	Weight of Collected musaparadisiacalinn	26	Gram (g)
4	Grinded musaparadisiacalinn	25.24	Gram (g)
5	Obtained extracted solution	246	milliliter (ml)

Table 3.1 Parameters Observed inPotassiumExtraction Process

The measured 246 ml of potassium nutrient solution had been extracted from 25.24 g of powdered banana peel. The obtained value of pH for the banana peel extraction is 8.43

According to the study, the pH of banana peel is 9.91 and the pH of ethanol is 7.33. The range of extract varies between the ranges 8-11. Hence it is within the range. It would not cause any affect, when it applied into the soil.





www.ijirset.com |A Monthly, Peer Reviewed & Refereed Journal| e-ISSN: 2319-8753| p-ISSN: 2347-6710|

Volume 14, Issue 5, May 2025

|DOI: 10.15680/IJIRSET.2025.1405211|



Figure 3.1 pH Graph for Banana Peel Extract.

4.1.2 Nitrogen Extraction Process

The neem extract range would be acidic in nature. The obtained pH Value was in acidic nature (<7). Due to the presence of ascorbic acid in neem leaves, the excess hydrogen ions exist in the neem leaves extracted solution. The decreased pH value is due to presence of excess hydrogen (H^+) ions. It ensured the extraction had been occurred.

Serial Number	Parameters	Values	Units
1	Initial p H	7.33	-
2	Final p H	5.49	-
3	Weight of collected azadirachtaindica	15	Gram (g)
4	Grinded azadirachtaindica	12.14	Gram (g)
5	Obtained extracted solution	117	Milliliter (ml)

The neem leaves were collected, dried and soaked in ethanol in 1:10 ratio i.e., 12.14 g in 120 ml of ethanol. The obtained extract amount was 117 ml from 12.14 g with pH 5.49.



Figure 3.2 pH Graph for Neem Leaves Extract





www.ijirset.com |A Monthly, Peer Reviewed & Refereed Journal| e-ISSN: 2319-8753| p-ISSN: 2347-6710|

Volume 14, Issue 5, May 2025

|DOI: 10.15680/IJIRSET.2025.1405211|

3.2 Algal Growth Parameters

p H increased from 7.02 to 8.22; turbidity increased from 15.3NTU to 54.1 NTU; water color shifted from pale yellow to green. The optimal growth conditions were on day 15 at p H 7.4, Turbidity 33.4 NTU, aThe algal growth was analyzed at below conditions,

Serial		
Number	Particulars	Conditions
1	Growth period	30 days
2	Light : dark	8:12 hrs
3	Aeration	Atmospheric air
4	Light	Sun light
5	Growth medium	Subsurface water
6	p H of growth medium	7.02
7	Turbidity of growth medium	15.3 NTU
8	Temperature	29.5 °C
9	Nutrient supplement	twice (10 th , 20 th day)
10	N	2 ml / 500 ml
11	K	2 ml / 500 ml
12	N:K	1:1
13	pH of Nitrogen Solution	5.49
14	pH of Potassium Solution	8.43

Table 3.3 Growth Conditions of Algal Culture

The observed pH values throughout the 30 days of algal growth period were given below.

Table 3.4 pH Values Observed during Growth Period

Serial Number	Day	pH Values
1	1	7.02
2	5	7.06
3	10	7.12





www.ijirset.com |A Monthly, Peer Reviewed & Refereed Journal| e-ISSN: 2319-8753| p-ISSN: 2347-6710|

Volume 14, Issue 5, May 2025

|DOI: 10.15680/IJIRSET.2025.1405211|

4	15	7.4
5	20	7.7
6	25	7.98
7	30	8.22

Here pH resulted as alkalinity i.e., 8.22 > 7.02 > 7.00

The pH, presence of hydrogen ions represents the algal growth by varying the pH value. A study states that, the pH level would increase after algal growth, because algae consume dissolved carbon dioxide, release oxygen and hydroxide alkalinity during photosynthesis. Due to the release of hydroxide ions, the pH range would increase.

The obtained pH value from algal growth analysis was 8.22 at 29.5°C. Initial pH was 7.02 (< 8.22). Hence alkalinity (>7) resembled. Here, the gradual increase in pH had been observed. This ensured the algal growth.



Figure 3.3 pH Graph of Observed Algal Growth

Serial Number	Day	Turbidity Values (NTU)
1	1	15.3
2	5	17
3	10	20
4	15	33.4
5	20	40.3
6	25	45
7	30	54.1

Table 3.5 Turbidity Values Observed During Growth Period





www.ijirset.com |A Monthly, Peer Reviewed & Refereed Journal| e-ISSN: 2319-8753| p-ISSN: 2347-6710|

Volume 14, Issue 5, May 2025

|DOI: 10.15680/IJIRSET.2025.1405211|

The algal growth makes the water turbid. Initial turbidity of growth medium was 15.3 NTU.Final turbidity of growth medium was 54.1 NTU.



Figure 3.4 Turbidity Graph of Observed Algal Growth

4.2.3 Color Analysis

Generally, the presence of algal growth is the cause for green resemblance in the water body. The green color is due to the presence of chlorophyll in the algae for the photosynthesis process. Hence, the green color resembled in growth medium ensured the algal growth. The color difference had been visible during the growth period from white to yellowish green. This ensured algal growth.

3.3 Weight Increase

The dry biomass increased from 9.9 g to 14.53 g, demonstrating significant growth using the extracted nutrients.

Particulars	Initial Weight (g)	Final Dry Weight (g)
Collected Algae	71.56	9.9
Cultured Algae	78.19	14.53

Table 4.6 Weight Analysis of Algal Biomass





www.ijirset.com |A Monthly, Peer Reviewed & Refereed Journal| e-ISSN: 2319-8753| p-ISSN: 2347-6710|

Volume 14, Issue 5, May 2025

|DOI: 10.15680/IJIRSET.2025.1405211|

These findings align with previous research showing that neem extracts aid n nitrification inhibition and banana peels serve as rich potassium sources for fertilizer production (Gajalakshmi et al., 2004; Katumbwe et al., 2020)





IV.CONCLUSION

The experiment effectively demonstrated the use of ethanol – extracted nutrients from plant waste to culture algae, resulting in a functional biofertilizer. this method supports sustainable agriculture by reducing reliance on chemical fertilizers and reusing organic waste. The produced biofertilizer is rich in essential nutrients and environmentally safe.

V. FUTURE SCOPE

- Testing biofertilizer quality and nutrient composition
- Applying biofertilizer to crops and measuring growth improvement
- Scaling the process for commercial applications

REFERENCES

1.Ammar, E. E., Aioub, A. A. A., Elesawy, A. E., Karkour, A.M., Mouhamed, M. S., Amer, A. A., & EI-Shershaby, N.A.(2022). Algae as biofertilizers: between current situation and future prospective. Saudi Journal of Biological Sciences.

2.Chatterjee, A., Singh, S., Agarwal, C., Yadav, S., Rai, R., & Rai, L. C. (2017). Role Of Algae As A Biofertilizer. In Algal Green Chemistry. (Ammar, 2022)

3.Gajalakshmi, S., &Abbasi, S.A. (2004). Neem Leaves as a Source of Fertilizer Cum Pesticide Vermicompost. Bioresource Technology, 92(3), 291-296

4.Godlewska.k., Michalak. I., & Ronga. D. (2021). Plant extracts in sustainable agriculture. Italian journal of agronomy, 16(2), 151-160.

5.Kattumbwe.N. F., Muvula. N. E., seya. K. E., Bafuanyembaka. B. I., Franck. K., Longwa . N. L., & Mukalay. J. B. (2020). Synthesis of a potassium fertilizer from banana peels and its fetility effect on onion growth and ripening in lumbumbashi. International journal of recent engineering research and development.

6.Sriraj. P., Toomsan. B., &Butnan.S. (2022). Effects of Neem Leaf Extrcton The Soil properties, Growth, Yield, And Inorganic Nitrogen Contents of Lettuce. Horticulturae, 8(11),1023. (kattubwe, 2020) (sriraj. P., 2022)









INTERNATIONAL JOURNAL OF INNOVATIVE RESEARCH

IN SCIENCE | ENGINEERING | TECHNOLOGY





www.ijirset.com