

**International Journal of Innovative Research in Science,
Engineering and Technology**

(An ISO 3297: 2007 Certified Organization)

Vol. 2, Issue 9, September 2013

Jarratt Second Order Method System of Nonlinear Equations.

Sandeep Singh

Dept. of Mathematics, D. A. V. College, Chandigarh, India

ABSTRACT: In this paper, we present second-order Jarratt method for computing zeros of system of nonlinear equations numerically. In this paper, we extending the idea of the proposed families of Jarratt method to system of nonlinear equations. It is proved that the above said families have second order of convergence. Several numerical examples are also given to illustrate the efficiency and the performance of the presented families.

Keywords: System of Nonlinear equations, Optimal Order of Convergence, Halley's method, Schroder's method, Jarratt method.

I. INTRODUCTION

Due to the fact that systems of nonlinear equations arise frequently in science and engineering they have attracted researcher's interest. For example, nonlinear systems of equations, after the necessary processing step of implicit discretization, are solved by finding the solutions of systems of equations. We consider here the problem of finding a real zero, $x^* = (x^*_1, x^*_2, \dots, x^*_n)^T$, of a system of non linear equations

$$\begin{aligned} f_1(x_1, x_2, \dots, x_n) &= 0; \\ f_2(x_1, x_2, \dots, x_n) &= 0; \\ &\vdots \\ &\vdots \\ f_n(x_1, x_2, \dots, x_n) &= 0; \end{aligned}$$

This system can referred in vector form by

$$F(X) = 0 \quad (1.1)$$

Where $F = (f_1, f_2, \dots, f_n)^T$ and

$$X = (x_1, x_2, \dots, x_n)^T$$

Let the mapping $F: D \subset \mathbb{R}^n \rightarrow \mathbb{R}^n$ assumed to satisfy the assumptions (1.1). $F(X)$ is continuously differentiable in an open neighborhood D of X^* .

There exists a solution vector X^* of (1.1) in D such that $F(X^*) = 0$ and $F'(X^*) \neq 0$. Then the standard method for finding the solution to equation (1.1) is the classical Newton's method [2-5] given by $X^{k+1} = X^k - F(X^k)/F'(X^k)$, $k = 0, 1, 2, \dots$ (1.2)

$$\left\{ \begin{aligned} y_n &= x_n - \frac{2 f(x_n)}{3 f'(x_n)} \\ x_{n+1} &= x_n \\ &- \frac{1 f(x_n)}{2} \left[\frac{(a_4^2 - 22a_4 a_5 - 27a_5^2) f'(x_n) + 3(a_4^2 + 10a_4 a_5 + 5a_5^2) f'(y_n)}{(a_4 f'(x_n) + 3a_5 f'(y_n))(3(a_4 + a_5) f'(y_n) - (a_4 + 5a_5) f'(x_n))} \right] \end{aligned} \right. \quad (1.1)$$

[10] Now extend this idea for system of equations, we have

$$\left\{ \begin{aligned} Y_n &= X_n - \frac{2 F(X_n)}{3 F'(X_n)} \\ X_{n+1} &= X_n \\ &- \frac{1}{2} \left[\frac{(a_4^2 - 22a_4 a_5 - 27a_5^2) F'(X_n) + 3(a_4^2 + 10a_4 a_5 + 5a_5^2) (F(Y_n))'}{(a_4 F'(X_n) + 3a_5 F'(Y_n)) (3(a_4 + a_5) F'(Y_n) - (a_4 + 5a_5) F'(X_n))} \right] F(X_n) \end{aligned} \right. \quad (1.2)$$

International Journal of Innovative Research in Science, Engineering and Technology

(An ISO 3297: 2007 Certified Organization)

Vol. 2, Issue 9, September 2013

II COVNERGENCE ANALYSES

We shall present the mathematical proof for the order of convergence of formula (1.2).

Lemma 1: Let $D \subseteq R^n \rightarrow R^n$ be P -times Frechet differentiable in a convex set $D \subseteq R^n$ then for any $X, H \in R^n$, the following expression holds:

$$F(X+H) = F(X) + F'(X)H + 1/2! F''(X)H^2 + 1/3! F'''(X)H^3 + \dots + 1/(p-1)!F^{p-1}(X)H^{p-1} + R_p. \quad (2.1)$$

Where

$$\|R_p\| \leq 1/p! \sup_{0 \leq t \leq 1} \|F^{(p)}(X + tH)\| \|H\|^p,$$

and $H_p = (h, h, \dots, h)$.

Now we analyze the behavior of (1.2) through the following theorem:

Theorem 2.1. Let $D \subseteq R^n \rightarrow R^n$ be four times Frechet differentiable in a convex set D containing the root r of $F(x) = 0$. Then, the sequence $x_k, k \geq 0 (x^0 \in D)$ obtained by using the iterative expression of method (1.2) converges to r with convergence order second if $a_4 \neq 3a_5$ or $a_4 \neq -3a_5$.

Proof: The Taylor's expansion (3.1) for $F(x)$ about x^k is

$$F(x) = F(x^k) + F'(x^k)(x-x^k) + 1/2! F''(x^k)(x-x^k)^2 + 1/3! F'''(x^k)(x-x^k)^3 + 1/4! F^{iv}(x^k)(x-x^k)^4 + O(\|x-x^k\|^5). \quad (2.2)$$

Let $e^k = x^k - r$. Then, setting $x = r$ and using $F(r) = 0$ in (2.2), we obtain

$$F(x^k) = F'(x^k)e^k - 1/2! F''(x^k)(e^k)^2 + 1/3! F'''(x^k)(e^k)^3 - 1/4! F^{iv}(x^k)(e^k)^4 + O(\|e^k\|^5). \quad (2.3)$$

Pre-multiplying by $F'(x^k)^{-1}$ to both sides of (2.3)

$$F'(x^k)^{-1}F(x^k) = e^k - 1/2! F'(x^k)^{-1}F''(x^k)(e^k)^2 + 1/3! F'(x^k)^{-1}F'''(x^k)(e^k)^3 - 1/4! F'(x^k)^{-1}F^{iv}(x^k)(e^k)^4 + O(\|e^k\|^5). \quad (2.4)$$

Now from (1.2), we yields

$$y^k - x^k = -e^k + 1/2! F'(x^k)^{-1}F''(x^k)(e^k)^2 - 1/6 F'(x^k)^{-1}F'''(x^k)(e^k)^3 + O(\|e^k\|^4). \quad (2.5)$$

Also,

$$(y^k - x^k)^2 = (e^k)^2 - F'(x^k)^{-1}F''(x^k)(e^k)^3 + O(\|e^k\|^4), \quad (2.6)$$

$$(y^k - x^k)^3 = -(e^k)^3 + O(\|e^k\|^4), \quad (2.7)$$

$$(y^k - x^k)^4 = O(\|e^k\|^4). \quad (2.8)$$

Here $e^m = (e, e, \dots, e)$, $e \in R^n$.

Taylor's expansion of $F(y^k)$ about x^k is.

$$F(y^k) = F(x^k) + F'(x^k)(y^k - x^k) + 1/2! F''(x^k)(y^k - x^k)^2 + 1/3! F'''(x^k)(y^k - x^k)^3 + O(\|y^k - x^k\|^4). \quad (2.9)$$

From (2.5)-(2.8) and (2.9), we obtain

$$F(y^k) = F(x^k) - F'(x^k)e^k + F''(x^k)(e^k)^2 - [1/3F'''(x^k) + 1/2F''(x^k)F'(x^k)^{-1}F''(x^k)(e^k)^3] + O(\|e^k\|^4). \quad (2.10)$$

Now from (1.2)

$$e^{k+1} = e^k - \frac{1}{2} \left[\frac{(a_4^2 - 22a_4a_5 - 27a_5^2)F'(X_n) + 3(a_4^2 + 10a_4a_5 + 5a_5^2)(F(Y_n))'}{(a_4F'(X_n) + 3a_5F'(Y_n)) (3(a_4 + a_5)F'(Y_n) - (a_4 + 5a_5)F'(X_n))} \right] F(X_n)$$

$$\text{Let } R(x^k) = [(a_4^2 - 22a_4a_5 - b_5^2)F'(X^k) + 3(a_4^2 + 10a_4a_5 + 5a_5^2)F'(Y^k)] F(X^k)$$

$$\text{and } w(x^k) = (a_4F'(X^k) + 3a_5F'(Y^k)) (3(a_4 + a_5)F'(Y^k) - (a_4 + 5a_5)F'(X^k))$$

$$R(x^k) = (4a_4^2 + 8a_4a_5 - 12a_5^2)(F'(x^k))^2 e^k - 2 [(a_4^2 + 10a_4a_5 + 5a_5^2) F''(x^k) F'(x^k) + (a_4^2 + 2a_4a_5 - 3a_5^2) F''(x^k) F''(x^k)] (e^k)^2 + O(\|e^k\|^3) \quad (2.11)$$

$$w(x^k) = (a_4^2 + 4a_4a_5 - 6a_5^2)(F'(x^k))^2 - 2 [(a_4^2 + 4a_4a_5 + 3a_5^2) F''(x^k) F''(x^k) - (2a_4 - a_5)a_5 F''(x^k)F'(x^k)] (e^k) + O(\|e^k\|^2). \quad (2.12)$$

From (2.11) and (2.12), we obtain

$$w(x^k)e^k - R(x^k) = -2 [(a_4 - 3a_5)^2 F''(x^k) F''(x^k) - (a_4 + 3a_5)^2 F''(x^k) F'(x^k)] (e^k)^2 + O(\|e^k\|^3). \quad (2.13)$$

From (1.2) and (2.13), we have

$$w(x^k)e^{k+1} = w(x^k)e^k - R(x^k) = -2 [(a_4 - 3a_5)^2 F''(x^k) F''(x^k) - (a_4 + 3a_5)^2 F''(x^k) F'(x^k)] (e^k)^2 + O(\|e^k\|^3). \quad (2.14)$$

International Journal of Innovative Research in Science, Engineering and Technology

(An ISO 3297: 2007 Certified Organization)

Vol. 2, Issue 9, September 2013

From the above equation it is clear that (1.2) is second order convergence if $a_4 \neq 3a_5$ or $a_4 \neq -3a_5$.
Special cases of formula (1.2):

(a) For $a_4 = 5$ and $a_5 = 1$, family (1.2) read as:

$$\left\{ \begin{array}{l} y_n = x_n - \frac{2f(x_n)}{3f'(x_n)} \\ x_{n+1} = x_n \\ - \left[\frac{4[7F'(X_n) - 15F'(Y_n)]}{(5F'(X_n) - 9F'(Y_n))(5F'(X_n) + 3F'(Y_n))} \right] F(X_n) \end{array} \right. \quad (2.15)$$

This is a new second-order method and satisfies the following error equation $w(x^k) e^{k+1}$

$$= -2 [4 F'(x^k) F''(x^k) + 64 F''(x^k) F'(x^k)] (e^k)^2 + O(\|e^k\|^3).$$

(b) For $a_4 = 1$ and $a_5 = -2$, family (1.2) read as:

$$\left\{ \begin{array}{l} y_n = x_n - \frac{2f(x_n)}{3f'(x_n)} \\ x_{n+1} = x_n \\ - \frac{1}{2} \left[\frac{[-21F'(X_n) + F'(Y_n)]}{(F'(X_n) - 6F'(Y_n))(9F'(X_n) - 3F'(Y_n))} \right] F(X_n) \end{array} \right. \quad (2.16)$$

This is a new second-order method and satisfies the following error equation $w(x^k) e^{k+1}$

$$= -2 [49 F'(x^k) F''(x^k) - 25 F''(x^k) F'(x^k)] (e^k)^2 + O(\|e^k\|^5).$$

III. NUMERICAL RESULTS

In this section, we shall check the performance of the present formula JS1(2:15) and JS2(2:16) the comparison is carried out with Newton's method and with HM and CM [8]. A mat lab program has been written to implement these methods. We use the following stopping criteria for computer programs:

1. $\epsilon = e^{-10}$.
2. $|F(X_n)| < \epsilon$

For every method, we analyze the number of iterations needed to converge to the required solution. The numerical results are reported in the Table 1.

We consider the following problems for a system of nonlinear equations.

Problem (a)

$$\begin{aligned} x_1^2 - 2x_1 - x_2 + 0.5 &= 0 \\ x_1^2 + 4x_2^2 - 4 &= 0 \end{aligned}$$

Problem (b)

$$\begin{aligned} x_1^2 + x_2^2 - 1 &= 0 \\ x_1^2 - x_2^2 + 0.5 &= 0 \end{aligned}$$

Problem (c)

$$\begin{aligned} x_2 x_3 + x_4(x_2 + x_3) &= 0 \\ x_1 x_3 + x_4(x_1 + x_3) &= 0 \\ x_2 x_1 + x_4(x_2 + x_1) &= 0 \\ x_1 x_2 + x_1 x_3 + x_2 x_3 - 1 &= 0 \end{aligned}$$

Problem (d)

$$\begin{aligned} e^{x_1} + x_1 x_2 - x_2 - 0.5 &= 0 \\ \sin(x_1 x_2) + x_1 + x_2 - 1 &= 0 \end{aligned}$$

**International Journal of Innovative Research in Science,
Engineering and Technology**

(An ISO 3297: 2007 Certified Organization)

Vol. 2, Issue 9, September 2013

Problem (e)

$$\begin{aligned} x_1 + 2x_2 - 3 &= 0 \\ 2x_1^2 + x_2^2 - 5 &= 0 \end{aligned}$$

Problem (f)

$$\begin{aligned} x_1 + e^{x_2} - \cos(x_2) &= 0 \\ 3x_1 - x_2 - \sin(x_2) &= 0 \end{aligned}$$

Problem (g)

$$\begin{aligned} x_1^2 + x_2^2 + x_3^2 - 9 &= 0 \\ x_1 x_2 x_3 - 1 &= 0 \\ x_1 + x_2 - x_3^2 &= 0 \end{aligned}$$

Solution (a)

$$r = (1.9006767263670658, 0.31121856541929427)^T$$

Solution (b)

$$\begin{aligned} r &= (0.5000000000000000, 0.8660254378443865)^T \\ s &= (-0.5000000000000000, -0.8660254378443865)^T \end{aligned}$$

Solution (c)

$$\begin{aligned} r &= (0.57735026918962576, 0.57735026918962576, 0.57735026918962576, -0.2886751459481288)^T \\ s &= (0.57735026918962576, 0.57735026918962576, 0.57735026918962576, -0.2886751459481288)^T \end{aligned}$$

Solution (d) $r = (0, 1)^T$

Solution (e)

$$r = (1.4880338717125849, 0.75598306414370757)^T$$

Solution (f)

$$r = (0, 0)^T$$

Solution (g)

$$\begin{aligned} r &= (2.2242448288477843, 0.28388497407293814, 1.58370776128252723)^T \\ s &= (0.28388497407293814, 2.2242448288477843, 1.58370776128252723)^T \end{aligned}$$

TABLE 1: Numerical results of problems (a) to (g) using different methods.

F(X)	X	NM	HM	CM	JS1	JS2
(a)	$(3,2)^T$	10	7	6	10	4
	$(1.6, 0)^T$	9	7	6	6	2
(b)	$(.7, 1.3)^T$	9	6	5	3	3
	$(-1, -2)^T$	9	6	5	3	3
(c)	$(-1, -1, -1, -1)^T$	9	6	5	5	3
	$(1, 1, 1, 7)^T$	9	6	5	5	5
(d)	$(.7, .9)^T$	9	6	5	6	4
	$(-0.1, 0.2)^T$	9	6	6	3	3
(e)	$(.99, .1)^T$	9	7	6	5	4
	$(1.9, 1.4)^T$	9	7	5	9	4
(f)	$(1.5, 2)^T$	10	7	6	div	4
	$(.3, .5)^T$	9	6	5	5	3
(g)	$(.2, .6, 1.5)^T$	9	6	5	5	16
	$(3, .5, 2)^T$	9	6	5	3	3

International Journal of Innovative Research in Science, Engineering and Technology

(An ISO 3297: 2007 Certified Organization)

Vol. 2, Issue 9, September 2013

IV. CONCLUSIONS

The presented formula (2.15) and (2.16) is simple to understand, easy to program and has the second order of convergence. We contribute to the development of iteration processes and propose of Jarratt's method. We now obtain a wide general class of Jarratt's methods which are without memory and have the same scaling factor of function as that Jarratt's method. Numerical tests have been performed, which not only illustrate the method practically but also serve to check the validity of theoretical results we have derived. The performance is compared with Newton method, CM [8] and HM [8].

REFERNCES

- [1] A.M., Ostrowski, (1960). Solutions of Equations and System of Equations, Academic Press, New York.
- [2] Ostrowski, A. M., (1966). Solution of equations and system of equations. Academic Press NY, London.
- [3] Ortega, J. M. and Rheinboldt, W. C.,(1970). Iterative solution of non-linear equations in several variables, Academic Press, Inc.
- [4] Traub, J. F., (1982). Iterative method for the solution of equations, Chelsea Publishing Company, New York.
- [5] Dennis Jr, J. E. and Schnabel, R. B., (1983). Numerical methods for unstrained optimization and nonlinear equations, Prentice Hall, Englewood Cliffs, NJ.
- [6] J.F.Traub,(1964),Iterative Methods for the Solution of Equations, Prentice-Hall, Englewood Cliffs, NJ.
- [7] H.T. Kung,J.F. Traub,(1974), Optimal order of one-point and multipoint iteration, Journal of the ACM 21, pp.643-651.
- [8] Sharma, R.k. Guha and J. R. , R. (2012), An efficient fourth order weighted-Newton method for systems of nonlinear equations, Springer science +Business Media, LLC.
- [9] E.Schroder's, (1870), Uber unendlichviele Algorithm zur Auffosung der Gleichungen, Ann. Math.,2,pp.317-365.
- [10] R. Bhel, V. Kanwar and K. Sharma,(2012), optimal equi-scaled families of Jarratt's method, International Journal of Computer Mathematics.